

Docking of GSK-3 β with novel inhibitors, a target protein involved in Alzheimer's disease

Akanksha Joshi, Archit Sharma and Rajesh Kumar*

Department of Biotechnology, University Institute of Engineering and Technology, Kurukshetra University, Kurukshetra, Haryana, India

ABSTRACT

Alzheimer's disease (AD) is a chronic, advancing malady associated with loss of memory or cognition. It is the noted causes of lethality worldwide, there are no such drugs which can cure AD till date and are ineffective in the later stages. Such known drugs only ease the symptoms but do not prevent the onset or progression of the AD. Alzheimer's is caused by the aggregation of the hyperphosphorylated tau which is one of the common characteristics of the neurodegenerative disorder. There are a number of kinases which hosts the excessive phosphorylation of tau protein. One of the kinase extensively targeted in the AD is GSK-3 β (Glycogen Synthase Kinase-3 β). As indicated by many studies that by applying appropriate docking methods, a number of phyto compounds have shown enhanced target selectivity than the conventional Alzheimer's drugs. This review summarizes the known drug targets in the AD, their conventional inhibitors and also the comparison between the current and future AD therapy based on their binding affinities. As a result, large libraries of compounds with inhibitory effect can be screened. It was also studied that Withanolide-A has the potential to be the future drug for Alzheimer's disease.

KEY WORDS: DOCKING; DRUGS; PHOSPHORYLATION; TAU PROTEIN; WITHANOLIDE-A

INTRODUCTION

Alzheimer's is a type of dementia associated with memory loss and other intellectual abilities, severe enough to intrude with regular routine. Alzheimer's disease report for 60 to 80 percent of dementia and the present Alz-

heimer's disease therapies impaired from in proficient effects on its symptoms such as perception notably in the subsequent stages of the disease (<http://www.alz.org>). According to the report prepared by Alzheimer's and related disorders society of India in 2010, there are 3.7 million Indians suffering with dementia while the

ARTICLE INFORMATION:


*Corresponding Author: rkumar2015@kuk.ac.in

Received 16th March, 2018

Accepted after revision 18th June, 2018

BBRC Print ISSN: 0974-6455

Online ISSN: 2321-4007 CODEN: USA BBRCBA

 Thomson Reuters ISI ESC / Clarivate Analytics USA and Crossref Indexed Journal

NAAS Journal Score 2018: 4.31 SJIF 2017: 4.196

© A Society of Science and Nature Publication, Bhopal India 2018. All rights reserved.

Online Contents Available at: <http://www.bbrc.in/>

DOI: 10.21786/bbrc/11.1/13

numbers are anticipated to bifold by 2030. The number of factors is thought to increase the progression of this disease, some of which are; increasing age, family history, previous severe head injuries etc. Over the past decade, much of the research on Alzheimer disease (AD) has focused on radical-effected oxidative stress and its importance in disease pathogenesis. Oxidative stress increases amyloid beta deposits in the brain which results in the synthesis of neurotoxic aggregates. The net effect of oxygen radicals is damaging as it may lead to neuronal cell death and contribute to AD (Smith et al. 1998). Flavonoids also possess antioxidant activity and they regulate the redox status and prevent damage caused by oxidative stress. Protein Kinases are recognised as encouraging target structures considering their involvement in AD breakthrough pathways like pathophysiological tau protein phosphorylation and amyloid beta toxicity. The sound interdependence of tau phosphorylation and pathology has led to the search for Tau protein kinase inhibitors such as GSK3- β and Tyrosine kinase Fyn, which phosphorylates tau and also plays a causative role in amyloid pathway. Hereafter, acting as potential therapeutic agents (Medina, 2018).

ROLE OF FLAVONOIDS IN THE TREATMENT OF AD

Nature has fascinated us with a lot of natural remedies in the form of fruits, leaves, bark, vegetables, and nuts, etc. The wide varieties of biologically active nutrients existing in these natural products play a vital role in defence and aid of various neurodegenerative diseases. Flavonoids are an array of non-nutrient polyphenolic compounds readily procured from plants. It was realized that the competence of flavonoids to upgrade neurological health was resolved by their antioxidant capability. Flavonoids are endowed with numerous biological activities like anti-inflammatory, anticoagulant, anti-cancer, anti-oxidants, and anti-spasmodic. There is an extensive role of flavonoids and even their metabolites in different signaling pathways by altering the phosphorylation state of target protein put forward their therapeutic potential and beneficial in neurodegeneration (Spencer, 2007). Increasing evidence shows their ability to improve brain function such as memory and learning by interacting with cellular as well as molecular components of the brain resulting in enhanced neuronal function and induce neurogenesis (Spencer, 2010; Baptista et al. 2014).

A study has found the role of plant-derived compounds such as myricetin and epicatechin-5-gallate in abrogating heparin-induced cluster of tau into filaments (Taniguchi et al. 2004). In drug discovery, the dominant secondary metabolites (terpenoids, phenolics, and alkaloids) are of probable remedial relevance. Certain fla-

vonoids such as indirubin and morin are capable of the inhibiting the activity of GSK-3 beta and thereby blocking tau hyperphosphorylation. Kinases are involved in tau phosphorylation and phosphatases reverse this action. Thus, flavonoids also portray a crucial aspect in modulating the activity of phosphatases (Baptista et al. 2014). Genistein (phytoestrogen), a beneficial intermediary for the treatment of AD as it imitates estrogen which is involved in the development of memory and learning along with its neuroprotective activities, (Hussain et al. 2018). It was found that eicosanoyl-5-hydroxytryptamide (EHT), a naturally occurring component of coffee beans accelerates the activity of serine/threonine protein phosphatase, PP2A and thus provide therapeutic benefits associated with AD (Asam et al. 2017).

MOLECULAR CAUSES OF AD

The key events that lead to AD : Beta-amyloid toxicity. The brain of a patient with the AD is characterized by amyloid toxicity. Amyloid beta denotes peptides of 36-43 amino acids long processed from an amyloid precursor protein (APP) which is digested by beta secretase and gamma secretase to yield amyloid beta (A β). This peptide is found in brains of patients suffering from Alzheimer's (Murphy et al. 2010 Hamley, 2012, Sauer, 2017). Some processes include disruption of amyloid beta aggregates, alterations in the precursor of amyloid beta protein processing through the inhibition of beta-secretase. Thus, modulating the beta-secretase activity is the one suggested a therapeutic avenue to treat AD (Yin et al. 2007). Certain flavonoids may guard to counter the effect of Alzheimer's disease by interrupting with the generation of beta-amyloid peptides into neurotoxic aggregates. It is a matter of contention that interfering with the activity of beta and gamma-secretase enzymes may disrupt their other functional roles besides playing an important part in amyloidogenic pathways.

Thus such interference using γ secretase can result in skin cancers and cognitive dysfunction (Kikuchi et al. 2017). The decades old theory which aims at implicating beta amyloid as the leading cause of Alzheimer's has been questioned by a group of scientists. Researchers have tried and failed to prevent Alzheimer's using drugs targeted at amyloid β protein. Due to the lack of the utility of amyloid- β -aspired approach in Phase III clinical trials, it was prerequisite to conceive substitute drug discovery strategies for alzheimer's (Folch et al. 2016). Solanezumab, a drug which acts on amyloid β protein failed some pivotal clinical trials. However, it is still anonymous whether the disease is caused by plaques or they are just the by- products (Ramsey, 2018).

A number of normal patients have been found with amyloid deposits in their brain. It was anticipated that


amyloid beta deposition is an anomaly of aging and does not correlate with the AD progression (Kametani et al. 2018). Therefore there is a compulsive need to search policies directed at reducing misfolded tau protein which is one of the disease-causing agents (Bruden et al. 2010). Tau is liable to be the more superior target than the amyloid β as it coordinates efficiently with cognitive impairment, provided clinical symptoms are tangible (Congdon et al. 2018).

TAU PROTEIN HYPERPHOSPHORYLATION

For more than a decade, researchers have found 'tau' protein as one of the causes other than the Beta-amyloid plaques (Underwood, 2016). Accordingly, tau hyperphos-

phorylation and accumulation of insoluble aggregates is strongly related to reduce cognitive performance. Hence, we can affirm that tau is a reliable marker of the neurodegenerative process (Fig.1). Incorporation of phosphate groups into tau depends on; tau's confirmation and equity amidst the activity of kinases and phosphatases (Kremer et al. 2011).

Changes in tau confirmation could lead to excessive phosphorylation resulting in the formation of neurotoxic aggregates and tau-mediated neurodegeneration (Dixit et al. 2008). Tau is the member of family of proteins intricately in stabilizing the microtubules. They are common in neurons of Central Nervous System and also present at low levels in CNS astrocytes and oligodendrocytes (Shin



et al. 1991). The tau proteins have been formed as a result of alternate splicing of MAPT (microtubule-associated protein tau) gene in humans and is positioned on chromosome 17 (Goedert et al. 1989; Jesus et al. 2016).

The hydrophilic nature of the tau protein and its existence as intrinsically disordered protein was unfolded by many biophysical studies. (Porowska et al. 2014). One of the critical function of tau protein is to prevent the depolymerization of microtubules by regulating its stability in two ways: isoforms, and phosphorylation. On the basis of the number of binding domains, six variants of tau protein (352-441 amino acids and apparent molecular weight between 60-74kDa) exist in human brain tissue (Martin et al. 2011). Out of six modifications, three isoforms have 3 tubulin binding domains and other three have 4 tubulin binding domains in the C-terminal half of tau, (Guo et al. 2017).

The domain structure of tau is such that its positively charged binding domain is located in carboxy-terminal which binds to the microtubule which is negatively charged. Tau is a phosphoprotein (i.e. posttranslationally modified) with 79 probable Serine (Ser) and Threonine (Thr) phosphorylation sites on the extended tau isoform. It has been reported in a study that phosphorylation is possible in about 30 sites in a normal tau protein. PKN, a serine/threonine kinase is one such enzyme among the plethora of kinases which regulates the phosphorylation of tau (Billingsley et al. 1997). As revealed by primary sequence analysis, the tau molecule has three major domains: N-terminal (acidic), a proline-rich region, C-terminal domain (basic). These domains are characterized on the basis of their amino acid character and even on their microtubule interactions. Thus, tau protein acts as a dipole with two domains having the opposite charge (Kolarova et al. 2012; Porowska et al. 2014). Extreme phosphorylation of the tau protein proceeds to the formation of Paired helical fragments (PHF's) due to the loss of affinity with microtubules and they bind with one another which further aggregates in neurofibrillary tangles via. Post-translational modifications. Thus, there is a strong correlation between abnormal phosphorylation and self-aggregation of tau (Guo et al. 2017)

When disorganized, this aside from being very soluble protein forms remarkably insoluble tangles or aggregates which commit to the number of neurodegenerative disorders. The mutations in posttranslational modifications are the main cause of this failure i.e. they form non-functional aggregates. One of the studies demonstrated that dephosphorylation of the hyperphosphorylated tau converts abnormal tau protein into a normal like protein which then regulates microtubule assembly (Iqbal et al. 2011). Therefore abrogating the abnormal tau and recovery of the microtubule organization are the most promising therapeutic interventions to combat AD.

GSK-3 BETA AS A DRUG TARGET

GSK-3 is encoded by two genes: GSK-3 β , located on chromosome 19 and GSK-3 α , positioned on chromosome 2. GSK-3 is ubiquitously expressed in mammals as well as in yeast (Medina et al. 2011). GSK3 mediates the augmentation of phosphate molecules to serine and threonine amino acid residues and for this reason termed as serine/threonine protein kinase. The kinase domain of these 2 isoforms are highly homologous ((Stambolic et al. 1994) but are demarcated in the N- and C-terminal regions. GSK3 β has a molecular mass of 46-47 kDa consisting of 433 and 420 amino acids in human and mouse respectively. The protein contains an N-terminal domain, a kinase domain, and a C-terminal domain. The substrate Binding domain (BD) provides GSK-3 β specific binding sites for the tumor suppressor p53 and other protein complexes (Atlas of Genetics and Cytogenetics in Oncology and Haematology). A number of protein kinases are involved in tau phosphorylation such as Cdk5 (Cyclin-dependent Kinase 5), JNK (C-Jun amino-terminal Kinase), CK1 (Casein Kinase1), Dyrk1A, AMPK (Adenosine-monophosphate activated protein kinase), MARK5 (Microtubule affinity-regulating Kinases), PKA (Cyclic AMP-dependent protein Kinase), GSK-3 β (Glycogen Synthase Kinase-3 β) (Crews et al. 2010). But a study has shown that 31% of the therapeutically favorable phosphorylation sites of tau protein are phosphorylated by GSK3 β (Martin et al. 2013).

The classical approach to treat misfolding of tau protein provides inhibition of protein kinases (Glycogen synthase kinase 3 β) which hosts tau phosphorylation. According to the 'GSK-3 hypothesis of AD', tau hyperphosphorylation, memory impairment and enhanced β -amyloid production is due to the overexpression of GSK-3, all of which are characteristic features of the AD. If this hypothesis is consolidated then, inhibition of GSK-3 β by novel inhibitors provides a better pathway against the effect of this destructing disorder (Hooper et al. 2008). There are two isoforms of GSK-3 gene; GSK-3 α and GSK-3 β . GSK3 β also exist as longer splice variants (Mukai et al. 2002; Schaffer et al. 2003). Moreover, GSK-3 β results in a neuronal decline in the AD because of the fact that it is a causal mediator of apoptosis. Increased level of such protein eventuated in the autopsy evaluation of brain of alzheimer's victims (Pei et al. 1997). It is also validated that a spatial and temporal pattern of enhanced GSK-3 expression corresponds with the evolvment of neurofibrillary tangles proceeding towards neurodegeneration (Leroy et al. 2002).

MOLECULAR DOCKING

Drug research is an important tool in the field of medicine. Utility of computers to anticipate the efficiency

of binding of a set of small molecules or ligands with the target is an important element of drug discovery and developmental process. There is an ample realm of software packages used to execute molecular docking such as Dock, Autodock, GOLD, ICM, Glide, AutoDock Vina, FlexX etc. Automated docking is generally used for prognosis of biomolecular complexes, in structure and function examination and in computer-aided drug designing. A dozen of mechanism is available, consolidating varied energy evaluation methods. Due to the enhanced docking speed, AutoDock 4.2 has been widely used for virtual screening. It is the ultimate current version which is based upon the Lamarckian genetic algorithm, a hybrid algorithm comprising of both the genetic as well as local search and is more enhanced and accurate than previous version AD3.0. Unlike AD3.0, Autodock 4.2.6 (henceforth AD4.2) and Auto Dock Vina 1.1.2 (henceforth AD Vina) have upgraded results and improved elucidation, (Collignon et al. 2011, Nataraj et al. 2017 and Alvarez et al. 2017).

Two main programs are involved in AutodockTools: Autodock for docking of the ligand within the set of grids (within the binding site) in the target protein and Autogrid for selection of grid parameters, size of the box, its location etc (<http://autodock.scripps.edu/>). It is particularly suitable for protein-ligand docking in which we presume the pose and orientation of a small molecule when it is articulated to a protein receptor. It is used to select likely drug candidates. Typically, ligands are drug candidates and the macromolecule is the protein or receptor of the known three-dimensional structure. In this docking simulation, the ligand being docked was kept as flexible while target protein was kept as rigid. The graphical user interface i.e. Autodock Tools was used to prepare, run, analyzes the docking simulations.

CURRENT AND FUTURE AD THERAPY

Till date there are no such drugs/treatments available that can cure AD completely. However, there are several medications developed for Alzheimer's disease that can temporarily attenuate the symptoms. The Food and Drug Administration (FDA), U.S. has affirmed two medications-acetylcholinesterase inhibitors and Memantine. Drugs such as tacrine, rivastigmine, galantamine, and donepezil are the widely used conventional drugs to treat AD (Islam et al. 2013). Memantine is a dissociative hallucinogenic and anesthetic drug of the adamantane class of chemicals that are currently used as an FDA approved drug in the treatment of AD (www.alz.org). Therefore, traditional drugs like memantine and donepezil are being extendedly used as the reference in molecular docking studies. Hence, the objective of eventual AD therapy is to discover such novel compounds

which can target the tau protein and so that can be utilized for the recovery of neurodegenerative loss (Schneide et al. 2008).

The study related to the AD is focused more towards the traditional medicinal plants and its components such as *Withania somnifera* (Ashwagandha), *Celastrus paniculatus* (Jyotismati), *Convolvulus pluricaulis* (Shankhpushpi), *Bacopa monnieri* (Brahmi). By analyzing the binding energies of various ligands such as acacatechin, catechin, galangin, scopoletin, silibinin, memantine (as standard), it was observed that flavonoids exhibit binding energy scaled between 7.07 kcal/mol to -4.85 kcal/mol. Silibinin demonstrate prominent binding energy -7.07 kcal/mol than the standard memantine (-5.89 kcal/mol) (Madeswaran et al. 2013). A phytocompound, Catechin (with binding energy -9.7 kcal/mol) was shown to be the potent target of GSK-3 β and showed the same drug-likeness as conventional drug Donepezil (with binding energy-8.9kcal/mol), (Alam et al. 2017).

Withania somnifera, a potential inhibitor of GSK-3 β

Withania somnifera commonly called Ashwagandha, Indian ginseng and wind cherry have been recognizes as an important herb in Indigenous and ayurvedic medical system. Historically, the plant has been used therapeutically for boosting the brain function including memory retrieval. It has a cognition promoting effect in adults and children (Singh et al. 2011). It consists of two components: withanolides and withanamides. Withanolide A is extracted from the roots of the plant and promotes antioxidant properties that protect nerve cells from harmful free radicals. Many clinical trials and excessive research on animals support the use of Ashwagandha for anxiety, cognitive and neurological disorders (Rajasekar et al. 2011). Withanolides have also been used for the treatment of AD (Khan et al. 2016). Withanolide A is used as an inhibitor of acetylcholinesterase activity and reduces beta-amyloid protein formation. Also, it has been involved in the regeneration of pre and postsynaptic neurons. Instead of the root extract, a study also suggested fruits and leaves of Egyptian plant have strong antioxidant activity (Mahrous et al. 2017)

FUTURE PERSPECTIVES

Several new therapeutic approaches are currently under investigation which aims at targeting proteins such as Apolipoprotein E which is also responsible for the accumulation and hyperphosphorylation of tau. Anti-tau immunotherapeutic agents have gained much focus due to their specificity and selectivity to combat AD. But a longer follow up period might be required to test the safety and efficacy as the results were promising. Moreover targeting either tau or amyloid beta individually is

not apparently the satisfactory approach and therefore, combinational therapies might be thought of as a new proposal, (Coman et al. 2017 Bittar et al. 2018).

CONCLUSION

Drug research is of utmost importance in the field of medicine. Consequently, the use of computers to foresee the efficiency of binding of a set of molecules or ligands with the target is an important element of drug development process. To explore potent and effective drugs for the treatment of AD, different phytochemicals were compared against the standard using Autodock4. Appropriate ligands were docked into the active site of the receptor GSK-3 β and analyzed for the effective protein-ligand interactions. Therefore molecular docking identified many more promising, efficacious, selective new drugs against Alzheimer's reducing the time span of complex drug discovery process. Appropriate experimental evidences such as ADMET analysis which testifies absorption, penetration and toxicity may also be considered further as a lead in drug discovery process.

ACKNOWLEDGEMENTS

The authors are thankful to the Department of Biotechnology, University Institute of Engineering and Technology, Kurukshetra (U.I.E.T) for providing the space and resources for this work.

CONFLICT OF INTEREST STATEMENT

Authors state no conflict of interest. All authors have read the journal's Publication ethics and publication malpractice statement available at the journal's website and hereby confirm that they comply with all its parts applicable to the present scientific work.

REFERENCES

Alam A, Tamkeen N, Imam N, Farooqui A, Ahmed M.M. Ali S. Malik M.Z. and Ishrat R (2017) Pharmacokinetics and Molecular Docking studies of Plant-Derived Natural Compounds to Exploring Potential Anti-Alzheimer Activity Available from: <https://arxiv.org/pdf/1709.10374>

Alvarez A.C., Costa A.M. and Vilarrasa J (2017) The Performance of Several Docking Programs at Reproducing Protein-Macrolide-Like Crystal Structures. *Molecules* 22(1), 36 DOI:10.3390/molecules22010136

Alzheimer's disease and tauopathies.npj Vaccines volume 3, Article number: 9 DOI:

Anja S, Eckhard M. (2008) Tau-Based Treatment Strategies in Neurodegenerative Diseases Neurotherapeutics. *The Journal of the American Society for Experimental NeuroTherapeutics*

[cited 2008 July]; 5: 443-457. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5084246> DOI: 10.1016/j.nurt.2008.05.006

Anna K, Justin V L, Tomasz J, Fred V L. (2011) GSK3 and Alzheimer's disease: facts and fiction. *Frontiers in Molecular Neuroscience* [cited 2011]; 4(17):1-10. Available from: <https://www.frontiersin.org/articles/10.3389/fnmol.2011.00017/full> DOI: 10.3389/fnmol.

Anna MP, Urszula W, Marcelina G, Anna F, Grazyna N.(2014) Tau Protein Modifications and Interactions: Their Role in Function and Dysfunction. *Int. J. Mol. Sci.* [cited 2014 Mar 18]; 15: 4671-4713. Available from: <http://www.mdpi.com/1422-0067/15/3/4671ag> DOI: 10.3390/ijms 15034671

Articles /PMC5425816 DOI: 10.1007/s12551-016-0247-1 articles DOI: 10.3389/fnagi.2016.00262

Arumugam M, Muthuswamy U, Kuppusamy A, Thirumalaisamy S, Varadharajan S, Puliyath J. (2012) Docking studies; In silico lipoxygenase inhibitory activity of some commercially available flavonoids commercially available flavonoids *Advances in Tau-focused drug discovery for Alzheimer's disease and related tauopathies Ban. J. Pha.* [cited 2012 Jan 15]; 6: 133-138. Available From: <https://www.banglajol.info/Index.php/7055> DOI: 10.3329/bjp.v6i2.9408

Arumugam M, Muthuswamy U, Kuppusamy A, Thirumalaisamy S, Varadharajan S, Puliyath J. (2013) Computational drug discovery of potential TAU protein kinase I inhibitors using in Silico Docking Studies *Ban. J. Pha.* 8: 131-135. Available from: <http://www.banglajol.info>

Asam K, Staniszewski A, Zhang H, Melideo S.L., Mazzeo A, Voronkov M, Huber K.L., Pérez E, Stock M, Stock J.B., Arancio O, and . Nicholls R.E. (2017) Eicosanoyl-5-hydroxytryptamide (EHT) prevents Alzheimer's disease-related cognitive and electrophysiological impairments in mice exposed to elevated concentrations of oligomeric beta-amyloid. *PLoS One* 12(12) DOI: 10.1371/journal.pone.0189413

Avila J, Jiménez JS, Sayas CL, Zabala MBS, Rivas G, Hernández F. (2016) Tau structures. *Frontiers in Aging Neuroscience* 8(262): 1-10. Available from: <https://www.frontiersin.org/>

Baptista FI, Henriques AG, Silva AMS, Wiltfang A, Silva OAB. (2014) Flavonoids as Therapeutic Compounds Targeting Key Proteins Involved in Alzheimer's Disease. *ACS Chem. Neurosci.* 5(2): 83-92. Available from: DOI:abs/10.1021/cn400213r.2

Billingsley ML, Kincaid RL. (1997) Regulated phosphorylation and dephosphorylation of tau protein: effects on microtubule interaction, intracellular trafficking, and neurodegeneration. *Biochem J.* 323(pt3):577-91

Bittar A, Sengupta U and Kayed R (2018) Prospects for strain-specific immunotherapy in

Brunden KR, Trojanowski JQ, Le VMY (2010) Advances in Tau-focused drug discovery for Alzheimer's disease and related tauopathies. *Nat Rev Drug Discov.* [cited 2010 apr 1]; 8(10): 783-793. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC/articles/PMC2787232> DOI: 10.1038/nrd2959

Collignon B, Schulz R, Smith JC, Baudry J. (2011) Task-parallel message passing interface implementation of Autodock4 for

- docking of very large databases of compounds using high-performance super-computers . J. Comp. Chem. [cited 2011 Sep 10]; 32: 1202–1209. Available from: <https://onlinelibrary.wiley.com> DOI: 10.1002/jcc.21696
- Coman H and Nemeş B (2017) New Therapeutic Targets in Alzheimer's Disease. International Journal of Gerontology 11(1), Pages 2–6 DOI: <https://doi.org/10.1016/j.ijge.2016.07.003>
- Congdon E.E, Sigurdsson E.M. (2018) Tau-targeting therapies for Alzheimer disease. Nature Reviews Neurology
- Crews L, Masliah E. (2010) Molecular mechanisms of neurodegeneration in Alzheimer's disease Human Molecular Genetics 19(1): 12–20. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2875049> DOI: 10.1093/hmg/ddq160.
- Dixit R, Ross JL, Goldman YE, Holzbaur EL. (2008) Differential regulation of dynein and kinesin motor proteins by tau Science, Feb; 319: 1086–1089. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18202255> DOI: 10.1126/science.1152993
- Folch J, Petrov D, Ettchet M, Abad S, Lopez ES, Garcia ML, Olloquequi J, Zarate CB, Auladell C, Camins A (2016) Current Research Therapeutic Strategies for Alzheimer's Disease Treatment. Neural Plasticity [cited Aug 2016]:1–15. Available from: <http://dx.doi.org/10.1155/2016/850169312>
- Goedert M, Spillantini MG, Jakes R, Rutherford D, Crowther RA. (1989) Multiple Isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. Neuron 3(4): 519–526. Available from: DOI: 10.1016/0896-6273(89)90210-9
- Guo T, Noble W, and Hanger D.P. (2017) Roles of tau protein in health and disease. Acta Neuropathol. 133(5): 665–704 DOI: 10.1007/s00401-017-1707-9
- Hamley IW (2012) The Amyloid Beta Peptide: A Chemist's Perspective. Role in Alzheimer's and Fibrillization. Chemical Reviews 112: 5147–5192. Available from: <https://pubs.acs.org> DOI: [10.1021/cr3000994](https://doi.org/10.1021/cr3000994)
- Hooper C, Killick R, Lovestone S. (2008) The GSK3 hypothesis of Alzheimer's disease. J Neurochem. 104(6):1433–1439
- Hussain G, Zhang L, Rasul A, Anwar H, Sohail M.U. Razzaq A, Aziz N, Shabbir A, Ali M and Sun T (2018) Role of Plant-Derived Flavonoids and Their Mechanism in Attenuation of Alzheimer's and Parkinson's Diseases: An Update of Recent Data Molecules, 23(4), 814 DOI:10.3390/molecules23040814
- Kametani F, Hasegawa M. (2018) Reconsideration of Amyloid Hypothesis and Tau Hypothesis in Alzheimer's Disease Front Neurosci.12: 25. DOI: 10.3389/fnins.2018.00025
- Kazunori K, Kiwami K, Takuya T, Taisuke T.(2017) Dysregulated Metabolism of the Amyloid- β Protein and Therapeutic Approaches in Alzheimer Disease. J Cellular Biochemistry 118(12): 4183–4190
- Khalid I, Fei L, Cheng XG, Inge GI. (2011) Tau in Alzheimer Disease and Related Tauopathies. Cur. Alzheimer Res. [cited 2011 May 9]; 7(8): 656–664. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3090074>
- Khan SA, Khan SB, Shah Z, Asiri AM. (2016) Withanolides: Biologically Active Constituents in the Treatment of Alzheimer's disease. Med Chem. 12(3):238–56. Available from:<https://www.ncbi.nlm.nih.gov/pubmed/26527154> DOI:10.2174/15734064116661510
- Kolarova M, Garcia-Sierra F, Bartos A, Ricny J, Ripova D. Structure and Pathology of Tau Protein in Alzheimer Disease (2012) International Journal of Alzheimer's disease 2012: 1–13. Available from: <http://dx.doi.org/10.1155/2012/731526>
- Leroy K, Boutajangout A, Authelat M, Woodgett JR, Anderton BH, Brion JP. (2002) The active form of glycogen synthase kinase-3 β is associated with granulovacuolar degeneration in neurons in Alzheimer's disease . Acta Neuropathol. [cited 2002 feb]; 103: 91–99. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11810173> DOI: 10.1007/s004010100435
- Mahrous RS, Ghareeb DA, Fathy HM, Abu EL-Khair RM, Omar AA. (2017) The Protective Effect of Egyptian *Withania somnifera* Against Alzheimer's. Med. Aromat Plants (Los Angeles). 6(2): 1–6. Available from: <http://www.omics.org/> DOI:10.4172/2167-0412.100028.
- Mark AS, Keisuke H, Karen H, Migu AP, Peggy LRH, Sandra LS, Massimo T, George P. (1998) Amyloid- β Deposition in Alzheimer Transgenic Mice Is Associated with Oxidative Stress. J Neurochemistry 70(5): 2212–5
- Martin L, Latypova X, Terro F. (2011) Post-translational modifications of tau protein: implications for Alzheimer's disease. Neurochem. Int. 58(4):458–471. Available from:<https://www.ncbi.nlm.nih.gov/pubmed/21215781> DOI:10.1016/j.neuint.2010.12.023
- Medina M (2018) An Overview on the Clinical Development of Tau-Based Therapeutics Int. J. Mol. Sci. 19, 1160 Available from <http://www.mdpi.com/journal/ijms> DOI: 10.3390/ijms19041160
- Medina M, Wandosell F. (2011) Deconstructing GSK-3: The Fine Regulation of Its Activity. International Journal of Alzheimer's disease. Feb; 2011:1–12. Available from: <http://dx.doi.org/10.4061/2011/479249>
- Mukai F, Ishiguro K, Sano Y, Fujita SC.(2002) Alternative splicing isoform of tau protein kinase I/glycogen synthase kinase 3 beta. J. Neurochem. ; 81(5):1073–1083
- Murphy MP, LeVine H. (2010) Alzheimer's Disease and the β -Amyloid Peptide. J Alzheimers Dis. 19(1):311. Available from: <https://www.ncbi.gov/pmc/articles/pmc28135509/>DOI:
- Natraj S, Pagadala, Syed K, Tuszynski J. (2017) Software for molecular docking: a review Biophys. Rev. 9(2): 91–102. Available from: <https://www.ncbi.nlm.nih.gov/pmc/> now everything hinges on an approach that's already run into trouble
- Overview of Alzheimer's Disease. Chicago (US): Alzheimer's Association; 2009. Alzheimer's Disease Facts and Figures; ©2009; [about 80p]. Available from: https://www.alz.org/national/documents/report_alzfactsfigures2009
- Pei JJ, Tanaka T, Tung YC, Braak E, Iqbal K, Grundke II. (1997) Distribution, levels, and activity of glycogen synthase kinase-3 in the Alzheimer diseased brain. J Neuropathol Exp Neurol.

- 1997 Jan; 56(1):70-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed>
- Rajasekar S, Elango R. (2011) Estimation of alkaloid content of Ashwagandha (*Withania somnifera*) with HPLC methods. *Journal of Experimental Sciences* 2(5): 39-41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/1903170>
- Ramsey L (2018) The search for new Alzheimer's treatments just faced another setback – and
- Sauer A (2017) Has Alzheimer's Research Been Wrong for 20 Years? <https://www.alzheimers.net>
- Schaffer B, Wiedau-Pazos M, Geschwind DH.(2003) Gene structure and alternative splicing of glycogen synthase kinase 3 beta (GSK-3beta) in neural and non-neural tissues. *Gene* 302:73–81. Available from: <http://europepmc.org/abstract/med/12527198> DOI: 10.1016/S0378-1119(02)01092-2
- Shin RW, Iwaki T, Kitamoto T, Tateishi J. (1991) Hydrated autoclave pretreatment enhances tau immunoreactivity in formalin-fixed normal and Alzheimer's disease brain tissues. *Lab. Invest.* 64 (5): 693–702. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/1903170>
- Singh N, Bhalla M, Jager PD, Gilca M. (2011) An Overview on Ashwagandha: A Rasayana (Rejuvenator) of Ayurveda. *African Journal of Traditional, Complementary and Alternative Medicines.* 8(5): 208-213. Available From: <https://www.ncbi.nlm.nih.gov/pubmed/21511111>
- nlm.nih.gov/ pmc/Articles/ PMC3252722 DOI: 10.4314/ ajt-cam.v8i5S.9
- Spencer JPE. (2007) The interactions of flavonoids within neuronal signaling pathways. *Genes Nutr.* 2(3): 257–273
- Spencer JPE. (2010) The impact of fruit flavonoids on memory and cognition. *British J Nutrition* 104: S40–S47
- Stambolic V, Woodgett JR. (1994) Mitogen inactivation of glycogen synthase kinase-3 beta in intact cells via serine 9 phosphorylation. *The Biochemical Journal.* 303(Pt 3): 701–4. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1137602>
- Taniguchi S, Suzuki N, Masuda M, Hisanaga SI, Iwatsubo T, Goedert M, Hasegawa M. (2004) Inhibition of Heparin-induced Tau Filament Formation by Phenothiazines, Polyphenols, and Porphyrins. *The Journal of Biological Chemistry* 280(9): 7614–7623. Available from: <http://www.jbc.org/content/280/9/7614> DOI: 10.1074/jbc.M408714200
- Underwood E. (2016) Tau protein—not amyloid—may be key driver of Alzheimer's symptoms. *Brain & Behavior, Memory Topic* Available from: <https://www.sciencemag.org/news/2016/05> DOI:10.1126/science.aaf9985.
- Yin YI, Bassit B, Zhu L, Yang X, Wang C, Li YM. (2007) γ -Secretase Substrate Concentration Modulates the A β 42/A β 40 Ratio Implications For Alzheimer Disease. *J Biological Chemistry* 282: 23639-23644