

Morphology and Seasonal variation of fowl tape worm, *Raillietina tetragona* (Molin, 1858) in Purba Bardhaman, West Bengal, India

Sreenita Ghosh, Anadi P. Nandi and Soumendranath Chatterjee*

Parasitology and Microbiology Research Laboratory, Department of Zoology, Burdwan University, Purba Bardhaman 713104, West Bengal, India

ABSTRACT

The tape worm, *Raillietina tetragona* (Molin 1858) is a common parasite of domestic fowl, *Gallus domesticus* in West Bengal. This cestode parasite plays a significant role in growth and reproduction of fowl. In the current study, Light microscopic (LM) examination showed that the present cestode has a minute pin like scolex or head (18.98µm – 19.76 µm) and a long flat body divided into immature, mature and gravid proglottids. Head bears a rostellum (2.94 -3.11µm), four acetabula (6.79µm - 7.4µm) and followed by a neck (15.1µm – 16.27µm). Seasonal variation of *R. tetragona* from *Gallus domesticus* in Purba Bardhaman, West Bengal, India has been studied through two years span (January, 2016 to December, 2017). Monthly variations of infection of *Raillietina tetragona* showed comparatively higher prevalence in hot summer months (58.33 - 70 %) and humid rainy season (80-95.24%) and lower prevalence during winter season (0 - 10%). Mean intensity of *Raillietina tetragona* was also higher in summer (1-2.5) and rainy season (3.36-10.2). The present study on the morphology and seasonal prevalence of *Raillietina tetragona* in Purba Bardhaman region of West Bengal showed that more awareness should be focused towards upgrading and maintenance of the local free ranging chicken breeds.

KEY WORDS: *RAILLIETINA TETRAGONA*, *GALLUS DOMESTICUS*, LIGHT MICROSCOPIC OBSERVATIONS, PREVALENCE, PURBA BARDHAMAN, INDIA

ARTICLE INFORMATION:

Corresponding Author: soumen.microbiology@gmail.com

Received 2nd July, 2019

Accepted after revision 7th Sep, 2019

BBRC Print ISSN: 0974-6455

Online ISSN: 2321-4007 CODEN: USA BBRCBA

Thomson Reuters ISI ESC / Clarivate Analytics USA



NAAS Journal Score 2019: 4.31 SJIF: 4.196

© A Society of Science and Nature Publication, Bhopal India

2019. All rights reserved.

Online Contents Available at: <http://www.bbrc.in/>

DOI: 10.21786/bbrc/12.3/25

INTRODUCTION

Domestic fowl, *Gallus domesticus* (Linnaeus 1758) (Galliformes: Phasianidae), is supposed to have derived from the South East Asian and wild Indian red forest chicken (Permin and Ranvig, 2001). These fowls are with high nutritional value. Sometimes these are domesticated traditionally under free-range management systems in villages where little or no supplementary foodstuff is provided and the poultry farmers are unaware of any veterinary care, which finally leads to the parasitic infections to the chicken (Gary and Richard 2012). According to Rohde (1994), there are some other factors like temperature, humidity, rainfall and parasite maturation which influence the parasitic infection in domestic fowl. Thus environment plays a significant function in the seasonal variability of these cestode parasites. During the present study *G. domesticus* were collected from Purba Bardhaman and surrounding rural areas while some other gastrointestinal cestode parasites were also recovered from the domestic fowl gut and *R. tetragona* was found to be the most prevalent species among all. This is also supported by the findings of Khan et al. (2016).

In the present survey morphological observations were made using light microscope (LM) and measurements of its cephalic regions were taken. Other morphological characters like shape of the scolex; rostellum and suckers, number of eggs in the egg capsules and shape of the eggs were also made. However, reports on seasonal prevalence of *Raillietina tetragona* in domestic fowl of Purba Bardhaman region of West Bengal could not be noticed in the accessible reports. The two year survey was carried out to determine the prevalence of this cestode parasite in *G. domesticus* occurring in and around Purba Bardhaman, West Bengal, India.

MATERIAL AND METHODS

Freshly dissected out guts of the domestic fowl, *Gallus domesticus* were purchased from the local markets of Bardhaman, West Bengal, brought to the laboratory, as soon as possible, and examined for the presence of *Raillietina tetragona* in each week of every month for a period of consecutive two year study period from January 2016 to December 2017. Live worms were washed in normal saline (0.85% NaCl) and freed from the adhering host materials, fixed and preserved in 70% alcohol. For identification and morphological observations, the adult cestodes were washed in normal saline, dried and flattened on filter paper, fixed in 70% alcohol and observed with the help of compound light microscope.

The prevalence of infection and mean intensity were calculated from the recorded data with the help of the following formulas

$$\text{Prevalence of infection} = \frac{\text{no. of infected } Gallus domesticus}{\text{total number of } Gallus domesticus \text{ examined (infected + uninfected)}} \times 100$$

(In percentage)

$$\text{Mean intensity} = \frac{\text{Total no. of } Raillietina \text{ sp. collected from infected intestine}}{\text{No. of infected } Gallus domesticus \text{ in the sample}}$$

RESULTS

General morphology of the cestode

Light microscopic (LM) study showed that the body of *R. tetragona* was typically divided into scolex, followed by a short unsegmented region, the neck, succeeded by a chain of proglottids consisting of immature, mature and gravid segments. Scolex is small pin like and provided with four well-formed hemispherical acetabula, armed with hooks and an armed disc shaped rostellum (Plate 1). In the mature proglottids the genital pores are unilateral (Plate 3) and in the gravid proglottids (Plate 4) eggs are found in egg capsules, each containing six to twelve eggs.

The current LM observation showed that the scolex of *Raillietina tetragona* was about 18.98µm – 19.76µm in width, neck was about 15.1µm– 16.27µm and rostellum was 2.94µm – 3.11µm in width and armed with one row of hooks. Four acetabula were measured about 7.08µm, 6.95µm, 6.79µm and 7.4µm in length and 4.12µm, 4.03µm, 3.17µm and 3.71µm in width respectively. Mature proglottids (Plate 2) were rectangular in shape (65.68µm – 68.96µm) and gravid proglottids (60.85µm– 61.44µm) were more or less square containing numerous egg capsules (Plate 5) which were more or less round or oval in shape and measured about 0.85µm– 1.09µm.

During the two year study period from January 2016 to December 2017, out of 382 fowl guts examined, 230 were found infested with *R. tetragona* and totally 907 parasites were collected. As *R. tetragona* was isolated all through the year from *Gallus domesticus*, characterization and seasonal prevalence were done for this species only. Monthly variations in respect to percentage of prevalence and the mean intensity of *R. tetragona* infection have been presented in Table 1 and total number of parasite present in *Gallus domesticus* in each season for the two year study period has been shown in Figure 1. The result showed that average prevalence percentage was 51.91%. The study revealed that the higher percentage of prevalence of *R. tetragona* occurred in summer season showing higher peak in May (61.58% -70%) and it showed its highest values during rainy season and the maximum value of percentage of prevalence was recorded in the month of July (93.33% - 95.24%). From September to November it tends to lower down and the lowest prevalence occurred in winter (0 -10%). The mean

Table 1. Monthly variation in % of prevalence and Mean intensity of infection of <i>Raillietina tetragona</i> infecting <i>Gallus domesticus</i>										
Month of collection	Number of gut examined		Infected host		Parasite present		Prevalence %		Mean Intensity	
	2016	2017	2016	2017	2016	2017	2016	2017	2016	2017
January -16	8	10	0	0	0	0	0	0	0	0
February -16	8	14	1	2	1	3	12.5	14.28	1	1.5
March -16	12	16	2	2	2	4	16.67	12.5	1	2
April -16	12	10	7	6	12	10	58.33	60	1.71	1.66
May-16	20	13	14	8	25	20	70	61.58	1.79	2.5
June-16	20	17	16	15	54	67	80	88.24	3.36	4.46
July-16	15	21	14	20	120	204	93.33	95.24	8.57	10.2
August -16	19	23	16	21	95	119	84.21	91.3	5.93	5.66
September -16	25	15	18	11	50	52	72	73.33	2.78	4.72
October-16	24	28	20	23	22	26	83.33	82.14	1.1	1.13
November -16	15	16	7	5	9	10	46.67	31.25	1.28	2
December -16	10	11	1	1	1	1	10	9.09	1	1
Total duration: 2 years	total no. of sample: 382		Total number of infected host: 230		Total number of parasite present: 907		Mean percentage of prevalence: 51.91%			
Period of study: January 2016 to December 2017										

intensity of infection also declined in the winter season (0-1) during the two year study period.

DISCUSSION

Raillietina tetragona has a cosmopolitan distribution occurring in pigeon, chicken and guinea fowl. This tape-worm completes its development in two hosts. Birds are the definitive hosts and ants, mostly of the species *Tetramorium*, and housefly of the species *Pheidole* and *Musca* act as the intermediate hosts (Horsfall 1938;

Soulsby 1982; Su 1986). Light Microscopic observations revealed that *Raillietina tetragona* was different from the other *Raillietina spp.* in respect to the shape of scolex, rostellum and acetabula. The hooks of the rostellum were placed in one row (Butboonchoo et al., 2016). This type of study was also carried by some scientists earlier (Hofstad et al. 1984; Sawada 1964 and 1965) but scanty literature is available. Lachhandama (2009) observed the morphological structures of *R. echinobothrida* from fowl gut and Mu et al. (2009) worked on *R. echinobothrida* and *R. tetragonal* from the gut of domestic







FIGURE 1. Seasonal prevalence (No. of parasite) of *Raillietina tetragona* present in *Gallus domesticus* during two year study period from January 2016 to December 2017




chicken. In the recent year, Butboonchoo *et al.* (2016) distinctly differentiated four different species of *Raillietina* spp. including *Raillietina tetragona* on the basis of Light Microscopic and Scanning Electron Microscopic observations. The present study showed similar findings.

Seasonal variability of *R. tetragona* in domestic fowl, *Gallus domesticus*, was checked throughout the year and comparatively higher percentage of prevalence and mean intensity of this parasite were observed in the summer and rainy season. Seasonal variability of cestodes were showed that changes in prevalence and infection rates could depend upon host, parasite and also upon the geographical location (Biswal *et al.* 2013). Oniye *et al.* (2001) in Nigeria reported a high prevalence of



Raillietina tetragona in the month of June and August. Fakae *et al.* (1991) from Eastern Nigeria reported 72.5 % prevalence of *R. tetragona* in *Gallus gallus* during dry season (November to April). These findings were supported by the previous results as reported by Frontovo, (2000), Fakae *et al.* (1991) and Oniye *et al.* (2001).

Seasonal prevalence of *R. tetragona* also varied with geographical location and other climatic conditions. According to Adang *et al.* (2008) *R. tetragona* was found to be the most common cestode parasite in domestic pigeon, occurring in 11 months of the annual cycle, showing highest prevalence in rainy season in Zaria, Northern Nigeria. Salam *et al.* (2010) and Sheikh *et al.* (2016) reported that the prevalence of *R. cesticillus* from *Gallus gallus domesticus* in Kashmir is highest in summer followed by autumn, spring and winter months respectively. Butt *et al.* (2014) reported that the average percentage of prevalence of *Raillietina cesticillus* in *G. domesticus* was 83.5% during July to November, 2013 in Hyderabad, Sindh, Pakistan. Accord-





ing to the observations of Sheikh *et al.* (2016) the high prevalence of *Raillietina sp.* infection was seen during warm summer followed by autumn, spring and winter months. Except winter, the high prevalence of cestode infection throughout the year is a strong evidence of unscientific management and control in domestic fowls. As a result an infection creating environment originates continuously. However, Shukla *et al.* (2012) reported that the maximum prevalence of *Raillietina sp.* in *G. domesticus* from Ahmednagar district, Maharashtra, India was found in winter followed by rainy season and minimum prevalence was noted in summer. According to Patil and Bhamare (2018), a high prevalence of *Raillietina sp.* in *G. domesticus* from Nashik district of Maharashtra, India occurred in winter season followed by summer season and low in rainy season. A possible explanation of this report is the availability of the intermediate host and oncosphere stages of *Raillietina sp.* increases during cold climatic condition. The oncosphere stages are taken by the intermediate hosts and grow into cysticercoids which become adult in the definitive host during summer season.

In our survey greater number of mature cestodes were obtained during summer and rainy seasons whereas in autumn mostly immature stages were obtained. High rate in the percentage of prevalence and mean intensity of *R. tetragona* in the summer and rainy season may be a result of the abundance of infective stage and the stage bearing intermediate hosts in the survey zone during these seasons. It seems that moisture and temperature favour the development of intermediate hosts, the development of eggs in soil as well as proliferate the reproductive rate of parasites (Møller, 2010). Low prevalence of *R. tetragona* in winter season may be the result

of elimination of multiple infections. This condition may occur due to intraspecific antagonistic reaction.

In fact, reports from other researchers around the world in different times showed that *Raillietina spp.* has a very high prevalence of infection in birds especially in indigenous chickens. (Ahmed and Sinha, 1993; Puttalakshamma *et al.* 2008; Nnadi *et al.* 2010; Sreedevi *et al.* 2016; da Silva *et al.* 2016). According to a recent report of Jajere *et al.* (2018) who found the guinea fowls have been heavily infected with *Raillietina spp.* and among the various species, *Raillietina tetragona* was most abundant having 72.8% prevalence of infection. When the infective stages of intestinal parasites pass out to the surroundings, they have to get over the environmental hazards before they reach their suitable hosts for further development (Gillett, 1974). So, favourable environmental conditions are also necessary for the effective transmission of the parasites (Smith, 1990) and seasonal variations in the social conduct of the host and their accumulation can also describe for seasonal dynamics of parasitic infections (Hosseini *et al.* 2004). The time period for obtaining table size, in case of indigenous breed, is longer than exotic breeds which are usually feed on artificial supplementary diets. So another reason of higher infection is may be due to improper management, the inadequacy of food grains for the local breeds and they have to feed on insects, mites and worms which may be the intermediate hosts for the cestode parasite and has been shown to improve susceptibility to parasitism in the system (Shukla and Mishra, 2013).

Moreover, earlier studies have shown that the immune systems of humans, rodents and birds got weaker by rough climatic conditions, under-nutrition or investment in reproduction (Lloyd 1995; Nelson *et al.* 2002) and due to low immunity level, hosts become more vulnerable to infections (Hillgarth and Wingfield, 1997). A longer breeding period of the host bird provides a selective advantage to the parasites (Dunn and Winkler 2010; Møller *et al.* 2010). Reports have shown that during the breeding season of host birds, antibody production and cell mediated immunity rate is getting weaker which is the reason for higher rates of parasitic infection (Hillgarth and Wingfield, 1997; Moreno *et al.* 2001). According to Sheikh *et al.* (2016) the host birds have shown a moderate resistance against the cestode parasites with the advancement of age because of the improved immune status in grown-ups than in young ones. Chicks hatch out during late spring. During summer and autumn they are in their young age when they get exposed to parasite infection. The young and the juvenile forms are susceptible to the parasite because they are immunologically weak. This may also affect the seasonal variability of *Raillietina tetragona*. Till now

there are many surveys on *Raillietina spp.* reported from different parts of the world but it still needs to execute more studies to follow the changing dynamics of helminth infection in domestic poultry management.

ACKNOWLEDGEMENT

Thankful acknowledgement is due to the Department of Zoology, University of Burdwan, for giving the laboratory facilities

Author contribution: Sreenita Ghosh has carried out the present work under the guidance of Dr. A. P. Nandi and Dr. Soumendranath Chatterjee (Professor of Zoology).

Compliance with ethical standards: Though our Institution does not have such ethics committee, domestic fowl has been slaughtered for the present work following the guiding principle of CPCSEA (The Committee for the Purpose of Control and Supervision of Experiments on Animals) established by the Act of the Indian Parliament.

Conflict of interest: All the authors have declared that in the present work there is no conflict of interest.

REFERENCES

- Adang K.L., Oniye S.J., Ajanusi O.J., Ezealor A.U., Abdu P.A. (2008) Gastrointestinal helminths of the domestic pigeons (*Columba livia domestica* Gmelin, 1789 Aves: Columbidae) in Zaria, Northern Nigeria. *Sci World J* Vol. 3 No 1: Pp 33–37
- Ahmed M.I., Sinha P.K. (1993) Prevalence of poultry helminthiasis in an arid zone in Nigeria. *Indian Vet J* Vol. 70: Pp 703–704
- Biswal D., Nandi A.P., and Chatterjee S. (2013) Temporal variation of the cestode, *Cotugnia cuneata* (Meggit, 1924) in their host, domestic pigeons, *Columba livia domestica* (Gmelin, 1789). *J Parasit Dis* Vol. 39 No 2: Pp 194–199
- Butoonchoo P., Wongsawad C., Rojanapaibul A., Chai J.Y. (2016) Morphology and Molecular phylogeny of *Raillietina* spp. (Cestoda: Cyclophyllidae: Davaineidae) from domestic chickens in Thailand. *Korean J Parasitol* Vol. 54 No 6: Pp 777–786
- Butt Z., Shaikh A.A., Memon S.A., Mal B. (2014) Prevalence of Cestode parasites in the intestine of local chicken (*Gallus Domesticus*) from Hyderabad, Sindh, Pakistan. *Journal of Entomology and Zoology Studies* Vol. 2 No 6: Pp 301–303
- Da Silva G.S., Romera D.M., Fonseca L.E.C., Meireles M.V. (2016) Helminthic parasites of chickens (*Gallus domesticus*) in different regions of São Paulo State, Brazil. *Braz J PoultSci* Vol. 18: Pp 163–168.
- Dunn P.O., Winkler D.W. (2010) Effects of climate change on timing of breeding and reproductive success in birds. In: *Effects of climate change on birds*. Pp 113–128 (Eds) Møller A.P., Fiedler W., Berthold P., Oxford University Press, Oxford.
- Fakae B.B., Umeorizu J.M., Orajaka L.J.E. (1991) Gastrointestinal helminth infection of the domestic fowl (*Gallus gallus*) during the dry season in eastern Nigeria. *J Zool Vol.* 105: Pp 503–508.
- Frantovo D. (2000) Some parasitic nematodes (Nematoda) of birds (Aves) in the Czech Republic. *Acta Societatis Zoologicae Bohemicae*
- Gary D.B. and Richard D.M. (2012) Intestinal parasites in backyard chicken flock 1 In: *Series of Veterinary Medicine- Large animal clinical sciences*. Vol 76, University of Florida. <http://edis.ifas.ufl.edu>
- Gillett J. (1974) Direct and indirect influences of temperature on the transmission of parasites from insects to man. In: *The effects of meteorological factors upon parasites*. Pp 79–95 (Ed) Taylor A.E.R., Muller R., Blackwell Scientific Publications, London.
- Hillgarth N., Wingfield J.C. (1997) Testosterone and immunosuppression in vertebrates: implications for parasite-mediated sexual selection. In: *Parasites and pathogens: effects on host hormones and behaviour*. Pp 143–155 (Ed) Beckage N.E., Chapman and Hall, New York.
- Hofstad M.S., Calnek B.W., Helmboldt C.F., Reid W.M., Yoder H.W. (1984) *Diseases of Poultry*. Iowa State University Press
- Horsfall M.W. (1938) Observations on the life history of *Raillietina echinobothrida* and of *R. tetragona* (cestoda). *The Journal of Parasitology* Vol. 24 No 5: Pp 409–421
- Hosseini P.R., Dhondt A.A., Dobson A. (2004) Seasonality and wildlife disease: how seasonal birth, aggregation and variation in immunity affect the dynamics of *Mycoplasma gallisepticum* in house finches. *Proc R SocLond B* Vol. 271: Pp 2569–2577.
- Jajere S.M., Lawal J.R., Atsanda N.N., Hamisu T.M., Goni M.D. (2018) Prevalence and burden of gastrointestinal helminthes among grey-breasted helmet guinea fowls (*Numida meleagris galeata*) encountered in Gombe state, Nigeria. *International Journal of Veterinary Science and Medicine* Vol. 6 No 1: Pp 73–79
- Khan A., Bhutto B., Shoaib M., Fahad S., Ahmed A., Khetrani B.I., Nizamani A.R., Zeb A., Rahman M.U., Khan S. (2016) Prevalence of gastrointestinal helminths in Banaraja fowls reared in semi-intensive system of management in Mayurbhanj district of Odisha, *Journal of Animal Health and Production* Vol. 4 No 1: Pp 26–30
- Lalchhandama K. (2009) On the structure of *Raillietina echinobothrida*, the tapeworm of domestic fowl. *Sci Vis* Vol. 9 No 4: Pp 174–182
- Linnaeus C. (1758) *System Nat.* ed. 10:58
- Lloyd S. (1995) Environmental influences on host immunity. In: *Ecology of infectious diseases in natural populations*. Pp 327–361 (Ed) Grenfell B.T., Dobson A.P., Cambridge University Press, Cambridge.
- Møller A.P. (2010) Host-parasite interactions and vectors in the barn swallow in relation to climate change. *Global Change Biol* Vol. 16: Pp 1158–1170

- Møller A.P., Flensted-Jensen E., Klarborg K., Mardal W., Nielsen J.T. (2010) Climate change affects the duration of the reproductive season in birds. *J AnimEcol* Vol. 79: Pp 777–784
- Moreno J., Sanz J.J., Merino S., Arriero E. (2001) Daily energy expenditure and cell-mediated immunity in pied flycatchers while feeding nestlings: interaction with moult. *Oecologia* Vol. 129: Pp 492–497
- Mu L., Li H.Y., Yan B.Z. (2009) Comparative study on morphology and development of two species of *Raillietina* from chicken. *Chinese Journal of Parasitology & Parasitic Diseases* Vol. 27 No 3: Pp 232–236
- Nelson R.J., Demas G.E., Klein S.L., Kriegsfeld L.J. (2002) Seasonal patterns of stress, immune function, and disease. Cambridge University Press, New York.
- Nnadi P.A., George S.O. (2010) A cross-sectional survey on parasites of chickens in selected villages in the subhumid zones of south-eastern Nigeria. *J Parasitol Res* Vol. 2010: Pp 141824
- Oniye S.J., Audu P.A., Adebote D.A., Kwaghe B.B., Ajanusi O.J., Nfor M.B. (2001) Survey of helminth parasites of Laughing Dove (*Streptopelia segalensis*) in Zaria, Nigeria. *African Journal of Natural Sciences* Vol. 4: Pp 65–66
- Patil S.D. and Bhamare A.V. (2018) Seasonal variation of cestode parasite *Raillietina* in an edible bird *Gallus domesticus* (L.). *Environment Conservation Journal* Vol. 19 No 3: Pp 77–80
- Permin A. and Ranvig H. (2001) Genetic resistance in relation to *Ascaridia galli* in chickens. *Veterinary Parasitology* Vol. 102 No 12: Pp 101–111
- Puttalakshamma G.C., Ananda K.J., Prathiush P.R., Mamatha G.S., Rao S. (2008) Prevalence of gastrointestinal parasites of poultry in and around Bangalore. *Vet World* Vol. 1: Pp 201–202
- Rohde K. (1994) Niche restriction in parasites: proximate and ultimate causes. *Parasitology* Vol. 109(suppl.): Pp S69–S84.
- Salam S.T., Mir M.S., Khan A.R. (2010) The prevalence and pathology of *Raillietina cesticillus* in indigenous chicken (*Gallus gallus domesticus*) in the temperate Himalayan region of Kashmir. *VeterinarskiArhiv* Vol. 80: Pp 323–328.
- Sawada I. (1964) On the genus *Raillietina* Fuhrmann, 1920 (I). *J Nara GakugeiUniv (Natural Science)* Vol. 12: Pp 19–36.
- Sawada I. (1965) On the genus *Raillietina* Fuhrmann 1920 (II). *J Nara GakugeiUniv (Natural Science)* Vol. 13: Pp 5–38
- Sheikh B.A., Ahmad F., Sofi T.A. (2016) Morphology and Prevalence of Some Helminth Parasites in *Gallus domesticus* from Gurez Valley of Jammu and Kashmir, India. *Journal of Fisheries and Livestock Production* Vol. 4: Pp 159
- Shukla S. and Mishra P. (2013) Gastro Intestinal Helminths Parasites of Local Chickens Samples from Tribal Areas of Madhya Pradesh, India. *Int. J. of Life Sciences* Vol. 1 No 4: Pp 284–287
- Shukla S.J., Borde S.N., Humbe A., Bhavare V.V. (2012) Seasonal variation of intestinal Tapeworms in *Gallus gallus domesticus* at Ahmednagar region. *IntMultidiscip Res J* Vol. 2 No 4: Pp 01–03
- Smith G. (1990) The population biology of the free-living phase of *Haemonchus contortus*. *Parasitology* Vol. 101: Pp 309–316.
- Soulsby E.J.L. (1982) Helminths, arthropods and protozoa of domestic animals, 7th edn. FLBS Barrierve Tindal. London, Pp-235–244
- Sreedevi C., Jyothisree C.H., Rama Devi V., Annapurna P., Jeyabal L. (2016) Seasonal prevalence of gastrointestinal parasites in desi fowl (*Gallus gallus domesticus*) in and around Ganavaram, Andhra Pradesh. *J Parasit Dis* Vol. 40: Pp 656–661
- Su X.L.Y. (1986) Studies on the life history of *Raillietina tetragona* Molin and its natural intermediate host in Xiamen. *Wuyi Science Journal* 06 (epub)