

In vitro Phytochemical Profiling of Anti Diabetic and Anti Inflammatory Activities of *Premna rajendranii* – An Endemic Species

Steffy Francis¹, Anand Gideon V², John Britto and J.Mariyaraj¹

¹PhD Research scholar, Department of Botany, Bishop Heber College, Trichy

²Associate Professor and Head, Department of Botany, Bishop Heber College, Trichy

³The Rapinat Herbarium and Centre for Molecular Systematics, St. Joseph's College, Trichy
(Affiliated to Bharathidasan University, Tiruchirappalli – 24.)

ABSTRACT

The present study aims to find out the presence of phytochemicals and evaluate the anti diabetic and anti-inflammatory activities of *Premna rajendranii*. Anti diabetic assay was evaluated using α - amylase inhibitory assay by DNSA method, anti-inflammatory activity was tested using the method of Mizushima et al with simple modifications and the methods of Peach and Tracey (1956), Gibbs (1974), Harborne (1984), Trease and Evans (1985), Edeoga et al.,(2005), Khandewal (2008), Kokate et al.,(2001), Sofowara (2009) and Tiwari et al.,(2011) were used to identify the nature of phytochemical constituents present in *Premna rajendranii*. Anti diabetic analysis of methanolic leaf extract of *Premna rajendranii* showed optimal activity when compared to standard drug. Maximum inhibition showed at 100 μ g (79.3%) and minimum inhibition showed at 25 μ g (44.82%). Anti-inflammatory activity of methanolic leaf extract of *P.rajendranii* has shown high inhibition than standard at 50 μ l i.e. 33.88 \pm 0.01. Phytochemical screening shows the presence of Alkaloids, Carbohydrates, Flavonoids, Steroids, Glycosides, Phenols, Proteins, Tannins, Saponins, and Terpenoids in the leaf of *Premna rajendranii*. Anti diabetic and anti-inflammatory analysis showed the presence of bioactive compounds and the medicinal values of *Premna rajendranii*.

KEY WORDS: *PREMNA RAJENDRANII*, *ANTI DIABETIC*, *ANTI-INFLAMMATORY*, *PHYTOCHEMICAL ANALYSIS*.

ARTICLE INFORMATION

*Corresponding Author: steffyclarefcc@gmail.com

Received 04th Oct 2019

Accepted after revision 21th Dec 2019

Print ISSN: 0974-6455 Online ISSN: 2321-4007

CODEN: BBRCBA

Thomson Reuters ISI Web of Science Clarivate Analytics
USA and Crossref Indexed Journal



NAAS Journal Score 2019 (4.38) SJIF: 2019 (4.196)
A Society of Science and Nature Publication, Bhopal India
2019. All rights reserved.
Online Contents Available at: <http://www.bbrc.in/>
DOI: 10.21786/bbrc/12.4/24

INTRODUCTION

Plants have an important role in the day to day life of mankind. Each and every plant contains phytochemicals either with medicinal properties or poisonous properties. Scientists and researchers are more focused on medicinal plants for the benefit of mankind. According to them each of the medicinal plant is a source of drug candidate. From ancient times India is considered a rich repository of medicinal plants and Indian traditional systems of medicine such as Ayurveda, Siddha and folk medicine are well known. Genus *Premna* has an indispensable value in Indian health care practice. Species of *Premna* are known for the preparation of famous Ayurvedic formulation Dashamoolaarishtam (Joshi et al., 2017). *Premna*'s medicinal properties have been used in Indian traditional system of medicine especially for diarrhoea, stomach and hepatic disorders. The various biological activities including antioxidant, antibacterial, anti-inflammatory, cytotoxic and hepatoprotective have been displayed both at extract and pure compound level (Rekha, 2015). During the taxonomic revision of Indian Verbenaceae, Rajendran and Daniel (2002) recognized 31 species and 6 varieties of *Premna*. Recently, PrabhuKumar et al.

(2013) reported the discovery of a new species *Premna rajendranii* from Western Ghats (Chinnar of Kerala and Madukkarai of Tamilnadu). Apart from this, a research team comprising Robi, Augustin, Sasidharan and Udayan (2013) rediscovered an endemic and rare species *Premna paucinervis* (C. B. Clarke) Gamble from the Vagamon hills along South Western Ghats of Kerala after a lapse of 140 years of its original type collection by R.H. Beddome (1872) from the Anamalais in the Western Ghats (Tamilnadu) (Bose, 2014). *Premna* included earlier in the family Verbenaceae, was recently transferred to the Lamiaceae family based on molecular data (A.P.G. IV. 2009; Francis et al., 2019).

MATERIALS AND METHODS

Collection and authentication: The plant material for our investigation was collected from the scrub jungles of the Madukkarai Hills, Coimbatore

District in Tamilnadu, and authenticated by Dr. S. John Britto S.J, at the Rapinat Herbarium and Centre for Molecular Systematics, St. Joseph's College (Autonomous), Tiruchirappalli. The voucher Specimen (RHT 68887) was deposited in the Rapinat Herbarium.

Extraction: The leaves were shade dried and powdered using mechanical grinder. The powder sample was stored in an air tight container and the portion of the powder was taken in test tube and solvent (Methanol) was added to it such that plant powder soaked in it and shaken well. The solution then filtered with the help of muslin cloth and filtered extract was taken and used for antidiabetic and anti-inflammatory analysis.

Preliminary Phytochemical Analysis: Test's for Alkaloids: Mayer's Test: To a few ml of filtrate, one or two drops of Mayer's reagent were added by the side of the test tube. A white creamy precipitate indicated the test as positive. Wagner's Test: To a few ml of filtrate, few drops of Wagner's reagent were added by the side of the test tube. A reddish brown precipitate confirmed the test as positive. Hager's Test: To a few ml of filtrate 1 or 2 ml of Hager's reagent (Saturated aqueous Solution of picric acid) was added. A prominent yellow precipitate indicated the test as positive. (Kokate et. al., 2001).

Test for flavonoids: Pears Test: To 2-3ml extract, was added zinc powder in a test tube, followed by drop wise addition of conc. HCL. Formation of purple red or cherry colour indicated the presence of flavonoids (Peach et.al., 1956). Lead acetate test: 1ml extract was treated with 1ml 10% lead acetate (Pb(OAc)₄) solution. Formation of yellow colour precipitate indicated the presence of flavonoids. Alkaline reagent test: To 2ml test solution, sodium hydroxide solution was added to give a yellow or red colour (Khandewal et.al. 2008). Conc.H₂SO₄ test: 5ml of dilute ammonia solution was added to the extract followed by conc.H₂SO₄. Yellow colour indicated the presence of flavonoids. Tests for Phenolic Compounds and Tannins: Ferric Chloride Test: The extract (50 mg) was dissolved in 5 ml of distilled water. To this few drops of neutral 5% Ferric Chloride solution was added. A dark green colour indicated the presence of phenolic compounds.

Potassium dichromate test: To the extract add 5% potassium dichromate solution. Positive result was confirmed by a formation of brown precipitate (for phenol).

Lead Acetate Test: The extract (50 mg) was dissolved in distilled water and to this 3 ml of 10% Lead Acetate solution was added. A bulky white precipitate indicates the presence of phenolic compounds (Treare et.al., 1985). **Braymer's Test:** To 2 ml extract, added 2 ml H₂O and followed with 2-3 drops of FeCl₃ (5%). Green precipitate proved presence of tannins. **Tests for Saponins:** **Foam Test:** To 1ml of extract, add 2ml of distilled water and shaken vigorously and allowed to stand for 10 min. There is the development of foam on the surface of the mixture. Then shake for 10 minutes, it indicates the presence of saponins (Khandewal et.al., 2008). **NaHCO₃ Test:** To extract a drop of sodium bicarbonate was added. The mixture was shaken vigorously and kept for 3 min. A honey comb like froth was formed and

it showed the presence of saponins. **Tests for Glycosides:** **Keller Kiliani Test** (Test for cardiac glycoside): To 2 ml extract, was added 1 ml glacial acetic acid, one drop 5% FeCl₃ and 1 ml conc. H₂SO₄. A brown ring of the interface indicated the presence of cardiac glycosides (Kokate et.al., 2001; Khandewal et.al., 2008). **Glycoside Test:** To small amount of extract, was added 1 ml water and shake well. Then aqueous solution of NaOH was added. Yellow color appeared that indicated the presence of glycosides (Treare et.al., 1985).

Molisch's Test: To 1ml of extract, 2drops of Molisch's reagent was added in a test tube and 2ml of con. H₂SO₄ was added carefully keeping the test tube slightly curved. Formation of violet ring at the junction indicated the presence of glycosides (Khandewal et.al., 2008).

Tests for Carbohydrates: **Molish's Test:** To 2 ml of filtrate two drops of alcoholic solution of α - naphthol was added, the mixture was shaken

Table: 1.1 Preliminary phytochemical analyses in *Premna rajendranii* (leaf)

| | Test | Chloroform | Acetone | Ethanol | Methanol | Aqueous |
|-----------------|---|------------|---------|---------|----------|---------|
| Alkaloids | Wager's | + | + | ++ | +++ | ++ |
| | Hager's | — | + | ++ | +++ | ++ |
| | Mayer's | — | + | ++ | +++ | + |
| Flavonoids | Pew's | + | + | + | ++ | — |
| | Lead Acetate | + | +++ | +++ | +++ | +++ |
| | NaOH | — | +++ | +++ | +++ | +++ |
| Phenol & tannin | Con.H ₂ SO ₄ | + | +++ | +++ | +++ | ++ |
| | FeCl ₃ | + | + | + | ++ | + |
| | K ₂ Cr ₂ O ₇ | + | + | + | + | + |
| Saponins | Lead Acetate | — | — | + | ++ | + |
| | Braymers | + | — | + | + | + |
| | Foam | — | — | — | — | — |
| Glycosides | NaHCO ₃ | — | — | — | — | — |
| | Keller kiiani | — | +++ | +++ | +++ | +++ |
| | Glycosides | — | +++ | +++ | +++ | +++ |
| Carbohydrates | Molish | — | ++ | +++ | +++ | ++ |
| | Molish | — | + | ++ | ++ | ++ |
| | Benedicts | — | — | + | + | — |
| Terpenoids | Salkowskis | — | + | + | ++ | + |
| Quinones | Quinones | — | — | — | ++ | +++ |
| Sterols | Salkowskis | — | — | — | — | — |
| | Keller kiiani | — | + | + | ++ | ++ |
| Protein | Biuret | — | + | + | + | + |

well and 1 ml of con. H_2SO_4 was added slowly along the sides of the test tube and allowed to stand. A violet ring indicated the presence of carbohydrates. Benedict's test: To 0.5 ml of filtrate, 1 ml of Benedict's reagent was added. The mixture was heated on a boiling water bath for 2 mins. A characteristic coloured precipitate indicated the presence of sugar. Test for Terpenoids: Salkowski's Test: 2 ml of chloroform and 1 ml of conc. H_2SO_4 was added to 1 ml of extract and observed for reddish brown color that indicated the presence of terpenoids. Tests for quinones: 1ml of extract was treated with alcoholic potassium hydroxide solution. Quinines give coloration ranging from red to blue. Test for sterols: Salkowski's Test: To 2 ml of extract, was added 2 ml chloroform and 2 ml conc. H_2SO_4 from the side of the test tube. Chloroform layer appeared red and acid layer showed greenish yellow fluorescence indicated the presence of sterols (Khandewal et.al., 2008).

Tests for Proteins and Amino Acids: Biuret Test: An aliquot of 2 ml of filtrate was heated with 1 drop of 2 % H_2SO_4 solution. To this 1 ml of ethanol (95%) was added, followed by excess of KOH Pellets. Pink colour in the ethanolic layers indicated the presence of proteins. **Conc. H_2SO_4 Test:** 2 ml extract was treated with few drops of conc. H_2SO_4 . Formation of white precipitate indicated the presence of proteins.

Xantho proteins Test: 2 ml extract was treated with few drops of conc. HNO_3 and NH_3 solution. Formation of reddish orange precipitate indicated the presence of xantho proteins.


Anti-Diabetic Assay (A-Amylase Inhibitory Assay)

By Dnsa Method: α -amylase was dissolved in phosphate buffer saline (PBS, 0.02 mol/L, pH6.8) at a concentration of 0.1 mg/ml. Various concentrations of sample solutions (0.25 ml) were mixed with α -amylase solution (0.25 mL) and incubated at 37 °C for 5 min. Then the reaction was initiated by adding 0.5 mL 1.0% (w/v) starch substrate solution to the incubation medium. After incubation at 37 °C for 3 min, the reaction was stopped by adding 0.5 ml DNS reagent (1% Dinitrosalicylic acid, 0.05% Na_2SO_3 and 1% NaOH solution) to the reaction mixture and boiling at 100 °C for 5 min. After cooling to room temperature, the absorbance (Abs) at 540 nm was recorded by a spectrophotometer. The inhibition percentage was calculated by the following equation:

Table 2. Antidiabetic Activity of Methanolic leaf extract of *Premna rajendranii*

| SL.NO: | Concentration in (μ g) | % of inhibition | |
|------------------------|-----------------------------|-----------------|-----------------|
| | | Sample | Standard |
| 1 | 25 | 44.82 \pm 0.5 | 61.7 \pm 0.01 |
| 2 | 50 | 61.37 \pm 0.6 | 67.3 \pm 0.01 |
| 3 | 75 | 64.82 \pm 0.4 | 70.9 \pm 0.01 |
| 4 | 100 | 79.3 \pm 0.7 | 87.4 \pm 0.21 |
| IC ₅₀ value | 288.8 | 246.7 | |

Figure 1. Antidiabetic Activity of *Premna rajendranii*



$$\text{Percentage Inhibition} = \frac{[(\text{Abs1} - \text{Abs2})/\text{Abs1}] \times 100}{\times 100}$$

Where, Abs1= control and Abs2= sample.

Anti-Inflammatory Activity: Method of Mizushima et al was followed with simple modifications. The reaction mixture (0.5 ml) consisted of 0.45 ml bovine serum albumin (3% aqueous solution) and varying concentration of compound (25, 50, 100, 200 µg/ml of final volume), pH was adjusted to 6.3 using small amount of 1N hydrochloric acid. The samples were incubated at 37°C for 20 min and then heated at 80°C for 2min. After cooling the samples, 2.5 ml phosphate buffer saline (pH 6.3) was added to each tube. The absorbance was measured using spectrophotometer at 660nm. The percentage inhibition of protein denaturation was calculated as follows:

$$\text{Percentage of inhibition} = \frac{[(\text{Abs Control} - \text{Abs Sample}) / \text{Abs control}]}{\times 100}$$

RESULTS AND DISCUSSION


The result of phytochemical profiling of leaves of *P.rajendranii* indicates the presence of Alkaloids, Flavonoids, Phenols, Tannins, Glycosides, Carbohydrates, Terpenoids, Quinones, Sterols and Proteins. High concentrations of bioactive compounds were found in methanolic, ethanolic, aqueous, acetone extracts while very low concentrations in chloroform extract. Alkaloids, Flavonoids and Glycosides were mainly seen in most of the samples except chloroform extraction (Table 1).

Diabetes mellitus is one of the major metabolic disorders (Keerthana et al., 2013), and is characterized by high blood glucose levels. According to World Health Organization, 180 million people are currently suffering from DM and the rate is rapidly growing. There are many effective medicines for DM but with serious side effects. Recently researchers are in the experiments to find effective medicine for DM without any side effect. More than 1200 plants have a potential antidiabetic capacity (Modak et. al., 2007; Bailes, 2002; Chitturi and George, 2002; Kesari et. al., 2007, Prabhu et al 2018). Methanolic leaf extract of *P.rajendranii* was investigated for in vitro antidiabetic activity with respect to

Table 3. Anti-inflammatory Activity of *Premna rajendranii*

| SL.NO: | Concentration in (µg) | % of inhibition | |
|--------|-----------------------|-----------------|------------|
| | | Sample | Standard |
| 1 | 25 | 6.88±0.02 | 9.79±0.01 |
| 2 | 50 | 33.88±0.01 | 13.54±0.02 |
| 3 | 100 | 34.98±0.03 | 44.86±0.05 |
| 4 | 200 | 36.08±0.1 | 87.78±0.01 |

Figure 2. Anti-inflammatory Activity of *Premna rajendranii*



inhibition of α -amylase. The concentrations of samples used for testing the inhibitory activity are 25, 50, 75 and 100 (μ g). The inhibition activity of test samples showed optimum activity when compared to commercial drug at different concentrations. The maximum inhibition showed at 100 μ g concentration was 79.3% and minimum inhibition showed at 25 μ g concentration was 44.82%. IC₅₀ value of sample (288.8) is higher than standard drug (246.7). Acarbose was used as standard drug (Table 2, Fig: 1). Protein denaturation is the major cause of Inflammation (Mizushima and Kobayashi, 1968) and it is a response to negative stimuli including injury, infection (Lumeng and Saltiel, 2011). Basically it was meant to destroy invading microorganisms, inactivate toxins, repairing and healing of injuries which might lead to life threatening hypersensitivity reactions. Inflammation is the major cause for several diseases including neurological, cardiovascular, intestinal, dental and renal disorders and also linked to diabetics, ageing, obesity, multiple sclerosis, pancreatitis, cancer (Sakat et al., 2010; Burns et al., 2001; Kuek et al., 2006; Grivennikov et al., 2010; Jenny, 2012; Hoque et al., 2012; Marchant et al., Wyss- Coray et al., 2012).

Prescribed medicines for inflammation are mainly non steroidal anti- inflammatory drugs. These drugs have certain side effects causing gastric bleeding, ulceration, renal failure. Medicinal plants have a prolonged history of use as a remedy for inflammation (Insel, 1996; Rang et al., 2007). *Premna serratifolia* and *Premna tomentosa* have been reported to possess anti-inflammatory activity (Alam et al., 1993; Habtemariam et al., 2015). Methanolic leaf extract of *P.rajendranii* has shown high inhibition than standard at 50 μ l i.e. 33.88 \pm 0.01. The inhibition activity of other test samples (25 μ l, 100 μ l, 200 μ l) showed moderate activity when compared to standard drug. Diclofenac was used as the standard drug.

CONCLUSION

Our study demonstrates that the *P.rajendranii* acts as an antidiabetic and anti-inflammatory agent. The antidiabetic and anti-inflammatory activities may be due to the presence of various

bioactive compounds especially flavonoids. More purification needs to be done and further research on *P.rajendranii* is necessary for isolating bioactive compounds and their mode of action.

ACKNOWLEDGEMENTS

The authors are thankful to Dr. S. John Britto S.J., Director, The Rapinat Herbarium and Centre for Molecular Systematics, Tiruchirappalli, and Dr. Anand Gideon for valuable support and providing Laboratory facilities.

REFERENCES

- Alam, M., Joy, S., Susan, T. and Ali, S.U., (1993). Anti-inflammatory activity of *Premna tomentosa* Willd. In albino rats. *Ancient science of life*, 13(1-2), p.185.
- Angiosperm Phylogeny Group, (2009). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society*, 161(2), pp.105-121.
- Bailes, B.K., (2002). Diabetes mellitus and its chronic complications. *AORN Journal*, 76(2), pp.265-282.
- Burns, J., Gardner, P.T., Matthews, D., Duthie, G.G., Lean, J. and Crozier, A., (2001). Extraction of phenolics and changes in antioxidant activity of red wines during vinification. *Journal of Agricultural and Food Chemistry*, 49(12), pp.5797-5808.
- Chitturi, S. and George, J., (2002). Hepatotoxicity of commonly used drugs: nonsteroidal anti-inflammatory drugs, antihypertensives, antidiabetic agents, anticonvulsants, lipid-lowering agents, psychotropic drugs. In *Seminars in liver disease* (Vol. 22, No. 02, pp. 169-184). Copyright© 2002 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel.:+ 1 (212) 584-4662.
- Francis, S., Gideon, V.A., Britto, S.J., Dessy, V.J., (2019). A Review on Two Endemic Species of Genus *Premna* and their Conservation Importance, *Journal of Drug Delivery and Therapeutics*, 9(4- s), pp.666-669.
- Grivennikov, S.I., Greten, F.R. and Karin, M., (2010). Immunity, inflammation, and cancer. *Cell*, 140(6), pp.883-899.
- Jenny, N.S., (2012). Inflammation in aging:

- cause, effect, or both?. *Discovery medicine*, 13(73), pp.451-460.
- Joshi, V.K., Joshi, A. and Dhiman, K.S., (2017). The Ayurvedic Pharmacopoeia of India, development and perspectives. *Journal of ethnopharmacology*, 197, pp.32-38.
- Habtemariam, S. and Varghese, G.K., (2015). A Novel Diterpene Skeleton: Identification of a highly aromatic, cytotoxic and antioxidant 5-methyl-10-demethyl-abietane-type diterpene from *Premna serratifolia*. *Phytotherapy Research*, 29(1), pp.80-85.
- Hoque, R., Malik, A., Gorelick, F. and Mehal, W., (2012). The sterile inflammatory response in acute pancreatitis. *Pancreas*, 41(3), p.353.
- Indrianingsih, A.W., Tachibana, S. and Itoh, K., (2015). In vitro evaluation of antioxidant and α -Glucosidase inhibitory assay of several tropical and subtropical plants. *Procedia Environmental Sciences*, 28, pp.639-648.
- Insel, P. A (1996). Analgesic-antipyretic and anti-inflammatory agents and drugs employed in the treatment of gout. *The Pharmacological Basics of Therapeutics*, 9, pp. 617-657.
- Keerthana, G., Kalaivani, M.K. and Sumathy, A., (2013). In-vitro α amylase inhibitory and anti-oxidant activities of ethanolic leaf extract of *Croton bonplandianum*. *Asian Journal of Pharmaceutical and Clinical Research*, 6(4), pp. 32-36.
- Kesari, A.N., Kesari, S., Singh, S.K., Gupta, R.K. and Watal, G., (2007). Studies on the glycemic and lipidemic effect of *Murraya koenigii* in experimental animals. *Journal of Ethnopharmacology*, 112(2), pp.305-311.
- Khandelwal, K.R., (2008). *Practical Pharmacognosy* Nirali Prakashan. Pune, India, 19, pp.49-70.
- Kokate, C.K., Purohit, A.P. and Gokhale, S.B. (2001). Carbohydrate and derived Products, drugs containing glycosides, drugs containing tannins, lipids and protein alkaloids. *Text Book of Pharmacognosy*, Nirali Prakashan, Pune.
- Kuek, A., Hazleman, B.L., Gaston, J.H. and Östör, A.J., (2006). Successful treatment of refractory polyarticular juvenile idiopathic arthritis with rituximab. *Rheumatology*. 45(11), pp.1448-9.
- Kumar, K.M.P., Sunilkumar, T., Sreeraj, V., Thomas, B., Balachandran, I. and Antony, V.T., (2013). A new species of the genus *Premna* L. (Lamiaceae) from Western Ghats of India. *Webbia*, 68(2), pp.127-131.
- Lumeng, C.N. and Saltiel, A.R., (2011). Inflammatory links between obesity and metabolic disease. *The Journal of clinical investigation*, 121(6), pp.2111-2117.
- Marchant, D.J., Boyd, J.H., Lin, D.C., Granville, D.J., Garmaroudi, F.S. and McManus, B.M., (2012). Inflammation in myocardial diseases. *Circulation Research*, 110(1), pp.126-144.
- Mizushima, Y. and Kobayashi, M., (1968). Interaction of anti-inflammatory drugs with serum proteins, especially with some biologically active proteins. *Journal of Pharmacy and Pharmacology*, 20(3), pp.169-173.
- Modak, M., Dixit, P., Londhe, J., Ghaskadbi, S. and Devasagayam, T.P.A., (2007). Recent advances in Indian herbal drug research guest editor: Thomas Paul Asir Devasagayam Indian herbs and herbal drugs used for the treatment of diabetes. *Journal of clinical Biochemistry and Nutrition*, 40(3), pp.163-173.
- Peach, K., Tracey, M.V., (1956). *Modern methods of plant analysis*. Berlin: Springer Verlag
- Prabhu, S., Vinodhini, S., Elanchezhian, C. and Rajeswari, D., (2018). Evaluation of antidiabetic activity of biologically synthesized silver nanoparticles using *Pouteria sapota* in streptozotocin-induced diabetic rats. *Journal of Diabetes*, 10(1), pp.28-42.
- Rang, H.P., Dale, M.M., Ritter, J.M., Moore, P.K., (2007). *Pharmacology*. Elsevier publication, Churchill Livingstone, 6, pp.226-232.
- Rekha, K., Richa, P.K., Babu, S. and Rao, M., (2015). A phytochemistry of the genus *Premna*: a review. *International Journal of Pharmaceutical and Chemical Sciences*, 4(3), pp.317-25.
- Sakat, S., Juvekar, A.R. and Gambhire, M.N. (2010). In vitro antioxidant and anti-inflammatory activity of methanol extract of *Oxalis corniculata* Linn. *International Journal of Pharmacy and Pharmaceutical Science*. 2(1), pp.146-155.
- Treare, G.E, Evans, W.C. (1985), *Pharmacognosy* 17th edn. Bahiv Tinal, London, 149.
- Wyss-Coray, T. and Rogers, J., (2012). Inflammation in Alzheimer disease—a brief review of the basic science and clinical literature. *Cold Spring Harbor Perspectives in Medicine*, 2(1), p.a006346.