

## Biochemical evaluation of chlorophyll content using different solvents in various plant species of Amravati, Maharashtra (India)

N.A. Wagay<sup>\*1</sup>, M.A. Dar<sup>1</sup>, T.S. Per<sup>2</sup> and Shah Rafiq<sup>3</sup>

<sup>1</sup>Botany Research Laboratory, Vidyabharati Mahavidyalya College, Camp, Amravati, M.S., India

<sup>2</sup>Department of Botany, Government Degree College Doda, J & K, India

<sup>3</sup>Plant Tissue Culture Laboratory, Department of Botany, University of Kashmir, J&K, India

### ABSTRACT

Plants act as transducers of the solar light in the chemical energy because they contain chlorophyll molecules and are of utmost importance because of their light absorbing property. These pigment molecules are helpful in the measuring plant productivity, maintaining photostasis, protecting from excess sunlight and are also indicators of phototoxicity, pollution and environmental stress. In the present study, 40 plant species of District Amravati were evaluated for their chlorophyll content by Arnon's method of spectrophotometry, using 80% Acetone and 95% Ethanol as solvents. In case of 80% Acetone maximum chlorophyll-*a* concentration was found in *Psidium guajava* and *Polyalthia longifolia* and in case of 95% Ethanol maximum level of chlorophyll-*a* concentrations were found in eight species; *Psidium guajava*, *Bauhinia purpurea*, *Moringa oleifera*, *Murraya paniculata*, *Plumeria rubra*, *Terminalia catappa*, *Peltophorum pterocarpum* and *Delonix regia*. 80% Acetone based samples showed more difference in concentrations of chlorophyll-*b* with respect to chlorophyll-*a*. The maximum concentration of chlorophyll-*b* was found in *Psidium guajava*, *Murraya paniculata* and *Bauhinia purpurea* and in 95% Ethanol based samples chlorophyll-*b* concentration was maximum in *Bauhinia purpurea*. On the basis of the observation of all the results it was evident that the maximum and minimum values of chlorophylls are same in both the solvents but overall the 95% Ethanol showed higher concentration of both chlorophyll-*a* & *b*.

**KEY WORDS:** AMRAVATI, CHLOROPHYLL CONTENT, SOLVENTS, SPECTROPHOTOMETER

**Article Information:** \*Corresponding Author: [nasir.wagay1989@rediffmail.com](mailto:nasir.wagay1989@rediffmail.com)

Received 15/03/2019 Accepted after revision 25/06/2019

Published: 30th June 2019 Pp- 523-530

This is an open access article under Creative Commons License,  
Published by Society for Science & Nature, Bhopal India.

Available at: <https://bbrc.in/>

Article DOI: <http://dx.doi.org/10.21786/bbrc/12.2/40>

## INTRODUCTION

The chlorophylls comprise a group of more than 50 tetrapyrrolic pigments with common structural elements and function (Scheer 2003). Pigments are functionally important molecules in photosynthetic organisms. They not only harvest the light energy necessary for carbon reduction but some serve to protect the organism from excess light. The balance of photosynthetic pigments is dynamic and contributes to the maintenance of photostasis within the cell (Huner *et al.* 1998). The ratio of chlorophyll-*a*, and chlorophyll-*b* in terrestrial plants has been used as an indicator of response to light or shade conditions (Porra, 1991; Vicas *et al.* 2010). The small proportion of chlorophyll-*a/b* is considered as sensitive biomarker of pollution and environmental stress (Tripathi and Gautam, 2007). The amount of extracted chlorophyll may provide information on the sensitivity of plants during cultivation and herbicide application, and even indicate the manner of phytotoxic activity of herbicides (Nikolić *et al.* 2007). In order to analyze and describe changes in the process of photosynthesis and detect stress in plants, various types of indicators of chlorophyll activity were used (Lichtenthaler, 1996). In tumor or cancer therapy chlorophyll or chlorophyll derivatives can be utilized as a photodynamic agent (Brandis *et al.* 2006). It can be studied, modified and synthesized in chemistry and physics disciplines for different applications i.e., electronic, photophysics, optoelectronic, electrochemistry etc (Nurhayati and Suendo, 2011).

The absorbance properties of pigments facilitate the qualitative and quantitative analysis of these molecules. Determination of the content of photosynthetic pigments in leaves is one of the key techniques in studying the process of photosynthesis and measuring plant productivity. Chlorophyll molecules absorb and re-emit light, a characteristic which was the basis for developing two basic methods: absorption and fluorescence monitoring of the optical activity of chlorophyll molecules. The method of chlorophyll fluorescence is used for monitoring photosynthesis *in situ* and *in vivo* and for estimating the impact of various stress factors (abiotic, biotic, xenobiotic) on this crucial process. It makes possible to differentiate plant genotypes resistant to the aforementioned stressful environmental factors, and also to assess the positive impact of various agricultural measures on plant health (mineral nutrition, use of herbicides, etc.). Methods for non-destructive quantification of chlorophyll in plant leaves are particularly valuable in assessing nitrogen content in plants because chlorophyll is one of the most important points of its accumulation, (Indira *et al.*, 2015).


The confirmation of the two forms of chlorophyll was practically confirmed by Tswett 1903 (Nešković *et al.*

2003). A hundred years later, we used spectrophotometry to analyze plant pigments extracted in various solvents. There is a trade-off between choosing the best solvent for efficient quantitative extraction of chlorophylls and use of a solvent best suited for spectrophotometric assay (James and Akaranta, 2011). The selection of solvent for extraction brings about a dilemma. A number of factors may affect the activity of a solvent: the time required for extraction, amount of plant material, percentage of moisture in plant material, preservation of extract in unchanged form, (Moran and Porath, 1980; Jelić *et al.* 1992), as well as the fact that the extraction of chlorophyll-*a* is a slower process than that of chlorophyll-*b*. Light, an important environmental factor, causes degradation of chlorophyll, so that extraction should be carried out in almost total absence of light (Jelić *et al.* 1992).

The absorbance of the chlorophylls is then quantitatively determined by spectrophotometry at the wavelengths of maximum chlorophyll-*a* and chlorophyll-*b* absorption i.e.,  $\lambda = 647$  nm and  $\lambda = 664$  nm (Moran and Porath, 1980), while the actual content of photosynthetic pigments is calculated according to Wellburn's formulas (1994). This procedure is based on the Lambert-Beer law on linear relationship between absorbance and concentration of pigments within a certain range. To understand the variation in chlorophyll-*a*, chlorophyll-*b* and total chlorophyll content using different solvents, forty native species of tropical region were selected. Conclusions were drawn based on the variation in chlorophyll content values.

## MATERIALS AND METHODS

Amravati is a district in the state of Maharashtra situated at 20°55'33" N and 77° 45'53" E at 343m (1,125ft.) asl. The Amravati district has an area of 270 km<sup>2</sup>. The study area has well demarcated four seasons as a hot summer, heavily raining monsoon, a brief autumn and a mild winter. The area has sub tropical and deciduous climatic conditions with ample rainfall in the monsoon resulting in a rich diversity of vascular plants (Figures 1, 2, and 3). 40 commonly grown plant species of Amravati district (viz., *Aegle marmelos*, *Acacia nilotica*, *Alstonia scholaris*, *Annona reticulate*, *Annona squamosa*, *Artocarpus heterophyllus*, *Anacardium occidentale*, *Azadirachta indica*, *Butea monosperma*, *Bauhinia purpurea*, *Buchanania lazan*, *Cassia siamea*, *Callistemon lanceolatus*, *Cinnamomum tamala*, *Citrus aurantium*, *Citrus limon*, *Dalbergiasisoo*, *Delonix regia*, *Eucalyptus globules*, *Ficus benghalensis*, *Ficus religiosa*, *Mangnifera indica*, *Moringa oleifera*, *Murraya paniculata*, *Murraya koenigii*, *Manilkara zapota*, *Peltophorum pterocarpum*,



*Plumeria rubra*, *Pongamia pinnata*, *Psidium guajava*, *Polyalthia longifolia*, *Saraca indica*, *Santalum album*, *Syzygium cummini*, *Tectona grandis*, *Terminalia catappa*, *Tecoma stans*, *Thevetia peruviana*, *Tamarindus indica*, *Zizyphus mauritiana*) were used for experimental purpose. The understudy plants were morphologically identified with the help of standard floras i.e., Flora of British India (Hooker, 1876); and in Maharashtra collected and recorded by Cooke (1967); Naik (1998); and Singh and Karthekeyan (2001) and authenticated by taxonomist Professor Dr. S.P. Rothe. These species are mostly preferred to grow in tropical regions. Healthy and uninfected species were collected at their stage of maturity; and care was also taken during sampling of leaves to avoid mechanical injuries. Fresh leaf samples were washed thoroughly first in tap water followed by distilled water in the laboratory, kept to dry in room temperature (18°C) and analyzed for the determination of chlorophylls (chlorophyll-*a* and -*b*).

For the estimation of chlorophyll content the procedure given by Arnon (1949) with slight modifications was used. Leaves were cut into small pieces and major veins and tough fibrous tissue was discarded. 100 mg of material was used for grinding. 10 ml of 80% acetone (acetone:water; 80 : 20 v:v) was added. For complete pulverization of tissue few grains of sand was added. The homogenate was filtered through filter paper and retentate was discarded while the extract (filtrate) was

collected in the test tube. Chlorophyll concentration was determined using spectrophotometer/ colorimeter. In this study, leaves of 40 different tree species commonly available in understudy region of Maharashtra were used for the quantification of chlorophyll content. Two different solvents 80% Acetone and 95% Ethanol were used separately for the extraction of chlorophyll. Absorbance readings of chlorophyll extracts were measured at two different wavelengths 645nm and 663nm respectively.



## RESULTS AND DISCUSSION

Based on the absorbance value, calculations were made using Arnon's (1949) equation and the amount of chlorophyll-*a*, chlorophyll-*b* and total chlorophyll were estimated and tabulated. The results of Chlorophyll-*a*, chlorophyll-*b* and chlorophyll-*a+b* concentrations using 80% acetone as solvent are shown in Tabulated form (Table 1) and as well as in Graphical form (Graph 1). The maximum chlorophyll-*a* concentration was found in *Psidium guajava* and *Polyalthia longifolia* i.e., 0.03g/kg and minimum concentration was found in *Alstonia scholars* i.e., 0.0024 g/kg. In most of the species maximum concentrations of chlorophyll-*a* was found in the range between 0.005-0.02 g/kg. Concentration of chlorophyll-*b* showed much variation in comparison to concentration of chlorophyll-*a*. The maximum concentration of chlorophyll-*b* was found in *Psidium guajava*, *Murraya paniculata* and *Bauhinia purpurea* which was more than 0.03g/kg and the minimum concentration was found in *Alstonia scholaris* about 0.002 g/kg. Most of the concentrations were found in the range between 0.1-0.3 g/kg or more specifically between 0.1-0.2 g/kg. The maximum level of chlorophyll-*a+b* found in *Psidium guajava* was about 0.06 g/kg and the minimum concentration was found in *Alstonia scholaris*, 0.005g/kg. In most of the species chlorophyll-*a+b* concentration was found in the range between 0.02-0.04 g/kg. About nine species showed concentration of 0.02 g/kg and ten species showed concentration above 0.04g/kg.

The results of Chlorophyll-*a*, chlorophyll-*b* and chlorophyll-*a+b* concentrations using 95% Ethanol as solvent are depicted in Graph 2 and in Table 2. The maximum level of chlorophyll-*a* concentrations were found in eight species such as (*Psidium guajava*, *Bauhinia purpurea*, *Moringa oleifera*, *Murraya paniculata*, *Plumeria rubra*, *Terminalia catappa*, *Peltophorum pterocarpum*, *Delonix regia*) in similar amounts 0.03 g/kg, and minimum level of chlorophyll-*a* was found in *Syzygium cumini*, 0.005 g/kg. In 12 species less than 0.015 g/kg concentrations were found. The remaining species showed the concentration more than 0.015 g/kg. The chlorophyll-*b* concentration was the maximum in *Bauhinia purpurea*, 0.039g/

Table 1. Chlorophyll-*a*, chlorophyll-*b* and chlorophyll-*a+b* concentrations using 80% acetone

S. No	Name of plant species	Chl- <i>a</i>	Chl- <i>b</i>	Chl- <i>a+b</i>
1	<i>Annona reticulata</i>	0.0085	0.0071	0.0157
2	<i>Annona squamosa</i>	0.0265	0.0195	0.0461
3	<i>Artocarpus heterophyllus</i>	0.0155	0.0250	0.0405
4	<i>Bauhinia purpurea</i>	0.0195	0.0325	0.0521
5	<i>Butea monosperma</i>	0.0131	0.0134	0.0265
6	<i>Cassia siamea</i>	0.0182	0.0133	0.0315
7	<i>Cinnamomum tamala</i>	0.0194	0.0228	0.0423
8	<i>Citrus limon</i>	0.0264	0.0165	0.0429
9	<i>Citrus aurantium</i>	0.0142	0.0110	0.0253
10	<i>Dalbergia sisoo</i>	0.0053	0.0047	0.0101
11	<i>Delonix regia</i>	0.0205	0.0159	0.0365
12	<i>Ficus benghalensis</i>	0.0174	0.0116	0.0291
13	<i>Ficus religiosa</i>	0.0121	0.0108	0.0229
14	<i>Mangifera indica</i>	0.0075	0.0076	0.0152
15	<i>Azadirachta indica</i>	0.0187	0.0102	0.0290
16	<i>Moringa oleifera</i>	0.0193	0.0264	0.0457
17	<i>Murraya koenigii</i>	0.0089	0.0087	0.0177
18	<i>Murraya paniculata</i>	0.0190	0.0328	0.0518
19	<i>Plumeria rubra</i>	0.0125	0.0122	0.0248
20	<i>Santalum album</i>	0.0189	0.0261	0.0450
21	<i>Syzygium cumini</i>	0.0103	0.0151	0.0255
22	<i>Tectona grandis</i>	0.0188	0.0252	0.0440
23	<i>Zizyphus mauritiana</i>	0.0215	0.0142	0.0358
24	<i>Tecoma stans</i>	0.0185	0.0230	0.0415
25	<i>Buchanan lanzan</i>	0.0153	0.0144	0.0298
26	<i>Anacardium occidentale</i>	0.0166	0.0193	0.0360
27	<i>Polyalthia longifolia</i>	0.0309	0.0275	0.0585
28	<i>Acacia nilotia</i>	0.0186	0.0222	0.0409
29	<i>Aegle marmelos</i>	0.0158	0.0136	0.0295
30	<i>Tamarindus indica</i>	0.0161	0.0212	0.0373
31	<i>Pongamia pinnata</i>	0.0105	0.0127	0.0233
32	<i>Psidium guajava</i>	0.0313	0.0368	0.0681
33	<i>Saraca indica</i>	0.0059	0.0060	0.0120
34	<i>Terminalia catappa</i>	0.0087	0.0075	0.0163
35	<i>Callistemon lanceolatus</i>	0.0105	0.0110	0.0215
36	<i>Eucalyptus globules</i>	0.0162	0.0216	0.0378
37	<i>Manilkara zapota</i>	0.0132	0.0168	0.0300
38	<i>Alistonia scholaris</i>	0.0024	0.0027	0.0051
39	<i>Thevetia peruviana</i>	0.0094	0.0073	0.0168
40	<i>Peltophorum pterocarpum</i>	0.0124	0.0172	0.0296



kg and minimum concentrations were found in about 6 species in the same amounts i.e., between 0.007-0.009 g/kg. In maximum species, concentrations were found in range of 0.03 g/kg. In ten species concentrations were found above 0.03 g/kg. The maximum level concentration of chlorophyll-*a+b* was found in *Bauhinia purpurea* i.e., 0.071g/kg and the minimum chlorophyll concentration was found in *Syzygium cumini* i.e., 0.013g/kg. In 7 species the concentrations were found more than 0.06 g/kg and in 12 species concentrations were found more than 0.05 g/kg. About 93% of species showed more than 0.02 g/kg chlorophyll- *a+b* concentration.

A graphical representation of comparison between chlorophyll *a+b* concentrations in these two different solvents; 80% Acetone and 95% Ethanol showed different results (Graph 3). Samples prepared using 95% Ethanol showed more concentrations of chlorophyll- *a+b* in most of the species. Only 8 species of 80% Acetone prepared samples showed slightly more chlorophyll- *a+b* concentration than 95% Ethanol prepared samples that too with minor differences. 95% Ethanol prepared samples showed very higher concentrations than 80% Acetone prepared samples in more than 25% species. In about 10 species both the solvents showed the same amount


of chlorophyll- *a+b* concentration. *Alstonia scholaris*, *Terminalia catappa* were the two species which showed a huge difference of chlorophyll *a+b* concentrations in two solvents, in which 95% Ethanol showed the higher concentration then 80% Acetone.

As chlorophyll is one of the important attributes, the observations of chlorophyll-*a*, *b* and total chlorophyll-*a+b* in all the 40 species have been estimated and compared. A graphical representation on the basis of results have also been plotted in order to understand the variations and differences among different concentrations observed. Different solvents such as ethanol, acetone, DMSO etc, have been used by researchers from time to time, to estimate the chlorophyll concentration in plants. In the present study two solvents have been used: 80% Acetone and 95% Ethanol. The observations were taken using UV spectrophotometry at 663nm and 645nm and a variety of results were obtained.

In 80% Acetone maximum chlorophyll-*a* concentration was found in *Psidium guajava* and *Polyalthia longifolia* 0.03g/kg and minimum concentration was found in *Alstonia scholaris* 0.0024 g/kg and in 95% Ethanol maximum level of chlorophyll-*a* concentrations were found in eight species such as (*Psidium guajava*,

Table 2. Chlorophyll-*a*, chlorophyll-*b* and chlorophyll-*a+b* concentrations using 95% Ethanol

S. No	Name of plant species	Chl- <i>a</i>	Chl- <i>b</i>	Chl- <i>a+b</i>
1	<i>Annona reticulata</i>	0.0138	0.0115	0.0254
2	<i>Annona squamosa</i>	0.0241	0.0309	0.0550
3	<i>Artocarpus heterophyllus</i>	0.0130	0.0221	0.0352
4	<i>Bauhinia purpurea</i>	0.0312	0.0398	0.0711
5	<i>Butea monosperma</i>	0.0240	0.0341	0.0582
6	<i>Cassia siamea</i>	0.0228	0.0153	0.0381
7	<i>Cinnamomum tamala</i>	0.0187	0.0245	0.0432
8	<i>Citrus limon</i>	0.0127	0.0083	0.0210
9	<i>Citrus aurantium</i>	0.0159	0.0125	0.0284
10	<i>Dalbergia sisoo</i>	0.0137	0.0099	0.0236
11	<i>Delonix regia</i>	0.0313	0.0363	0.0676
12	<i>Ficus benghalensis</i>	0.0224	0.0152	0.0377
13	<i>Ficus religiosa</i>	0.0166	0.0119	0.0285
14	<i>Mangifera indica</i>	0.0120	0.0137	0.0253
15	<i>Azadirachta indica</i>	0.0112	0.0078	0.0191
16	<i>Moringa oleifera</i>	0.0312	0.0318	0.0630
17	<i>Murraya koenigii</i>	0.0149	0.0197	0.0347
18	<i>Murraya paniculata</i>	0.0314	0.0278	0.0592
19	<i>Plumeria rubra</i>	0.0314	0.0276	0.0590
20	<i>Santalum album</i>	0.0286	0.0218	0.0504
21	<i>Syzygium cumini</i>	0.0059	0.0071	0.0130
22	<i>Tectona grandis</i>	0.0219	0.0221	0.0441
23	<i>Zizyphus mauritiana</i>	0.0137	0.0138	0.0275
24	<i>Tecoma stans</i>	0.0237	0.0215	0.0452
25	<i>Buchanan lanzan</i>	0.0194	0.0153	0.0348
26	<i>Anacardium occidentale</i>	0.0300	0.0348	0.0649
27	<i>Polyalthia longifolia</i>	0.0253	0.0175	0.0429
28	<i>Acacia nilotia</i>	0.0270	0.0176	0.0446
29	<i>Aegle marmelos</i>	0.0132	0.0089	0.0221
30	<i>Tamarindus indica</i>	0.0234	0.0272	0.0507
31	<i>Pongamia pinnata</i>	0.0154	0.0160	0.0314
32	<i>Psidium guajava</i>	0.0313	0.0348	0.0661
33	<i>Saraca indica</i>	0.0094	0.0122	0.0216
34	<i>Terminalia catappa</i>	0.0313	0.0318	0.0631
35	<i>Callistemon lanceolatus</i>	0.0104	0.0087	0.0191
36	<i>Eucalyptus globules</i>	0.0273	0.0204	0.0478
37	<i>Manilkara zapota</i>	0.0228	0.0259	0.0487
38	<i>Alistonia scholaris</i>	0.0240	0.0333	0.0573
39	<i>Thevetia peruviana</i>	0.0281	0.0196	0.0477
40	<i>Peltophorum pterocarpum</i>	0.0313	0.0313	0.0627



*Bauhinia purpurea*, *Moringa oleifera*, *Murraya paniculata*, *Plumeria rubra*, *Terminalia catappa*, *Peltophorum pterocarpum*, *Delonix regia*) in similar amounts 0.03 g/kg, and minimum level of chlorophyll-*a* was found in *Syzygium cumini* 0.005 g/kg. In both the solvents chlorophyll-*a* concentration was found to be almost same. 80% Acetone based samples showed more difference in concentrations of chlorophyll-*b* with respect to chlorophyll-*a*. The maximum concentration of chlorophyll-*b* was found in *Psidium guajava*, *Murraya paniculata* and *Bauhinia purpurea* which were more than 0.03g/kg and the minimum concentration was found in *Alstonia scholaris* about 0.002 g/kg and in 95% Ethanol based samples chlorophyll-*b* concentration was maximum in *Bauhinia purpurea* 0.039g/kg and minimum concentrations were found in about 6 species in the same amounts in the range between 0.007-0.009 g/kg. *Alstonia scholaris* showed the least concentration of chlorophyll-*a* & *b* in 80% Acetone solvent.

On the basis of the observation of all the results it is evident that the maximum and minimum values of chlorophylls are same in both the solvents but overall the 95% Ethanol showed higher concentration of both chlorophyll-*a* and -*b*. Similar types of studies were also done by José Francisco *et al.*, (2008) in which they

determined chlorophyll concentrations in tropical tree species by Portable Chlorophyll Meter with appropriate adjustment equations. Faisal and Anis *et al.*, (2006) reported higher amount of chlorophyll-*a* ( $0.91 \pm 0.19$  mg/g FW) and chlorophyll-*b* ( $0.61 \pm 0.09$  mg/g FW) in micro propagated plants of *Psoralea corylifolia* compared to chlorophyll-*a* ( $0.83 \pm 0.31$  mg/g FW) and chlorophyll-*b* ( $0.53 \pm 0.14$  mg/g FW) in seedlings. Dere (1998) have also investigated the level of chlorophyll-*a* in fresh water forms of some algal species. Indira *et al.*, (2015) have also estimated the chlorophyll content of *Tridax procumbens* grown in normal and polluted region in which they reported that the chlorophyll content in normal and polluted regions is 2.99mg/g and 2.56 mg/g respectively.

This method of chlorophyll quantification is reliable but time consuming and requires great precision. The main disadvantage of the method is that the process of extraction can result in erroneous qualitative and quantitative determination of the content of pigments (due to photochemical reactions, impact of ambient oxygen, chlorophyllase activity, pheophytinization caused by acids from plant tissue, etc (Jelić *et al.*, 1992; Wellburn, 1994). Difficulties in comparing the results obtained by different extraction techniques sometimes raise the question of validity of research conclusions. A particular problem is posed by the fact that the amounts of chlorophyll measured after extraction with various solvents are hard to compare because different formulae are used for content calculation (Lichtenthaler, 1988). The defined absorption coefficients in these formulae are based on measurements made with outdated or imprecise spectrophotometers that are still in use. Therefore, the results obtained by different groups of researchers may differ, even when using the same extraction solvents, and be incomparable for several reasons: (I) differences in spectrophotometer resolutions in the range of red light wavelengths (II) the accuracy of readings of selected wavelengths, and (III) water content in analyzed tissues (Jelić *et al.*, 1992).

## CONCLUSION

Chlorophyll from 40 different tree species was extracted and estimated. Considering the results obtained in this work, chlorophyll content in *Psidium guajava* leaves was higher and almost similar in both the solvents with less variation followed by *Bauhinia purpurea*, *Murraya paniculata*, *Annona squamosa*, *Polyalthia longifolia*, *Acacia nilotica*, *Cinnamomum tamala*. Species like *Delonix regia*, *Anacardium occidentale*, *Terminalia catappa*, *Alstonia scholaris*, *Thevetia peruviana*, *Peltophorum pterocarpum* showed very high difference in their concentration in 95% Ethanol and 80% Acetone, it suggests that different

plants may need different solvents for isolation; it may be because of their biochemical constituents while as some plants showed a similar concentration in both the solvents. Chlorophyll-*a*, chlorophyll-*b* and total chlorophyll amount showed different values in the present study varying from 0.01 g/kg to 0.08 g/kg in these understudy tree species. By this we can conclude that different plant species have different requirements of photosynthetic pigments which have a direct relation to the photosynthetic activity and the rate of photosynthesis.

## ACKNOWLEDGEMENTS

Authors are sincerely thankful to Professor S.P. Rothe (Ex-Head, Department of Botany, Shri Shivaji Science College, Akola) for his help in identification and authentication of several plant species. Authors are also thankful to Dr. V.R. Deshmukh (Ex-Head Department of Botany, Vidyabharati Mahavidyalaya College, Amravati) for providing lab facilities during this study.

## CONFLICT OF INTEREST

The authors declare no competing interests to any person, agency or institution.

## REFERENCES

Arnon D. (1949) Copper enzymes isolated chloroplasts, polyphenoloxidase in *Beta vulgaris*. Plant Physiology Vol. 24 : Pages 1-15.

Brandis, Salomon A.S.Y. and Scherz A. (2006) Chlorophyll sensitizers in photodynamic therapy. Advances in Photosynthesis and Respiration Vol. 25: Pages 461-483.

Cooke T (1967) The Flora of the Presidency of Bombay. Vol I & II. Botanical Survey of India, Calcutta.

Faisal M. and Anis M. (2006) Thidiazuron induced high frequency axillary shoot multiplication in *Psoralea corylifolia*. Biologia Plantarum. Vol. 50: Pages 437-440.

Hooker JD (1872-1897) (Repr. 1998) The Flora of British India. Vol. I - VII. London.

Huner N.P.A., Oquist G., Sarhan, F. (1998) Energy balance and acclimation to light and cold. Trends in Plant Science Vol. 3 Issue 6: Pages 224-2309.

Indira P., Shamshad A.S. and John P.M. (2015) The effect of air pollution on some biochemical factors of some plant species growing in Hyderabad. International Journal Pharma Bio Sciences Vol. 6: Pages 1349-1359.

James A.O. and Akaranta O. (2011) Inhibition of Corrosion of Zinc in hydrochloric acid solution by Red Onion Skin Acetone extract. Research Journal of Chemical Sciences Vol. 1 Issue 1: Pages 31-37.

Jelić G., Bogdanović M. and Nikolić B. (1992) Metode za određivanje fotosintetskih pigmenta. Acta herbologica. Vol. 1: Pages 235-242.

José Francisco de Carvalho Gonçalves, Ulysses Moreira dos Santos Junior e Emerson Alves da Silva. (2008) Evaluation of a portable chlorophyll meter to estimate chlorophyll concentrations in leaves of tropical wood species from Amazonian forest. Hoehnea. Vol. 35: Pages 185-188.

Lichtenthaler H.K. (1988) *In vivo* chlorophyll fluorescence as a tool for stress detection in plants. In H.K. Lichtenthaler (Ed.), Applications of Chlorophyll Fluorescence Pages 129-142.

Lichtenthaler H.K., Lang M., Sowinska M., Heisel F., Miede J.A. (1996) Detection of vegetation stress via a new high resolution fluorescence imaging system. Journal of Plant Physiology Vol. 148: Pages 599-612.

Moran R. and Porath D. (1980) Chlorophyll determination in intact tissues using N, N-dimethylformamide. Plant Physiology. Vol. 65 Issue 3: Pages 478-479.

Naik VN & Associates. (1998) Flora of Marathwada. Vol. 1 & 2, Amrut Prakashan, Aurangabad.

Nešković M., Konjević R., Čulafić L. (2003) Fiziologija biljaka (Plant Physiology). Beograd: NNK International.

Nikolić B., Drinić G., Jovanović V., Janjić V., Stojaković S. (2007) Different aspects of inhibition of growth and photosynthesis of maize (*Zea mays* L.) by the phosphonate herbicide sulphosate, 1: Root manipulation. Acta herbologica Vol. 16: Pages 29-39.

Nurhayati and Veinardi Suendo. (2011) Isolation of chlorophyll *a* from spinach leaves and modification of center ion with Zn<sup>2+</sup>: Study on its optical stability. Jurnal Matematika & Sains. Page 16.

Porra R.J. (1991) Recent advances and re-assessments in chlorophyll extraction and assay procedures for terrestrial, aquatic, and marine organisms, including recalcitrant algae. In: Scheer H (ed) Chlorophylls. Pages 31-57.

Scheer H. (2003) In: Beverley Green, Parson WW (2003) Light-harvesting antennas in photosynthesis. Springer Netherlands Page 29.

Singh NP, Lakshminarasimhan P, Karthikeyan S, Prasanna PV (2001) Flora of Maharashtra State, Dicotyledones. Vol. I & II. BSI Publications, Calcutta.

Sukran Dere, Tohit Gunes, Ridvan Sivaci. (1998) Spectrophotometric determination of chlorophyll - A, B and total carotenoid contents of some algae species using different solvents. Turkish Journal of Botany Pages 2213-2217.

Tripathi A.K. and Gautam M. (2007) Biochemical parameters of plants as indicators of air Pollution. Journal of Environmental Biology Vol. 28: Pages 127-132.

Vicas S.I., Laslo V., Pantea S., Bandict G.E. (2010) Chlorophyll and carotenoids pigments from Mistletoe (*Viscum album*) leaves using different solvents. Fascicula Vol. 2: Pages 213-218.

Wellburn A.R. (1994) The spectral determination of chlorophylls *a* and *b*, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. Journal of Plant Physiology Vol. 144: Pages 307-313.