

An Updated Review on the Spiders of Order Araneae from the Districts of Western Ghats of India

Misal P. K^{*1}, Bendre N. N², Pawar P. A³, Bhoite S. H⁴ and Deshpande V. Y⁵

^{1,2,3,4,5}*Department of Zoology, Yashwantrao Chavan Institute of Science, Satara.*

^{1,2,3,4,5}*Drongo-Nisarga Sanshodhan, Sanvardhanaani Sanrakshan Sanstha., Satara.*

ABSTRACT

Spiders belonging to order araneae are one of the biggest groups of predaceous organisms in the animal kingdom. Along with their diverse structure, they are popular for their tough silk and spider venom which is of pharmaceutical importance. Also they are playing a major role in lower food web in the ecosystem to maintain the ecological equilibrium and are one of the best bioindicators of natural ecosystems. Now a days, due to the habitat destruction caused by natural calamities and anthropogenic activities, the number of spiders species are found to be declined, as a result number of species of spiders are getting extinct before they are explored. One of the major hotspots of biodiversity of India is Western Ghats, known for their high species diversity. Taking into account the importance of spiders as bioindicators of environment, the present article was aimed to review on Araneae diversity from Satara Sangli and Kolhapur districts lying in the Northern Western Ghat regions of Maharashtra, India. This study shall help to record the number of species found in this regions. This review provides a checklist of total 27 families of 101 genera representing 178 species along with some new species recorded with special reference to their behaviour, habitat and variations among them. The review mainly provides a baseline information for the future study on diversity of spiders. From this review it is concluded that Northern Western Ghats being one of the biggest hotspots of diversity have a large number of spiders species, which are very less as compared to the spider diversity of India as suggested by the world catalogue of spiders, suggesting for an urgent need to explore further the areneae diversity in order to maintain the ecological equilibrium and the products obtained from them.

KEY WORDS: ARACHNIDA, ARANEA, SPIDER, BIODIVERSITY, WESTERN GHATS.

Article Information: *Corresponding Author: pkmspider@gmail.com

Received 15/10/2019 Accepted after revision 19/12/2019

Published: 30th Dec 2019 Pp-844-854

This is an open access article under Creative Commons License,

Published by Society for Science & Nature, Bhopal India.

Available at: <https://bbrc.in/>

Article DOI: <http://dx.doi.org/10.21786/bbrc/12.4/314>

INTRODUCTION

Spiders are the ancient Arthropods belonging to the class Arachnida having history of about 350 million years ago. These are the air breathing arthropods known to have special characteristics containing eight legs, two to eight eyes and spinnerets to spin the silk. They differ from class Insecta due to the absence of antennae and wings. Class Arachnida contains orders- Scorpiones, Schizomida, Amblypygi, Uropygi, Opiliones, Pseudoscorpiones, Palpigradi, Solifuge, Ricinulei, Acari and Araneae, (Tikader, 1982; Sebastian and Peter 2009). Spiders belonging to order Araneae differ from other orders of Arachnids by the presence of pedicel that joins the cephalothorax and the abdomen. Their special characteristic is the presence of spinnerets which secrete silk, this character differs them from the mites, which also secrete the silk by the glands in their palp as there are no spinnerets. Order Araneae constitutes the largest order, which ranks 7th in all over diversity of species. Spiders are very small to large sized arthropods with some special features within environment and ubiquitous in terrestrial habitat, (Turnbull, 1973; Preston, 1984; Nyfeller, 1987).


Arachnids are an important but generally poorly studied group of phylum Arthropoda. Many authors are making a lot of efforts to bring forward the research on spider biology and educate its importance to society. Many reviews have been

collected from international context which have helped in supporting research on spiders, which are meant to have precious role in ecology as they are mostly predatory in nature which help in reducing insect pests and invertebrate population in most ecosystems, (Jennings, 1986; Wise, 1993; Russell 1999; Maloney, 2003 and Plantnick 2019). Many researchers have conducted baseline studies involving role of spiders as biological, ecological indicators of natural ecosystem (McIver, 1990; Churchill, 1997; Maelfait, 1998) and in conservation planning, (Kremen, 1993). Allred (1969) and Allred (1976) also reported diversity of spiders in Arizona and Utah after new power plant installations in Nevada at the Nevada Nuclear test site to indicate any harmful changes in the ecosystem due to this plant.

Wise (1993) recorded importance of spiders in ecological web. Spiders are widely distributed group of predaceous organisms in the animal kingdom, (Riechert, 1984). Knowing the importance of presence of diverse Araneo fauna and its role in the ecosystem, the present review was aimed on the spider diversity of Satara, Sangli, Kolhapur districts which are the important locations of Western Ghats of Maharashtra known for their varied and rich biodiversity. According to World Spider Catalogue (Version 20.5) by Plantnick (2019), there are 48,334 species belonging to 4,143 genus representing 120 families. The pioneer contribution in Arachnology of Indian Spiders was given by European arachnologist, Stoliczka (1869).

Earliest contribution in arachnology was given by Blackwell (1867), Karsh (1873), Simon (1887), Thorell (1895), Pocock (1900), Malhotra (1980) and Tikader (1980, 1982). Gajbe (1978-2009) gave precious contributions on the Spiders of India. According to Indian Spiders checklist given by Siliwal (2005), 1442 species belonging to 361 genus of 59 families have been recorded and in 2012, Keswani recorded 1685 species of 438 genus belonging to 60 families, out of them, 91 species are of Infraorder Mygalomorphae and 1595 species of Araneomorphae. As per Tikader (1974), knowledge on diversity and distribution of spiders in Maharashtra is sparse, as compared to other regions of the world. Gazetteer of India which

Figure 1. Spider- Dorsal and ventral view



included 90 species, 14 families. Hippargi (2011b) recorded 19 families from Lonar, 25 families from Melghat and 31 families from Southern Tropical thorn forest, Solapur. From Western Ghats, Sebastian (2012) recorded 275 species belonging 139 genus and 39 families. Wasankar (2016) recorded 37 species of spiders belonging to 25 genera representing 17 families from Buldhana. 21 species representing 19 genus of 13 families recorded by Gajbe (2016) from Nagpur. 71 species of 15 families by Maheshwari (2018), and 104 species of 18 Families by Deshmukh (2014) recorded from Satpura ranges. From Amravati 31 species of 8 genera of 1 family were recorded by Deshmukh (2018). From Vidarbha in cotton fields 70 species belonging to 29 genera of 7 families were recorded, and in orange fields total 64 species of 13 families were recorded by Deshmukh (2017). Gajbe (2016) recorded 13 species of 9 families from Chandrapur and from Akola 26 species were recorded by Shirbhate (2017) (Table No 2).

Looking at the scenario of diversity of spiders from Maharashtra, a data from 2012-2019 was compared with India (Keswani,2012) which suggests that a lot of species still have to be explored on large scale before any natural calamities or any other environmental hazards make any species extinct without being explored Fig No.4. So this review has been carried out on to present the current status of spider diversity in three regions of Maharashtra, Satara, Sangli and Kolhapur which are the main parts of Northern Western Ghats known for one of the hottest biodiversity spots of India, [Fig No.2]. Along

with their importance as bioindicators they are also known popularly for Spider venom, which plays precious role in pharmacological extracts, by preying larva and adults of mosquito which carry malarial parasite (Choi, 2004; Ndava,2018). Spiders contain venom which is rich in proteins, containing disulfide peptides which have affinity and specificity towards specific subtype of ion channels and receptors. According to a report by Honor,(2015) only 0.01 % of its enormous pharmacological component has been explored till now. This discovery has accelerated the hope of development of novel molecules from such animal sources for drug discovery. This has also been innovatively used in medical,pharmaceuticals, and therapeutics, as a non polluting biopesticide, (Bode, 2001; Fry, 2009; Saez,2010; Herzig,2011;Windley, 2012; King and Hardy 2013; Mammola,2017; Li, 2017; Pineda,2018; Halarhkar,2018 and Luddecke,2019)

Recently various non chemical control methods are used in integrated pest management. Spiders in the rice field control the population of plant hoppers and leafhoppers, also in the cotton fields. And it is observed that web weavers controls the population of adult white fly and hunters control the population of nymph of pests in the field, without damaging plants. Thus they maintain the equilibrium in the ecosystem by playing major role in lower food web in the ecosystem, (Hamamura,1969; Sasaba,1973; Gavarra,1973; Samal, 1975; Kobayashi,1977; Chiu,1979;Vanden,1982; Holt,1987; Tanaka,1989; Jeyaparvathi, 2013; Lee;

Figure 2. Map showing three districts of Maharashtra-Sangli,Satara,Kolhapur(part of northern western ghats)



Table 1. List of Family wise Distribution of Spiders

Sr.No	Name of Family	Genus
01	Araneidae	i)Araneus ii)Argiope iii) Cyclosa iv)Gasteracantha v)Eriovixia vi)Larinia vii)Neoscona ix) Nephila x)Parawixia xi) Plebs xii)Guizygiella xiii) Cyrtophora xiv)Arachnura xv) Telecantha xvi)Polys
02	Salticidae	i)Aemonea ii)Bavia iii)Brettus iv) Currhotus v)Epeus vi)Epocilla vii) Hasarius viii) Hyllus ix)Menemerus x) Phintella xi)Marpissa xii)Myrmarachne xiii) Plexippus xiv)Portia xv) Rhene xvi)Telamonia xvii)Thiania
03	Lycosidae	i)Geolycosas ii)Hippasa iii) Lycosa iv)Pardosa v) Architosa vi)Evippa
04	Thomisidae	i)Amyciaea ii)Misumena iii)Oxytate iv) Ozyptila v)Runcinia vi) Thomisus vii)Xysticus viii)Tmarus
05	Sparassidae	i)Heteropoda ii)Olios
06	Theridiidae	i)Achaearanea ii)Argyodes iii) Chikunia iv)Enoplognatha v)Episinus vi) Euryopsis vii)Nesticodes viii) Phoroncidia ix)Phycosoma x) Rhomphaea xi)Steatoda xii)Theridion xiii)Theridula xiv)Latroedectus xv)Propostira
07	Oxyopidae	i)Oxyopes ii)Peucetia
08	Pholcidae	i)Artema ii)Crossopriza iii)Pholcus
09	Pisauridae	i)Pisaura ii)Nilus iii)Thalassius iv)Dolomedes v)Perenethis
10	Tetragnathidae	i)Leucauge ii)Opadometa iii)Tetragnatha
11	Gnaphosidae	i)Gnaphosa ii) Scotophaesis iii) Poecilochroa iv)Scopoides v)Zelotes
12	Eresidae	i)Stegodyphus
13	Corinnidae	i)Castianeira
14	Hersiliidae	i)Hersilia
15	Philodromidae	i)Philodromus ii)Tibellus iii)Tibilus
16	Ctenidae	i)Ctenus
17	Dipluridae	i)Diplura
18	Linyphiidae	i)Nerene
19	Euticharidae	i)Chieracanthium
20	Sicariidae	i)Loxosceles
21	Ulboridae	i)Uloborus Zosis
22	Zodariidae	i)Mallinella ii)Zodarion
23	Miturgidae	i)Cheiracanthium
24	Scytodidae	i)Scytodes
25	Clubionidae	i)Clubiona
26	Theraphosidae	ii)Unknown
27	Nephilidae	iii)Nephila
Total	27	101

Found species of spiders in study area -

- o *Araneus mitificus* , *Stegodyphus sarasinorum* , *Hyllussemicupreus*
- o *Argiope aemula* , *Argiope anasuja* , *Castianeira zetes*
- o *Cyclosa confrag* , *Neoscona bengalensis* , *Scytodes fusca*

Cotes, 2018; Tabasum, 2018 and Ndava, 2018). Spiders silk is used to explore evolution, biomaterial engineering, to check physical properties of protein fibres (Hinman, 2000; Heim, 2009; Mammola, 2017). Spiders are very careful about habitat changes. Their occurrence depends on the microclimatic characters, vegetation of biotic and abiotic factors. Their diversity is going to decrease due to habitat destruction, anthropogenic activities, environmental changes, cannibalism, territoriality, natural calamities hence they are on the verge of extinction, (Pinkus, 2006; Cardosa, 2010; Lee; Cardosa, 2017; Tabasum, 2018; Halarnkar, 2018). Taking into consideration the importance of spiders as biocontrol agents, bioindicators of ecosystem and therapeutic potential of Venom and economic importance of silk produced by them, it is a matter of concern to protect the diversity of spiders found in such area of varied biodiversity and pave the way to explore their pharmaceutical potential. The species level spider list used here is taken from literature, based

on i) modern literature ii) new species records with genitalia dissection and morphological, behavioural characters. All the valid Genus and species are summarised. The record of spiders of Satara, Sangli, Kolhapur has not yet been listed before so it was short of references. Total 27 families representing 101 genera belonging to 178 species have been recorded by authors in a diversity of papers. Some new species of spiders have been also recorded from Satara district. These families are present in every habitat. Infra-order Araneomorphae shows its dominance over Mygalomorphae. They show different types of web pattern. New species are recorded according to morphological and reproductive characters also the notes of its life stages, keys of identification, and their behavior were studied. The organisms morphology, habit, habitat, its life cycle study is crucial for its study and identification. More, (2010) recorded dominant families are Oxyopidae, Thomisidae, Araneidae, and Salticidae from the Radhanagari wildlife sanctuary, Chandoli national park, Koyana wildlife sanctuary.

Bendre, investigated total 11 species belonging to 6 families. Kulkarni, (2011) studied Ecology and behaviour of *Argiope* species during September to November 2010. Kulkarni, (2011) recorded new species of spider belonging to Family, Oxyopidae- *Oxyopes sataricus*. Kulkarni, (2012) recorded the spider *Arachnura angora* from Maharashtra. Kulkarni, (2012) recorded new flower spider




Figure 3. Spider anatomy and some families of order Araneae




Figure 4. Comparison of species wise diversity of spiders of year 2012 (Keswani) from India with three districts of Maharashtra till 2019

Table 2. Diversity of Spiders in Maharashtra from some areas-

Areas in Maharashtra	Study Year	Family	Genus	Species
Buldhana, Wasankar, (2016).	2014-2015	17	25	37
Nagpur, Gajbe, (2016).	2014-2015	13	19	21
Satpura Ranges, Deshmukh, (2014), Maheshwari, (2018).	2012-2013	18	-	104
Amravati, Deshmukh, (2018).	2014-2015	15	-	71
Vidarbha (Cotton fields and orange fields respectively), Deshmukh, (2017).	2017-2018	1	8	31
Chandrapur, Pawan, Gajbe, (2016).	2011-2012	7	29	70
Akola, Shirbhate, (2017).		13	32	64
	2012	9	12	13
	2015-2016	1	11	26

Thomisus sikkimensis with note on its courtship behaviour. Karanjkar in his minor project recorded 20 Families representing 40 Genus and 90 species from Satara Tehsil. Mirza, (2012) recorded a new species of the Mygalomorphae spiders belonging to family Idiopidae (trapdoor spiders). More, (2013) recorded 24 families representing 150 species of 78 Genera from Bamnoli region of Koyana wildlife Sanctuary during the study year 2011- 2013. Kulkarni, (2013) recorded 63 Genus of 25 Families and have given Generic checklist which showing Araneidae as a dominant family. Kulkarni, (2014) recorded new species *Tylorida satarensis*, with addition of field observations of that species. Kulkarni, (2015) described habit and conservation status of male *Tylorida satarensis*.

Kulkarni, (2015) recorded Genus *Siler Simon, 1889*. More (2015) recorded 90 species belonging to 54 genus from Zolambi region of Chandoli National Park. More (2015) recorded 18 families representing 61 species from Vakoba Devrai region of Radhanagari Wildlife Sanctuary. More (2015) investigated 16 families representing 58 species belonging to 38 genus from Rundiv, Sidheshwar and Ramnadi area of Chandoli national park. Lanka, (2015), investigated 18 families during year 2011-2013 and in that most dominant families are Araneidae representing 16 species and Salticidae representing 10 species. More (2016) recorded 18 families representing 70 species belonging to 48 genus from Dandoba hill forest. Lanka (2017), recorded 106 species belonging to 78 genus of 24 families from Radhanagari wildlife sanctuary. Kulkarni, (2017)

reported a new species of spiny spider *Meotipa sahyadri* belonging to family Theridiidae. Also he have given the phylogenetic analysis of that spider and relationships between *Meotipa* with *Chryso. Sarwade*, (April 2017) studied diversity and seasonal variation of spiders from Mhaishal lake and Brahmanath lake Khanderajuri, from Miraj Tehsil of Sangli. Depending on the habitat, climatic conditions, altitude and latitude diversity of spiders also changes. As they are shy creatures, hide from others. These primitive spiders are on the way of extinction and rarely found. They must be documented as they are less examined by the peoples.

ACKNOWLEDGMENTS

The authors are thankful to Sidharth Kulkarni, Ph.D student of Biological Sciences, George Washington University, Vishal Deshpande, Satara and Department of Zoology and Fisheries of Yashavantrao Chavan Institute Of Science, Satara for the guidance and support to carry out this research.

REFERENCES

- Allred, D.M. (1969) Spiders Of The National Reactor Testing Station. Great Basin Naturalist 29:105-108.
- Allred, D.M., Gertsch W.J.. (1976). Spiders And Scorpions From Northern Arizona And Northern Utah Journal Of Arachnology 3: 87-99.
- Bendre N.N., Shaikh A.A., And Sasane P.U., (2013) Spider Diversity In Botanical Garden Of Y.C. Institute Of Science, Satara; Proceedings

- Of National Conference On Environmental Biotechnology , ISBN- 978-81-8498-412-6.
- Blackwell, J. (1867);Description Of Seven New Species Of East Indian Spiders Recieved From The Rev. O P. Cambridge. *Annals And Magazine Of Natural History.* (3)14: 36-45.
- Bode F, Sachs F, Franz MR. (2001);Tarantula peptide inhibits atrial fibrillation. *Nature.*409(6816):35-36 DOI 10.1038/35051165.
- Cardoso, P. Arnedo, M A, Triantis, KA. (2010);Drivers of diversity in Macaronesian spiders and the role of species extinctions. *Journal of Biogeography.* 37(6):1034-1046.
- Chiu, S.C. (1979) ;Biological control of the brown planthopper. In: *Brown Planthopper, Threat to Rice Production in Asia.* International Rice Research Institute, Los Baños, Laguna, Philippines, pp. 335-355.
- Choi SJ, Parent R, Guillaume C, (2004), Isolation and characterization of Psalmopeotoxin I and II: Two novel antimalarial peptides from the venom of the tarantula *Psalmopoeus cambridgei*. *FEBS Lett.*2004; 572:109-117.
- Churchill, T. B. (1997);Spiders As Ecological Indicators: An Overview For Australia. *Memoirs Of The Museum Of Victoria.* 56 (2): 331-337.
- Coddington J A , Levi H W (1991);Systematics And Evolution Of Spiders (Araneae). *Ann Rev Ecol Syst.* 22:565-592.
- Cotes B, González M, Benítez E,(2018) ; Spider Communities and Biological Control in Native Habitats Surrounding Greenhouses. *Insects;*9(1):33.. doi:10.3390/insects9010033.
- Deshmukh U. S. , (2018); Araneidae (Orb Weavers) From Salbardi Forest Dist. Amravati, (Ms), India. *International Journal Of Current Engineering And Scientific Research (IjcesrIssn (Print): 2393-8374, (Online): 2394-0697, Volume-5, Issue-1, 2018.*
- Deshmukh U. S.,Raut N.M. ,(2014); Seasonal Diversity And Status Of Spiders (Arachnida: Araneae) In Salbardi Forest (Satpura Range), Maharashtra, India; *Journal Of Entomology And Zoology Studies;* 2 (6): 278-281, ISSN 2320-7078.
- Deshmukh U. S. (2017); Can Spiders Be The Biological Control Measure: A Survey Through Orange And Cotton Fields; *Indian J.Sci.Res.* 12 (2): 011-016, ISSN: 0976-2876 (Print) , ISSN: 2250-0138(Online).
- Lanka L.P,(2017) ; An Addition To Spider Fauna From The Vicinity Of Radhanagari Wildlife Sanctuary Of Kolhapur District; *International Journal Of Scientific Engineering And Research (IJSER) ISSN (Online): 2347-3878.*
- More S. (2015); Diversity Of Spiders From Zolambi Region Of Chandoli National Park; *IOSR Journal Of Pharmacy And Biological Sciences (IOSR-JPBS) E-ISSN: 2278-3008, P-ISSN:2319-7676. Volume 10, Issue 2 Ver. 1 (Mar -Apr. 2015), PP 30-33 .*
- Lanka L.P (2015); To Investigate Functional Assemblages Of Spiders From Western Ghat Ranges Of Radhanagari Wildlife Sanctuary Kolhapur, Maharashtra *EUROPEAN ACADEMIC RESEARCH Vol. III, Issue 4/ July 2015, ISSN 2286-4822*
- Fry, B.G., Roelants, K., Champagne, D.E.,(2009);The toxicogenomic multiverse: Convergent recruitment of proteins into animal venoms. *Annu. Rev. Genomics Hum. Genet.,* 10, 483-511. [CrossRef] [PubMed].
- Gajbe P.U.(2016); Record Of Some Spiders (Arachnida: Araneae) From Karhandla In Nagpur District, Maharashtra ; *Journal On New Biological Reports JNBR* 5(3) 133 - 138
- Gajbe P.U.,(2016); Record Of Some Spiders (Arachnida: Araneae) Found In Ghodazari, Chandrapur District, Maharashtra; *IJBAT, Special Issue-(6), ISSN 2347 - 517X (Online).*
- Gajbe, P. ,(2003); Checklists Of Spiders (Arachnid; Araneae) Of Madhya Pradesh And C Hattisgarh. *Zoos. Print Journal* 18 (10): 1223-1226.
- Gajbe, U. A. (1995a);Spiders Fauna Of Conservation Areas: Fauna Of Kanha Tiger Reserve, Madhya Pradesh. *Z00ological Survey Of India, Publication: 27-30.*
- Gajbe, U. A. (1995b);Spiders, Fauna Of Conservation Areas: Fauna Of Indravati Tiger Reserve, Madhya Pradesh. *Zoological Survey Of India, Publication: 53-56.*
- Gajbe, U. A. (1999);Studies On Some Spiders Of The Family Oxyopidae (Araneae: Arachnida) From India: Records Of Zoological Survey Of India 97(3): 31-79.ISSN 2319 - 1104.
- Gavarra, M., Raros R.S. , (1973); Studies on the biology of the predator wolf spider, *Lycosapseudoannulata* Böst. *Str.(Araneae: Lycosidae).* *Philippine Entomologist* 2, 6: 427.
- Halarnkar M H,(2018), Distribution, Diversity and Ecology of Spider Species At Two Different

- Habitats. *Int J EnvironSciNatRes*;8(5):555747. DOI: 10.19080/IJESNR.2018.08.555747.
- Hamamura, T. ,(1969) ;Seasonal fluctuation of spider populations in paddy fields. *Acta Arachnol.* 22, 2: 40-50.
- Heim M, Keerl D, Scheibel T. ,(2009); Spider silk: from soluble protein to extraordinary fiber. *Angewandte Chemie International Edition* 48(20):3584–3596 DOI 10.1002/anie.200803341.
- Herzig, V, Wood D.L.A., Newell F. ,(2011); 2.0, an updated online resource for spider toxin sequences and structures. *Nucl. Acids Res.*, 39, D653–D657. [CrossRef] [PubMed].
- Hinman MB, Jones JA, Lewis RV. ,(2000); Synthetic spider silk: a modular fiber. *Trends in Biotechnology* 18(9):374–379 DOI 10.1016/S0167-7799(00)01481-5.
- Holt, J., Cook A.J.,(1987); Simulation analysis of brown planthopper (*Nilaparvata lugens*) population dynamics on rice in the Philippines. *Jour. Applied Ecology* 24:87-102.
- Jeyaparvathi S, Baskaran S, Bakavathiappan G ,(2013); Biological control potential of spiders on the selected cotton pests. *International Journal of Pharmacy & Life Sciences*; 4(4):2568-2572.
- Karanjkar D.M. (2008) Biodiversity Of Arachnids In Satara Tehsil Minor Research Project Submitted To University Grants Commission Western Regional Office Ganeshkhind Pune.
- Karsch, E. (1873); Verzeichniss Westfälischer Spinnen (Araneiden) *Verh. Naturh. Ver. Preuss. Rhein. Westfal.* 10: 113-160.
- Keswani S,(2012); Checklist Of Spiders (Arachnida: Araneae) From India-2012, *Indian Journal Of Arachnology*, Vol.1 (1):1-129.
- King GF, Hardy MC. ,(2013); Spider-venom peptides: structure, pharmacology, and potential for control of insect pests. *Annual Review of Entomology* 58(1):475–496 DOI 10.1146/annurev-ento-120811-153650.
- Kobayashi, S., (1977) ;Changes in population density of spiders in paddy field during winter. *Acta Arachnol.* 27 (special number): 247-251.
- Kremen, C. ,(1993); Biological Inventory Using Target Taxa: Case Study Of The Butterflies Of Madagascar. *Ecological Applications*.
- Kulkarni S.S.,(2012); New Record Of Flower Spider *Thomisus sikkimensis* With A Note On Its Courtship Behaviour. *Bugs R All*.
- Kulkarni S, Lewis T (2015); Description of male *Tylorida sataraensis* Kulkarni, 2014 (Araneae, Tetragnathidae) with notes on habits and conservation status. *Biodiversity Data Journal* 3: e4451. doi: 10.3897/BDJ.3.e4451.
- Kulkarni S.S. (2016), A New Cavernicolous Assassin Bug Of The Genus *Bagauda bergrothi* (Heteroptera: Reduviidae: Emesinae) From The Western Ghats ,India; *Zootaxa* 4127(2): 365-375, ISSN 1175-5334.
- Kulkarni S, Yadav S (2015) Bridging the distributional gap of *Tylorida striata* (Thorell, 1877) and newsynonymy (Araneae: Tetragnathidae). *Biodiversity Data Journal* 3: e4878. doi: 10.3897/BDJ.3.e4878.
- Kulkarni S.S. , (2017); The Spiny Theridiid Genus *Meotipa* Simon, 1895 in India, With Description Of A Strange New Species With Translucent Abdomen And Phylogenetic Analysis About The Genus Placement (Araneae , Theridiidae); *Zootaxa* 4291(3): 504-520, ISSN 1175-5334.
- Kulkarni S.S. ,(2012); New Record Of The Spider *Arachnura angora* From Maharashtra With Additional Morphological Description. (Araneae: Araneidae).; *Rec. Zool. Surv. India* :112(Part-2)113-115.
- Kulkarni S.S.,(2015); First Record Of Genus *Siler* Simon, 1889 (Araneae: Salticidae) From India; *Journal Of Threatened Taxa* 7(10):7701-7703.
- Kulkarni S.S., (2012); A New Species Of The Genus *Oxyopes*, Latrielle (Araneae: Oxyopidae) From Sahyadri Ranges Of Western Ghats; (2012) Record Of Zoological Survey India-112(Part-2):35-37,(2012) ISSN 0375-1511.
- Kulkarni S.S., Deshpande V.Y. ,(2013); Preliminary Studies On The Spider Fauna Of Satara City ,Maharashtra, India (Araneae)., India ., *Proceedings Of National Seminar On Applied Research In Basic Sciences*.
- Kulkarni S.S.,(2011); Ecology And Behavioral Changes In Spider *Argiope anasuja* (Female); *Advances In Applied Science Research*, , 2 (3): 380-383 ISSN: 0976-8610 CODEN (USA): AASRFC
- Kulkarni SS, Lewis TR. (2015); Description of male *Tylorida sataraensis* Kulkarni, (Araneae, Tetragnathidae) with notes on habits and conservation status. *Biodivers Data J.* 2015;(3):e4451. Published 2015 Feb 16.

- doi:10.3897/BDJ.3.e4451
- Kulkarni, S.(2014). A new species of the genus *Tylorida* Simon, 1894 (Araneae: Tetragnathidae) from a rocky outcrop in the northern Western Ghats, India. *Journal of Threatened Taxa*, 6(3), 5558-5561. <https://doi.org/10.11609/JoTT.o3606.5558-61>
- Lee, H. P., Kim J.P.,JunJ.R. , (1993a.);Utilization of insect natural enemies andspiders for the biological control in ricepaddy field, community structure of insectpest and spiders, suppress effect on insectpest by natural enemies, and their overwintering habitats in rice paddy field. *RDA Jour. Agri. Sci. (Agri. Inst. Coop.)*35: 261-274.
- Li R, Yan Z, Wang J, (2017); De novo characterization of venom apparatus transcriptome of *Pardosapseudoannulata* and analysis of its gene expression in response to Bt protein. *BMC Biotechnol.* 2017;17(1):73. Published 2017 Nov 7. doi:10.1186/s12896-017-0392-z.
- Lüddecke T.(2019) Phylogeny-Guided Selection of Priority Groups forVenom Bioprospecting: Harvesting Toxin Sequences in Tarantulas as a Case Study;11, 488; doi:10.3390/toxins11090488 www.mdpi.com/journal/toxins.
- Maheshwari N.R., Patil S.R., (2018); Spider Fauna From North Maharashtra, India; *Flora And Fauna*. Vol.24No.ISSN 2456-9364(Online).
- Maloney, D., Drummond, F.A., Alford, R., (2003) ;Spider Predation In Agroecosystems: Can Spiders Effectively Control Pest Population?, MAFES 339 (Maine Agricultural And Forest Experiments Station) Technical Bulletin, 340 190, www.umaine.edu/mafes/elec_pubs/techbulletins/Tb190.pdf
- Mammola,(2017); Record breaking achievements by spiders and the scientists who study them. *PeerJ* 5:e3972; DOI 10.7717/peerj.3972.
- Maelfait, Jean-Pierre & Hendrickx, Frederik. (1998). Spiders as bio-indicators of anthropogenic stress in natural and semi-natural habitats in Flanders (Belgium): Some recent developments. *Proceedings of the 17th European Colloquium of Arachnology*.
- Mciver, J .D ., Moldenke, A .R ., Parsons, G .L . ,(1990); Litter Spiders As Bio-Indicators Of Recovery After Clearcutting In A Western Coniferous Forest . *Northwest Environmental Journal* 6 : 410-412 .
- Mirza Z, Sanap R., Vaze V., (2012), A New Species Of The Trapdoor Spiders Genus *Idiopis* Perty, 1833 (Araneae: Idiopidae) From The Western Ghats, With A Key To The Idiops Of India; *Revista Ibérica De Aracnología*, N° 21 (31/12/2012): 914. Grupo Ibérico De Aracnología (S.E.A.). ISSN: 1576 - 9518.
- Mirza ZA, Sanap RV, Bhosale H., (2014); Preliminary Review of Indian Eumenophorinae (Araneae: Theraphosidae) with Description of a New Genus and Five New Species from the Western Ghats. *PLoS ONE* 9(2): e87928. doi:10.1371/journal.pone.0087928
- More S., (2015); Diversity Of Spider Fauna From Bannoli Region Of Koyna Wildlife Sanctuary; *International Journal Of Science And Research (IJSR)* ISSN (Online): 2319-7064.
- More S.B., (2010); Spider Biodiversity In Western Ghats; Symposium Cum Workshop On Arachnology With Reference To Spiders, 28th Nov. To 1st Dec.2010.
- More S.B. , (2016), Diversity Of Spiders From Dandoba Hill Forest (Sangli); *IOSR Journal Of Pharmacy And Biological Sciences (IOSR-JPBS)* E-ISSN:2278-3008, P-ISSN:2319-7676. Volume 11, Issue 2 Ver. II PP 32-35, www.iosrjournals.org
- More S.B., (2015); Spider Diversity Of Rundiv, Sidheshwar And Ramnadi Area Of Chandoli National park, IJBAT, Issue (3), Vol. II, May 2015: 297-299 ISSN 2347 - 517X.
- More S.B., (2015); Inventorization Of Spider Diversity From Vakoba , Devrai Region Of Radhanagari Wildlife Sanctuary; *International Journal Of Science And Research (IJSR)* ISSN (Online):2319-7064, Index Copernicus Value (2013):6.14|Impact Factor (2013):4.438.
- Ndava J.,(2018);The future of mosquito control: The role of spiders as biological control agents: A review;ISSN: 2348-5906,CODEN: IJMRK2IJMR 2018; 5(1): 06-11,IJMR.
- Nyffeler, M., Benz G., (1987); Spiders In Natural Pest Control: A Review. *Journal Of Applied Entomology*, 103: 321-329.
- Peter J., Kulkarni S.(2016); An Unexpected New Species Of The Genus *Pseudopoda* (Araneae, Sparassidae, Heteropodinae) From The Western Ghats In India; *Zookeys* 577: 55-62, Doi: 10.3897/Zookeys.577.7848 .
- Pineda, S.S., Chaumeil P.A.,Kunert, A.(2018);

- ArachnoServer 3.0: An online resource for automated discovery, analysis and annotation of spider toxins. *Bioinformatics* 2018, 34, 1074–1076. [CrossRef] [PubMed]
- Pinkus MA, Ibarra-Nuñez G, Parra-Tabla V. (2006); Spider diversity in coffee plantations with different management in Southeast Mexico. *Journal of Arachnology*. 34(1):104–112.
- Platnick N. I., (2019); The World Spider Catalog, Version 20.5. American Museum Of Natural History, Online At [Http://Research.Amnh.Org/Iz/Spiders/Catalog](http://Research.Amnh.Org/Iz/Spiders/Catalog). DOI: 10.5531/Db.Iz.0001.
- Pocock R. I., (1900); The Fauna Of British India, Including Ceylon And Burma. Arachnida. London.
- Preston-Mafham K, Preston-Mafham R., (1995); Spiders Of The World. New York: Blandford Press Ltd. 1984. 24. Wise DH Spiders In Ecological Webs. Cambridge University Press, 1995: XIII + 328.
- Raiz T.N., (2018); Assessment Of Spider Diversity And Composition Along The Tungabhadra irrigation Channel At Ballari, Karnataka. *International Journal on Biological Sciences* 9 (1) : 36–44, ISSN No.: 0976–4518; NAAS Rating 3.14; UGC List of Journal No. 7775.
- Riechert S.E., Lockley T., (1984); Spiders As Biological Control Agents. *Annual Review Of Entomology* 29: 299–320.
- Russell-Smith A., (1999); The Spiders Of Mkomazi Game Reserve. In: Coe M, McWilliam N, Stone G, Parker M, Eds. *Mkomazi: The Ecology, Biodiversity And Conservation Of A Tanzanian Savanna*. Royal Geographical Society, London, P. 197–222.
- Saez NJ, Senff S, Jensen JE, (2010); Spider-venom peptides as therapeutics. *Toxins* 2(12):2851–2871 DOI 10.3390/toxins2122851.
- Samal, P., Misra B.C (1975) ; Spiders: The most effective natural enemies of the brown planthoppers in rice. *Rice Entomol. Newsletter* 3: 31.
- Sarwade A.B., Kamble N.A. (2017); Diversity And Seasonal Variation In Spider Abundance From Mirajtahsil, Sangli, Maharashtra; *Asian J. Biol. Life Sci. | Jan-Apr 2017 | Vol-6 | Issue-1*.
- Sasaba, T., Kiritani K., Kawahara S., (1973); Food preference of Lycosain paddy fields. *Bull. Kochi Inst. Agr. For. Sci.* 5: 61–63.
- Sebastian P. A, Peter KV, (2009) Spiders Of India. Universities Press (India) Private Limited.
- Shirbhate M. V. And Shirbhate A. M. (2017); Diversity And Distribution Of Spider Fauna (Family- Araneidae) In And Around Katepurna Sanctuary, Akola, India ; *Environment Conservation Journal* 18(3) 45–52, 2017 ISSN 0972–3099 (Print) 2278–5124 (Online) .
- Siliwal M, Molur S, Biswas B.K., (2005); Indian Spiders (Arachnida: Araneae): Updated Checklist. *Zoo's Print Journal* 20, 1999–2049.
- Simon, E., (1887i); Etude Sur Les Arachnides De l'Asie Méridionale Faisant Partie Des Collections De l'Indian Museum (Calcutta). I. Arachnides Recueillis À Tavoy (Tenasserim) Par Moti Ram. *Journal Of The Asiatic Society Of Bengal* 56: 101–117.
- Stoliczka F., (1869); Contribution Towards The Knowledge Of Indian Arachnoids. *Journal Of The Asiatic Society Of Bengal* 56, 101–117.
- Tanaka, K., (1989); Movement of the spiders in arable land. *Plant Protection* 43, 1:34–39.
- Thorell T., (1895); Descriptive Catalogue Of Spiders Of Burma, British Museum London, 1–406.
- Tikader B. K., Malhotra, M.S., (1980); The Fauna Of India. Spiders (Thomisidae And Lycosidae). *Zoological Survey Of India, Calcutta*: 44pp.
- Tikader, B. K., (1980); Fauna Of India - Araneae: Spiders, Vol. I (Araneidae & Gnaphosidae). *Zoological Survey Of India*. 448 Pp.
- Tikader, B. K., (1982); Fauna Of India - Araneae: Spiders, Vol. II (Thomisidae And Lycosidae). *Zoological Survey Of India*. 533 Pp.
- Turnbull, AL., (2003); Ecology Of The True Spiders (Araneomorphae). *Ann. Rev. Entomol.* 18. 305–348. 10.1146/Annurev.En.18.010173.001513.
- Van den B., R., Messenger P.S., Gutierrez A.P., (1982) A; n Introduction to Biological Control. Plenum Press, New York, USA. 247 pp.
- Wasankar P.P, Kakde V. R., (2016) ; Faunal Diversity Of Spider Found In Dnyanganga Sanctuary Of District Buldhana, Maharashtra, India ; *Online International Interdisciplinary Research Journal, {Bi-Monthly}*, ISSN 2249–9598, Volume-VI, May 2016 Special Issue.
- Whiteman, Honor. (2015) "Newly identified compounds in spider venom could help treat chronic pain." *Medical News Today. MediLexicon, Intl., Web*.
- Windley M J, Herzig V, Dziemborowicz SA, (2012); Spider-venom peptides as bioinsecticides. *Toxins*. 2012;4:191–227.