Dental Communication



Biosc.Biotech.Res.Comm. Vol 13 No (4) Oct-Nov-Dec 2020 Pp 1837-1841

Oral Health Knowledge and Attitude Among Students of King Abdulaziz University Jeddah Saudi Arabia

Heba Ashi¹, Farah Al Beiruty² Rahaf Bakheet² and Nawaf Almarzouki³ ¹Department of Dental Public Health, Faculty of Dentistry, King Abdulaziz University Jeddah, Saudi Arabia., ²Faculty of Dentistry, King Abdulaziz University, Jeddah, Saudi Arabia and

³Department of Ophthalmology, King Abdulaziz University Hospital, Jeddah, Saudi Arabia.

ABSTRACT

Maintenance of oral health and positive attitudes has an integral role in the success of dental treatment and has a direct effect on the prognosis. This study aimed to determine if dentists have more dedication to their oral health than non-dentists, as there is limited data surrounding this topic in Saudi Arabia. An electronic survey was conducted to collect the data from anonymous participants in both dental and non-dental fields. Main contents of survey questions were brushing, flossing, washing, fillings, crowns, extractions, orthodontics, halitosis, gum status, toothache and checkups. Chi square and Monto Carlo tests were used to analyze the results which detected significant differences on extractions, orthodontic appliance, halitosis and gum status for dental students with corresponding scores of (18.9%, 64.9%, 16.2%, 20.3% respectively) whereas non-dental students scored (33.6%, 45.8%, 29.8%, 40.5% respectively). With P-value of 0.025 for extractions, 0.009 for orthodontic appliance, 0.031 for halitosis, and 0.006 for gum status. The most prevalent problem was orthodontics. A significant difference was found between dental and non-dental students in terms of tooth brushing and flossing (75.7% VS 55.8% and 71.5% VS 23% respectively) (p = 0.046, P< 0.001 respectively). Missing tooth and halitosis were higher among males than females (50% VS 25.3% and 50% VS 22% respectively) with significant difference (p=0.011, 0.003 respectively). we highly recommend more oral hygiene health education programs especially for non-dental students. Further research is needed to assess oral health problems based on clinical examination and comprehensive detailed interviews to overcome internal validity errors that might occur in an electronic based survey.

KEY WORDS: DENTAL, ORAL, HEALTH, ATTITUDE, SAUDI ARABIA.

INTRODUCTION

Attitudes of dental students towards oral hygiene affect their habits and help improvements of oral health of their

Article Information:

Corresponding author email: hmashi@kau.edu.sa Received 24/10/2020 Accepted after revision 09/12/2020 P-ISSN: 0974-6455 E-ISSN: 2321-4007 Thomson Reuters ISI Clarivate Analytics Web of Science ESCI Indexed Journal

Identifiers & Pagination:

Vol 13(4) E-Pub 31st Dec 2020 Pp- 1837-1841 This is an open access article under Creative Commons License Attribution International (CC-BY 4.0) Published by Society for Science & Nature India DOI: http://dx.doi.org/10.21786/bbrc/13.4/30 society and patients (Sasanka et al., 2020). Good oral health depends on proper oral hygiene which helps in improving the quality of life (Halawany, 2012). It affects appearance, allows people to perform their daily activities without psychological or social limitations (Shah and El Haddad, 2015). Oral health relies on many factors such as personal attitudes, behaviors and knowledge. Dentists should be a role model for oral health attitudes and practices to their communities (Mekhemar et al., 2020). Dentists have a great role in this by providing prevention, treatment and helping people change their behavior and attitudes towards oral hygiene (Moheet and Farooq, 2013). However, poor oral health awareness was reported among Saudi dental students (Baseer and



Rahman, 2014). In addition, gingival diseases were found to be highly prevalent among dental students (Ahmad et al., 2019). There is an observed need to increase oral health awareness, positive attitudes and correct practices of dental students (Ahmad et al., 2019). Therefore, teaching students the skills needed for good oral health is as important as giving them knowledge (Halawany et al., 2015). In literature, the focus was mainly on treatment of the oral diseases rather than prevention, (Levin and Shenkman, 2004, Glick et al., 2012, Al-Nasser and Lamster, 2020).

The aims of this study are to increase awareness of oral hygiene importance to both dental and non-dental students, determine the most prevalent oral health problems among the groups in order to formulate a targeted community service. This in addition to assessing oral hygiene, attitudes and compliance among dental students and non-dental students and among female dental students or male ones in King Abdul-Aziz University Jeddah Saudi Arabia.

MATERIAL AND METHODS

This comparative cross-sectional study included 206 students at King Abdul-Aziz University Jeddah KSA. Electronic database search was done using Google Scholar and PubMed to gather background information and data related to the research question and determine the gap of knowledge. Then, formulating an anonymous electronic survey with multiple choice questions for dental and non-dental students to answer during the period from 29/01/2020 to 31/08/2020, followed by data analysis and segregation of the results. Arabic language was chosen for this survey to avoid any possibilities of language barriers, followed by data analysis and segregation from the survey. Students were stratified according to specialty and gender. Data were analyzed using the statistical package for the social sciences (SPSS), version 25. Categorical variables were presented as numbers and percentage. Results were compared using Chi-square test, Monto Carlo test and Fisher Exact test. All tests were 2-tailed, and a p-value of <0.05 was considered statistically significant.

Table 1. Oral	hygiene and oral health a	ccording	; to dental sp	ecialty	(n = 206)	
		Dental students (n=74)		Non dental students (n=131)		P-value
		N	0/0	N	0/0	
Brushing	Not daily	3	4.1%	9	7.0%	0.046*
	Once	9	12.2%	28	21.7%	
	Twice	56	75.7%	72	55.8%	
	More than twice	6	8.1%	20	15.5%	
Floss	I don't	17	23.0%	93	71.5%	<0.001*
	Once	51	68.9%	29	22.3%	
	Twice	4	5.4%	6	4.6%	
	More than twice	2	2.7%	2	1.5%	
Missing	I didn't	60	81.1%	87	66.4%	0.058**
tooth	1-2	12	16.2%	28	21.4%	
	3-4	1	1.4%	13	9.9%	
	5-6	0	0.0%	2	1.5%	
	>6	1	1.4%	1	0.8%	
Orthodontics	Yes, currently	6	8.1%	11	8.4%	0.022*
	Yes, in the past	42	56.8%	49	37.4%	
	Never	26	35.1%	71	54.2%	
Halitosis	Yes	11	14.9%	39	29.8%	0.016**
	No	62	83.8%	92	70.2%	
Gum	Excellent	57	77.0%	72	55.0%	0.006**
	Bleeding during brushing	15	20.3%	53	40.5%	
	Bleeding mostly	0	0.0%	3	2.3%	
Note. All var	iables are summarized as r	number a	nd percentag	e	1	
The test of si	gnificance was carried out	at 0.05	level			
*Chi- Square	test was used					
**Monto Carl	o test was used					
Significant re	esults are in bold					

Ethical considerations: the study was approved by Research Ethics Committee of the Faculty of Dentistry (REC-FD) Consent was governed automatically by participating in the survey, as it was declared before taking it.

RESULTS AND DISCUSSION

Out of 206 participants, 88.3% were females, (86.4%) with average age of 20-30 years old. Descriptive data are shown in table 1 with 73.3% postgraduate and 63.6% were non dental students. In regard of oral hygiene measures, a significant difference was found between dental and non-dental students in terms of tooth brushing and flossing (p = 0.046, P < 0,001 respectively). However, no significant difference was found between dental and non-dental students comparing the oral and tooth conditions. In regards of dental visits and compliance, the dental students showed higher compliance than the non-dental students (Table 1).

The most common oral health problem among both dental (64.9%) and non- dental students (45.8%) was malocclusion followed by bleeding on brushing (p= 0.009 and p=0.006 respectively) (Table 2, fig.1).

Table 2. Prevalence of the most common Oral health

problems in dental students and non-dental students

(n-205)

(1-203)							
	Dental students (n=74)	Non dental students (n=131)	P-value				
Fillings	82.4%	75.6%	0.254*				
Crowns	13.5%	9.9%	0.434*				
Missing tooth	18.9%	33.6%	0.025*				
Orthodontics	64.9%	45.8%	0.009*				
Halitosis	16.2%	29.8%	0.031*				
Bleeding during	20.3%	40.5%	0.006**				
brushing							
Bleeding mostly	0.0%	2.3%					
Receding gum	2.7%	2.3%					
Toothache	17.6%	26.7%	0.137*				

Note. All variables are summarized as percentage The test of significance was carried out at 0.05 level *Chi- Square test was used **Monto Carlo test was used Significant results are in bold

When comparing between both genders, missing tooth and halitosis were higher among males than females (50% VS 25.3% and 50% VS 22%) with significant difference (p=0.011, 0.003) (Table 3, fig 2).

This comparative cross-sectional study was conducted at King Abdul-Aziz University including 206 dental specialty and non-dental specialty students. This study assesses the knowledge and attitudes towards oral Health among Dental Students and Non-dental Students. To summarize our findings, dental specialty students had better attitudes towards oral hygiene and health, having less oral health problems than non-dental specialty students. Female students also had better oral attitudes with less prevalence of oral problems than males. As regard to oral health and attitudes among dental specialty students, our findings supported the conclusions of Gufran and colleagues. Most of our dental students reported brushing their teeth twice daily. In 2015, Gufran and fellows reported that 82% of their subject students brushed their teeth twice daily (Gufran et al., 2015).

Most of our students who reported not brushing their teeth on a daily basis were of a non-dental specialty. In addition, flossing was higher among dental students suggesting better oral health and attitudes among them. This corresponds to the findings of Santhosh Kumar et al., who found that dental students had a better attitude towards oral health and a better behavior when compared to pharmacy students (S. Kumar et al., 2012). Also another study showed that dental students had better knowledge, attitude and practice than medical students (H. Kumar et al., 2017).

In addition, a study by Baseer found a significant difference between dental hygiene behaviors among government and private universities students which was supported with a difference between the clinical and preclinical dental education (Baseer et al., 2013). In fact, knowledge among dental students is not satisfactory and they need more awareness about oral self-hygiene practice which is the same as what a study by Lavanya et al. concluded (Lavanya & Nallamilli, 2014).Regarding the prevalence of oral health problems in dental students and non-dental students, missing tooth, halitosis, bleeding on brushing and spontaneous bleeding were more frequently reported among non-dental students. This collaborates what Ahmad and colleagues reported in 2019; a study that documented a high prevalence of gingival diseases among all study subjects (Ahmad et al., 2019).

On the other hand, our study reported an unexpected higher prevalence of orthodontics among students of dental specialty. Similarly a study by Wagle et al. concluded that dentists had better oral health than general population and that the prevalence of decayed and unfilled teeth was lower among dentists, (Wagle et al., 2014).When it comes to gender-based oral health comparison among our sample, the prevalence rates of not brushing teeth daily, not using floss and halitosis were higher among males than females. Furthermore, the prevalence rates of missing tooth and halitosis were higher among male subjects than females. Therefore, we can conclude that females are more committed to oral health and hygiene. This could be a result of their concern about how their mouth and teeth look. This was similar to the findings of a study by Kumar et al. where they found that females had better knowledge and oral health practices than males (Kumar et al., 2017).

Note. All variables are summarized as percentage

		Males (N=24)		Females (N=182)		P-valu
	-	N	0/0	N	0/0	1
Brushing	Not daily	5	21.7%	7	3.9%	0.006*
	Once	5	21.7%	32	17.7%	
	Twice	11	47.8%	118	65.2%	
	More than twice	2	8.7%	24	13.3%	1
Floss	I don't	19	79.2%	92	50.8%	0.072*
	Once	5	20.8%	75	41.4%	
	Twice	0	0.0%	10	5.5%	1
	More than twice	0	0.0%	4	2.2%	1
Missing tooth	I didn't	12	50.0%	136	74.7%	0.064*
-	1-2	8	33.3%	32	17.6%	
	3-4	3	12.5%	11	6.0%	
	5-6	0	0.0%	2	1.1%	
	>6	1	4.2%	1	0.5%	
Orthodontics	Yes, currently	3	12.5%	14	7.7%	0.267**
	Yes, in the past	7	29.2%	84	46.2%	
	Never	14	58.3%	84	46.2%	1
Halitosis	Yes	12	50.0%	39	21.4%	0.030*
	No	12	50.0%	142	78.0%	
Gum	Excellent	11	45.8%	119	65.4%	0.205*
	Bleeding during brushing	11	45.8%	57	31.3%	
	Bleeding mostly	1	4.2%	2	1.1%	1
	No	21	87.5%	136	74.7%	1
Checkup	Yes, every 6 months	3	12.5%	42	23.2%	0.689*
	Yes, once per year	5	20.8%	28	15.5%	
	No, only during pain	13	54.2%	91	50.3%	1
	Never	3	12.5%	20	11.0%	
Note. All varia The test of sig	bles are summarized as num nificance was carried out at test was used	iber an 0.05 lev	d percentage vel			

Significant results are in bold



Similarly another study by Jaber et al. found that male students had good knowledge but poor practice toward oral health, (Jaber et al., 2017). Our paper showed the difference in oral health and attitudes, prevalence of oral problems among dental and non-dental students, female and male dental students. Thus, we highly recommend more oral hygiene health education programs especially for non-dental students.

We obtained our data using an online survey which has some limitations such as possibility of inaccurate information as the survey is in the form of multiplechoice questions.

The survey could be answered by the same person twice as it was anonymous which could affect the integrity of the results. Limited number of questions were used as increasing the questions might bore the participant which could affect the accuracy of the results. Non-dental students might provide us with non-accurate answers because they are not keen in the dental field. Dental and non-dental students could be biased either to their side or against which could affect the integrity of the results. Respondents may not feel comfortable providing answers that present themselves in an unfavorable manner.

CONCLUSION

Dental specialty students had better attitude towards oral hygiene, oral health and having less oral health problems than non-dental specialty students, so we highly recommend more oral hygiene health education programs especially for non-dental students. Female students were showing more positive attitude than males with the prevalence of oral health problems were more among male students. During literature search, we found limited data on oral health and attitudes of dental students in Saudi Arabia. About eleven studies conducted in Saudi Arabia were found, none of them was in Jeddah or King Abdulaziz University specifically. Further research is needed to assess oral health problems based on clinical examination and comprehensive detailed interviews to overcome internal validity errors that might occur in an electronic based survey.

REFERENCES

Ahmad, F. A., Alotaibi, M. K., Baseer, M. A., et al. (2019). The Effect of Oral Health Knowledge, Attitude, and Practice on Periodontal Status among Dental Students. European Journal of Dentistry, 13(3), 437–443. https:// doi.org/10.1055/s-0039-1697109

Al-Nasser, L., & Lamster, I. B. (2020). Prevention and management of periodontal diseases and dental caries in the older adults. Periodontology 2000, 84(1), 69–83. https://doi.org/10.1111/prd.12338.

Baseer, M. A., Rahman, G., Kawaey, Z. Al, et al. (2013). Evaluation of Oral Health Behavior of Female Dental Hygiene Students and Interns of Saudi Arabia by Using Hiroshima University Dental Behavioural Inventory (HU-DBI). Oral Health and Dental Management, 12(4), 255–261. https://doi.org/10.4172/2247-2452.1000522 Baseer MA, Et Rahman, G. (2014). Oral health attitudes and behavior among a group of female Saudi dental students. Saudi Journal of Oral Sciences, 1(1), 25. https://doi.org/10.4103/WKMP-0056.124182

Glick, M., Monteiro Da Silva, O., Seeberger, G. K.,et al. (2012). idj_12009 278..291. https://doi.org/10.1111/ idj.12009.

Gufran, K., Indrajit, M. G., Aljeaidi, Z., et al. (2015). Self-care of Oral Health Status and Uses of Interdental Aids among Dental Students: A&tnbsp;Cross-Sectional Study. International Journal of Dental and Medical Specialty, 2(4), 5. https://doi.org/10.5958/2394-4196.2015.00024.2

Halawany, H. S. (2012). A review on miswak (Salvadora persica) and its effect on various aspects of oral health. In Saudi Dental Journal (Vol. 24, Issue 2, pp. 63–69). Elsevier. https://doi.org/10.1016/j.sdentj.2011.12.004. Halawany, H. S., Abraham, N. B., Jacob, V., et al. (2015). The perceived concepts of oral health attitudes and behaviors of dental students from four Asian countries. Saudi Journal for Dental Research, 6(2), 79–85. https://doi.org/10.1016/j.sjdr.2014.09.002.

Jaber, M. F., Khan, A., Elmosaad, Y., et al. (2017). Oral health knowledge, attitude and practices among male Qassim university students. International Journal Of Community Medicine And Public Health, 4(8), 2729. https://doi.org/10.18203/2394-6040.ijcmph20173316 Kumar, H., Behura, S. S., Ramachandra, S., et al. (2017). Oral health knowledge, attitude, and practices among dental and medical students in Eastern India -A comparative study. Journal of International Society of Preventive and Community Dentistry, 7(1), 58–63. https://doi.org/10.4103/jispcd.JISPCD_30_17

Kumar, S., Tadakamadla, J., Kumar, S., et al. (2012). Inviting submissions: Special issue-Periodontal and Peri-Implant Diseases View project Kingdom of Saudi Arabia. In Orofacial Sciences The Journal of the School of Dental Sciences Universiti Sains Malaysia Arch Orofac Sci (Vol. 7, Issue 1). https://www.researchgate. net/publication/266010587.

Lavanya, R., & Nallamilli, S. (2014). Oral hygiene practices and habits among dental students and staff in a dental college, India. https://www.researchgate.net/publication/261646570_Oral_hygiene_practices_and_habits_among_dental_students_and_staff_in_a_dental_college_India.

Levin, L., & Shenkman, A. (2004). The Relationship Between Dental Caries Status and Oral Health Attitudes and Behavior in Young Israeli Adults. Journal of Dental Education, 68(11), 1185–1191. https://doi.org/10.1002/ j.0022-0337.2004.68.11.tb03864.x.

Mekhemar, M., Conrad, J., Attia, S., et al. (2020). Oral Health Attitudes among Preclinical and Clinical Dental Students in Germany. International Journal of Environmental Research and Public Health, 17(12), 4253. https://doi.org/10.3390/ijerph17124253

Moheet A, I., & Farooq, I. (2013). Self-reported differences between oral health attitudes of pre-clinical and clinical students at a dental teaching institute in Saudi Arabia. Saudi Dental Journal, 25(4), 149–152. https://doi.org/10.1016/j.sdentj.2013.07.001

Sasanka, K., Jayaraj, G., & Ganapathy, D. (2020). Awareness of Oral Hygiene among Dental Students-A Survey. 14(4), 5494–5502.

Shah, A., & ElHaddad, S. (2015). Oral hygiene behavior, smoking, and perceived oral health problems among university students. Journal of International Society of Preventive and Community Dentistry, 5(4), 327. https:// doi.org/10.4103/2231-0762.161765.

Wagle, M., Trovik, T. A., Basnet, P., et al (2014). Do dentists have better oral health compared to general population: A study on oral health status and oral health behavior in Kathmandu, Nepal. BMC Oral Health, 14(1), 23. https://doi.org/10.1186/1472-6831-14-23.