

Biosc.Biotech.Res.Comm. Vol 13 No (4) Oct-Nov-Dec 2020 Pp 1947-1953

Oxidative Degradation of Biodiesel: Modelling of Different Paths and Effect of Antioxidants on two Mechanisms

Ishty Malhotra^{1*} and Mansi Srivastava²

¹Department of Chemical Engineering and Technology, Indian Institute of Technology, Varanasi, India ²Department of Chemical Engineering, Indian Institute of Technology, Delhi, India

ABSTRACT

Since past many years, biodiesel has been considered as a locally delivered, clean-consuming, inexhaustible substitute for oil diesel. Utilizing biodiesel as a vehicle fuel expands energy security since it contains large amount of methyl esters in the form of especially poly-unsaturated methyl esters. Due to presence of low oxidative stability, they can get easily oxidised and their long storage results into degradation. In order to control this oxidation process, antioxidants are used. Various studies have been performed concentrating upon the oxidative stability but very few focussed upon the mechanisms involved in oxidative degradation and the effect of antioxidants on reactions. The present study models the two different pathways of oxidative degradation of biodiesel for 5 years using the MATLAB 2020 and study of consumption of antioxidants to control the reaction mechanism has taken place. Comparison has been done between the oxidation of biodiesel in the absence and presence of antioxidant via the explanation of two different mechanisms. Change in concentration and consumption of RH (biodiesel organic matter) and antioxidant has been studied during the two different paths of oxidation. It has been concluded that antioxidant has affected the path one process more as compared to the path two process during the degradation mechanism of biodiesel in 5 years. Whereas consumption of antioxidant in path one is only approximate 3% as compared to 99% in path two. Therefore, oxidative degradation carried from path one can be controlled with lower amount of antioxidant whereas no such effect of antioxidant takes place during reactions of path two even after its complete consumption.

KEY WORDS: ANTIOXIDANTS, BIODIESEL, OXIDATION, METHYL-ESTERS, DEGRADATION.

INTRODUCTION

Since, many years, research has been performed for variety of energy resources which is nearly as old as the man's need of energy. Energy has now become the

Article Information:

Corresponding author email: *anjalim4897@gmail.com* Received 30/10/2020 Accepted after revision 12/12/2020 P-ISSN: 0974-6455 E-ISSN: 2321-4007 Thomson Reuters ISI Clarivate Analytics Web of Science ESCI Indexed Journal

Identifiers & Pagination:

Vol 13(4) E-Pub 31st Dec 2020 Pp- 1947-1953 This is an open access article under Creative Commons License Attribution International (CC-BY 4.0) Published by Society for Science & Nature India DOI: http://dx.doi.org/10.21786/bbrc/13.4/47 most important part to be considered for global economic growth. To meet the energy requirements, the use of fossil fuels is highly discouraged due to its contribution in greenhouse emission gas to environment, increasing instability in supply and rising production cost (Bannister et al., 2011). Similarly, now-a-days, petroleum reserves are diminishing at a high rate and petroleum fuelled engines are affecting the environment in negative manner, due to which, importance of alternative fuels is increasing. Therefore, urgent need has raised to replace fossil fuels to fulfil the demand of the world. Biofuels said to have more focal points over petroleum derivatives and other sustainable wellspring of energy since it lessens reliance on unfamiliar oil, it is delivered locally subsequently



Malhotra & Srivastava

making work and elevating the economy, and has zero or unimportant outflows (Muhammad et al., 2018; Kåberger, 2018).

Hereby, biodiesel has become as an attractive and essential energy resource. Biodiesel contains large percentage of unsaturated methyl esters, which are especially in the form of poly-unsaturated methyl esters. They have comparatively lower oxidation stability than petroleum diesel. When the biodiesel is stored over time, their oxidative stability becomes a matter of concern. The storage problems that are merely caused by storage conditions are exposure to air, light, temperature as well as presence of metals with catalytic effect on oxidation (Sharma et-al, 2009; Karavalakis et al., 2010). To check the biodiesel quality control, oxidation stability is used as a parameter. The oxidation degradation process of biodiesel depends upon the factors such as fatty acids used in production, humidity, temperature and light. The biodiesel deterioration during operation has become the concern for many industries in recent years (Maia et al., 2011; Delgado et al., 2019).

Products formed during oxidation, not only affects the life in storage but also cause fuel filters clogging. To improve the oxidation stability, antioxidants are used by removing the free radical formed at initiation stage. The use of antioxidants helps in delaying of onset of oxidation with free radicals to form stable compounds which do not contribute to oxidation chain reactions. Natural antioxidants like tocopherols, sterols etc are found in vegetable oils, which are strongly affected by production and refining of biodiesel and parent oils respectively. Antioxidants stabilized the fatty peroxy radicals and hereby, the chain reaction gets stopped. Antioxidant do not inhibit but retards the polymerisation. Indicators like formation of colour, presence of soluble gums and insoluble in the fuel helps in stability. (Razon et al., 2016; García et al., 2017; Valenga et al., 2019).

Various work has been done on biodiesel oxidative mechanism which explains that oxidation of biodiesel involves two different pathways but very few focussed upon the effect of antioxidants on the oxidation mechanism. The modelling of reactions to determine the consumption of antioxidant and the organic compound hasn't been taken into popular account. (Rashed et al., 2015; Sazzad et al., 2016; Varatharajan et al., 2017). Hereby, this paper discusses the two different pathways through which oxidation of biodiesel can take place and their modelling for 5 years in MATLAB 2020 in order to understand the amount of degradation that takes place during the oxidation in presence and absence of antioxidant in two different pathways. This will also help to study the oxidation path which gets affected by the presence of antioxidants (Varatharajan et al., 2017).

MATERIAL AND METHODS

The oxidation mechanism of different hydrocarbons has been a subject to various research papers. Any mechanism while processing mainly involves chemical free radical process. Any hydrocarbon contains sites which respond with oxygen by means of the autoxidation component with the old-style revolutionary chain response steps of initiation chain propagation and termination. The initiation step is the development of a free radical that can respond straightforwardly with oxygen to structure a peroxide or hydroperoxide. General mechanism on the basis of which oxidation of unsaturated fatty acid takes place is initiation step. Initially, formation of peroxides takes place, which get additionally corrupted utilizing pathways and the auxiliary items framed utilizing them like acids, aldehydes and polymers. (Saeed et al., 2012).

In the Initiation Period, initiators, for example, heat, light, high energy radiation, metal particle impetuses eliminate a hydrogen ion from the unsaturated fat structure and produce free radicals as appeared in R1. RH speaks to a natural substrate being oxidized (FAME), R* is a carbon-focused alkyl radical got from RH, and HOO* is hydro-peroxyl radical. The carbon-focused radical (R*) shaped in the commencement response are exceptionally receptive, and joins with accessible oxygen, giving peroxy radicals (ROO*) in R2 depicted under Propagation Reaction (Herbinet et al., 2008; Riazi et al., 2017; Riazi et al., 2019; Srinivasan et al., 2019).

The peroxy free radical isn't as receptive as the carbon free radical, yet regardless are adequately responsive and rapidly eliminates another hydrogen by assaulting the frail piece of the FAME structure to shape a hydroperoxide in R3 for example (ROOH*) radical and a R*. The new R* from the above condition would then be able to respond with diatomic oxygen to make the peroxy radicals in R2, and the chain response proliferates. Response of at least two free radicals' prompts bring about end part. This is called Termination Reaction. The essential or chain breaking antioxidants agents responds with this peroxy (ROO*) and hydroperoxide (ROOH*) extremists and intrudes on the proliferation response by arrangement of antioxidant radical for example A*. Essential antioxidants don't restrain the initiation response of the biodiesel degradation (McCormick et al., 2007; Noirbent et al., 2020).

This A* framed, therefore follows two ways, first way prompts the response with antioxidant radical with the diatomic oxygen prompting arrangement of AO2*. This AO2* responds and structure various items. It has been expected by various writings that two pathways would happen alongside rudimentary thought of response. Second way prompts the response between the antioxidant radical with peroxide radical prompting development of ROOA. The estimation of rate consistent has been accepted hypothetically however considering the response component of oxidation of RH (Maia et al., 2011; Focke et al., 2011; Varatharajan et al., 2017).

In this study, two different above discussed pathways has been related to the impact of antioxidant. Modelling of the listed reactions has been done in order to evaluate the effect of antioxidant in the reaction on the whole process. Using MATLAB 2020, study has been performed in considering the cases in which antioxidant concentration during the reactions has been converted from 0.09 mol/m3 to the final concentration. The concentration of RH (organic substrate) has been taken as 0.03 mol/m3. Hereby, during the modelling, the requirement of rate

constants has been raised which have been fulfilled by the previously available studies (Waynick et al., 2005, Zuleta et al., 2012). Table 1 describes the reactions of two pathways along with the assumed kinetic rate constant using different literatures.

Table 1. Reaction Mechanism						
Path Number	Reaction Number	Reaction Scheme	Rate Constant Value			
1,2	R1	$\rm RH + O_2 \rightarrow \rm R^* + \rm HOO^*$	1*10 ⁻⁴ (mol/cm3) s ⁻¹			
1,2	R2	$R^* + O_2 \rightarrow ROO^*$	1*10 ¹⁰ (mol/cm3) s ⁻¹			
1,2	R3	$ROO^* + RH \rightarrow ROOH + R^*$	1*10 ⁻⁴ (mol/cm3) s ⁻¹			
1,2	R4	Antioxidant + $RO0^* \rightarrow A^* + ROOH$	1*10 ⁴ (mol/cm3) s ⁻¹			
1	R5	$R^* + R^* \rightarrow R-R$	1*10 ¹¹ (mol/cm3) s ⁻¹			
1	R6	$A^* + 0_2 \rightarrow A02^*$	1*10 ⁻² (mol/cm3) s ⁻¹			
1	R7	$AO_2^* + AO_2^* \to P$	1*10 ⁻² (mol/cm3) s ⁻¹			
2	R8	$ROOH + A^* \rightarrow ROOA$	1*10 ⁻⁸ (mol/cm3) s ⁻¹			



RESULTS AND DISCUSSION

The modelling part has been done using MATLAB 2020, in which the amount of degradation of organic substrate and antioxidant has been studied that takes place for 5 years, if an oxidation mechanism would happen for both the paths. Determination of the exhausted amount of antioxidant would also take place during the two different pathways in the modelling. As explained above, the RH oxidation decreases in presence of antioxidant. Similar, antioxidant study of safflower biodiesel has also been proved by various literatures which shows the relation of stability of biodiesel with induction period in regards with antioxidants. Other than this, the impact of

BIOSCIENCE BIOTECHNOLOGY RESEARCH COMMUNICATIONS



various antioxidants on refined biodiesel is like the first biodiesel: PY, PG, and TBHQ (tert-butylhydroquinone) gave the best outcome, trailed by BHA (butylated hydroxyanisole), BHT (butylated hydroxytoluene), DTBHQ, IB, and a-tocopherol showed that antioxidants don't inhibit the mechanism but slow down the process. (Tang et al., 2008; Xin et al., 2008; Karavalakis et al., 2010; Focke et al., 2011; Varatharajan et al., 2017; Devi et-al, 2018).

Similar modelling of reactions of biodiesel oxidation has been done that shows the consumption of biodiesel in the presence and absence of antioxidants in two different pathways for 5 years. Table 2 data obtained from the

Malhotra & Srivastava

modelling in MATLAB 2020 describes the percentage in which the exhaustion of RH takes place in two different cases. It has been seen that, degradation of RH becomes slower in Path 1 in the presence of antioxidant, whereas, not much diference has been seen in Path 2 for presence and absence of antioxidant (Devi et-al, 2018).

Table 2. Degradation of RH for Different Cases w.r.t Time.						
Time,	Percentage Ex	Percentage Exhausted in Path 1		Percentage Exhausted in Path 2		
Days	Presence of	Absence of	Presence of	Absence of		
	antioxidant	antioxidant	antioxidant	antioxidant		
200	93.3667	97.0733	98.9933	99.0500		
400	95.5700	99.5866	99.9005	99.9088		
600	98.0600	99.9483	99.9903	99.9913		
800	98.0500	99.9936	99.9990	99.9991		
1000	98.6966	99.9992	99.9999	99.9999		
1200	99.1366	99.9999	100	100		
1400	99.4266	100	100	100		
1600	99.6166	100	100	100		
1800	99.7460	100	100	100		





As shown in the plots obtained from modelling in MATLAB, in the Path 1 case, complete degradation of RH takes place in more than 1800 days in presence of antioxidant (Figure 3) and in approximately 1400 days in absence of antioxidant (Figure 4). It clearly shows

Figure 5: Degradation of RH in the second path in presence of antioxidant





that antioxidant lowers the rate of degradation of RH. Whereas for Path 2, complete degradation of RH takes place in 1200 days only in exponental plot, although the presence and absence of antioxidant doesn't effects much during the degradation mechanism (Figure 5, Figure 6).

1950 OXIDATIVE DEGRADATION OF BIODIESEL

Such study has also been done for soybean oil fatty acid methyl esters (SME) using five different antioxidants. It has been proved that expanding antioxidants stacking (concentration) indicated sharp expansions in action for loadings up to 1000 ppm followed by more modest expansions in action at higher loadings (Varatharajan et al., 2017).

Table 3. Degradation of antioxidant for Two Different

Paths w.r.t Time.

Time, Days	Percentage Exhausted		
	Antioxidant	Antioxidant	
	in Path 1	in Path 2	
200	1.0166	38.9666	
400	1.8444	66.2688	
600	2.3244	81.4777	
800	2.9888	89.9692	
1000	2.8577	94.4855	
1200	2.9988	97.0800	
1400	3.0933	98.4277	
1600	3.1544	99.1544	
1800	3.1955	99.5544	

Figure 7: Degradation of antioxidant in the first path Degradation of Antioxidant During the First Pathway mol/cm3 0.0905 0.09 0.0895 0.089 ation of Antio 0.0885 0.088 0.0875 0.087 0.0865 500 1000 1500 2000 Time, days

It is clearly observed in table 2 that, antioxidant gets used more in case of second path in order to carry out oxidative stability i.e., it consumed to 99.55% of its initial value. Whereas, only 3.19% of antioxidant gets consumed to carry out the same amount of oxidative stability of RH in case of path one. Hereby, it can be said that if path one is followed, it may decrease the cost of antioxidant to maintain the stability of same amount of RH in exponentially manner (Figure 7, Figure 8). The mechanism followed during the oxidation path of biodiesel becomes dependent of various factors. Since, addition of antioxidants causes storage stability and about association between the physical and compound attributes of biodiesel with natural variables, for example, light, metal pollution, shading changes, and dregs development. The difference between the consumption of antioxidant in different paths arises, due to formation of products like aldehyde, ketones etc, if secondary

Other than this, various other researches involve execution of antioxidants assessed for treating biodiesel got from soybean oil, rapeseed (low-erucic) oil, utilized cooking oil, fat (meat), and palm oil feedstocks proved that different antioxidants provide different change in concentration of same origin of biodiesel. However, no such work came across which focusses upon the routes of oxidation and effect of antioxidants on different routes. However, the present study becomes relevant as comparable to previous researches since the common antioxidant and common biodiesel concentration change takes place during the modelling (Dunn, 2005; Dunn, 2008; Ashok et al., 2017; Varatharajan et al., 2017). During the modelling part, in the presence of antioxidant, since the antioxidant gets reacted during the reaction and product formation takes place that shows the consumption of antioxidants as well. Table 3, Figure: 7 and Figure 8, data obtained from simulation part describes the percentage in which the exhaustion of antioxidant takes place in the two different cases (Varatharajan et al., 2017).



mechanism is followed (Kivevele et al., 2011; Rashed et al., 2015).

CONCLUSION

Degradation of aromatic compounds is the common phenomenon observed in industries. Such degradation also takes place for hydrocarbon compounds like biodiesel. The objective of the paper is to study about the consumption of antioxidant at different paths of biodiesel oxidation along with the degradation of biodiesel (RH) during the two paths. Both the studies have taken place keeping the duration of 5 years. It has been concluded that antioxidant has affected the path one process more as compared to the path two process during the degradation mechanism of biodiesel in 5 years. Whereas consumption of antioxidant in path 1 is only approximate 3% as compared to 99% in path two. Therefore,

oxidative degradation carried from path one can be controlled with lower amount of antioxidant whereas no such effect of antioxidant takes place during reactions of

Malhotra & Srivastava

path two even after its complete consumption. This data, can become useful for the industries which stores any organic substrate (like biodiesel), since the storage of such compounds can become harmful specially when chance of degradation like oxidation, corrosion etc are common for few compounds. To obtain concentration vs. time plot for biodiesel and antioxidants, the concentrations may be measured by techniques like Spectrophotometry or HPLC. Such type of mathematical models together with enthalpy balance, can be developed to predict the potential hazardous oxidation leaks in storage tanks. Such studies would help in developing safer practices.

REFERENCES

Ashok B., Nanthagopal K., Jeevanantham A.K., Bhowmick P., Malhotra D. and Pranjal Agarwal. (2017). An assessment of *Calophyllum inophyllum* biodiesel fuelled diesel engine characteristics using novel antioxidant additives. Energy Conversion and Management. 148. 935-943.

Bannister C.D, Chuck C.J, Bounds M., and Hawley J.G, (2011). Oxidative Stability of Biodiesel Fuel. Proceedings of the Institution of Mechanical Engineers, Part D: Journal of Automobile Engineering. 225 (D1). 99-114. Delgado S.N., Encinar J.S. and González J.F. (2019). Saffower Biodiesel: Improvement of its Oxidative Stability by Using BHA and TBHQ. Energies. 12. 1-40 Devi A., Das V.K. and Deka D. (2018). Evaluation of the effectiveness of potato peel extract as a natural antioxidant on biodiesel oxidation stability. Industrial Crops and Products. 123. 454-460,

Dunn R.O. (2005). Effect of antioxidants on the oxidative stability of methyl soyate (biodiesel). Fuel Processing Technology. 86. 1071– 1085

Dunn R.O. (2008). Antioxidants for improving storage stability of biodiesel. Bio-fuel bio-product and biorefinery. 2. 304–318

Focke W.W., Westhuizen I.V.D., Grobler A.B.L., Nshoane K.T., Reddy J.K. and Luyt A.S. (2012). The effect of synthetic antioxidants on the oxidative stability of biodiesel. Fuel. 94. 227-223

García M., Botella L., Gil-Lalaguna N., Arauzo J., Gonzalo A. and Sánchez J.L. (2017). Antioxidants for biodiesel: Additives prepared from extracted fractions of bio-oil. Fuel Processing Technology. 156. 407-414

Herbinet O., Pitz W.J. and Westbrook C.K. (2008). Detailed chemical kinetic oxidation mechanism for a biodiesel surrogate. Combustion and Flame. 154. 507–528.

Kåberger T. (2018). Progress of renewable electricity replacing fossil fuels. Global Energy Interconnection. 1(1). 48-52

Karavalakis G., Hilari D., Givalou L., Karonis D. and Stournas S. (2010). Storage stability and ageing effect of biodiesel blends treated with different antioxidants.

Energy. 36. 369-374.

Kivevele T.T., Mbarawa M.M., Bereczky A., Laza T. and Madarasz J. (2011). Impact of antioxidant additives on the oxidation stability of biodiesel produced from Croton Megalocarpus oil. Fuel Processing Technology. 92. 1244–1248

Maia E.C.R., Borsato D., Moreira I., Spacino K.R. Rodrigues P.R.P. and Gallina A.L., (2011). Study of the biodiesel B100 oxidative stability in mixture with antioxidants. Fuel Processing Technology. 92. 1750-1755.

McCormick R.L., Ratcliff M., Moens L. and Lawrence R. (2007). Several factors affecting the stability of biodiesel in standard accelerated tests. Fuel Processing Technology. 88. 651–657.

Muhammad U.L, Shamsuddin I.M, Danjuma A., Musawa1 R.U. and Dembo U.H. (2018). Biofuels as the Starring Substitute to Fossil Fuels. Petroleum Science and Engineering. 2(1). 44-49.

Noirbent G. and Dumur F. (2020). Recent Advances on Copper Complexes as Visible Light Photoinitiators and (Photo) Redox Initiators of Polymerization. Catalysts. 10 (953). 1-37.

Rashed M. M., Kalam M. A., Masjuki H. H., Rashedul H. K., Ashraful A.M., Shancita I. and Ruhul A.M. (2015). Stability of biodiesel, its improvement and the effect of antioxidant treated blends on engine performance and emission. RSC Advances. 46. 1-60

Razon, Knothe G. and Luis F. (2016). Biodiesel fuels. Progress in Energy and Combustion Science, 58, pp. 36-59.

Riazi H., Shamsabadi A.A., Grady M., Rappe A.M., and Soroush M. (2017). Experimental and Theoretical Study of the Self-Initiation Reaction of Methyl Acrylate in Free-Radical Polymerization. Industrial and Engineering Chemistry Research. 1-29.

Riazi H., Shamsabadi A.A., Grady M., Rappe A.M., and Soroush M. (2019). Method of Moments Applied to Most-Likely High-Temperature Free-Radical Polymerization Reactions. Processes. 1-25.

Saeed, Pullen J. and Khizer. (2012). An overview of biodiesel oxidation stability. Renewable and Sustainable Energy Reviews. 16. 5924-5950.

Sazzad B. S., Fazal M. A., Haseeb A.S.M.A. and Masjuki H.H. (2016). Retardation of oxidation and material degradation in biodiesel: a review. RSC Advances. 65, 1-47

Sharma, Jain S. and M.P. (2009). Stability of biodiesel and its blends: A review. Renewable and sustainable energy reviews. 14. 667-678.

Srinivasan S., Rappe A.M. and Soroush M. (2019). Chapter 4 - Theoretical insights into thermal selfinitiation reactions of acrylates. In computational quantum chemistry. (Edited by) Soroush M. Elsevier. 99-134.

Tang H., Guzman R.C.D., Salley S.O. and Simon K.Y. (2008). The oxidative stability of biodiesel: Effects of FAME composition and antioxidant. Lipid Technology. 20(11). 249-252

Valenga M.G.P, Boschen N.L, Rodrigues P.R.P. and Maia G.A.R, (2019). Agro-industrial waste and *Moringa oleifera* leaves as antioxidants for biodiesel. Industrial Crops and Products. 128. 331-337.

Varatharajan K. and Pushparani D.S. (2017). Screening of antioxidant additives for biodiesel fuels. Renewable

and Sustainable Energy Reviews. 82(3). 2017-2028 Waynick and Andrew J. (2005). Characterization of biodiesel and oxidation products. The Coordinating Research Council. Report.

Xin J., Imahara H. and Saka S. (2009). Kinetics on the oxidation of biodiesel stabilized with antioxidant. Fuel. 88(2). 282-286.

Zuleta E.C., Baena L., Riosa L.A. and Jorge A.C. (2012). The Oxidative Stability of Biodiesel and its Impact on the Deterioration of Metallic and Polymeric Materials: a Review. Journal of Brazilian Chemical Society, 23(12), 2159-2175.