

Evaluation of Piprazine Based Nickel Complex as Better BSA Protein Binder

Bidyut Kumar and PVPS Arun²


¹Department of Chemistry, School of Applied Sciences, Centurion University of Technology and Management, Odisha, India

²Department of Biotechnology, School of Engineering and Technology, Centurion University of Technology and Management, Odisha, India

ABSTRACT

In this work, a Schiff base ligand namely 4-((2-(piprazine-1-yl)ethyl)pent-2-en-2-ol (HL) has been synthesized. Further the reaction between HL and NiCl₂·6H₂O salts has resulted in Schiff base complex with general formula [Ni(L)].Cl. The ligand HL and the corresponding Ni(II) complex was further characterized using various spectroscopic techniques, such as, FT-IR, ESI-MS, and elemental analysis. In the recent scenario, research in the field of protein dynamics is very crucial as these are the major part of the biological systems, and interactions of those macromolecules are responsible for various pathological diseases. Most of the metals inside the biological system shows some kind of affinity to bind with those macromolecules. Herein, it is our unmet need to address the issue by preparing some metal complexes. Therefore, fluorescence spectroscopy was introduced to check the protein binding study of BSA using Stern-Volmer equation.

Graphical Abstract



KEY WORDS: SCHIFF BASE, BSA USING STERN-VOLMER.

INTRODUCTION

Biological processes drives to recognize the fundamental principles by the use of transition metals and its

functionality due to its diverse nature in the quest of the scientist, which ultimately assistances to advance various structural and more significantly functional model systems.[1-3] Besides studying diverse biological procedures these metal ions are tend to retain the assets like antifungal, antibacterial, antiproliferative, antimicrobial and anticancer activity. These transition metal ions also perform a crucial part in terms of structural organization and overall functionality.[4-6].

Moreover, small metal complexes are quite interactive with essential protein is a current topic of main research as there is adequate potential for the progress of new therapeutic agents predominantly screening antitumor

ARTICLE INFORMATION

Received 11th Oct 2020 Accepted after revision 28th Dec 2020
Print ISSN: 0974-6455 Online ISSN: 2321-4007 CODEN: BBRBCA

Thomson Reuters ISI Web of Science Clarivate Analytics USA and Crossref Indexed Journal



NAAS Journal Score 2020 (4.31)
A Society of Science and Nature Publication,
Bhopal India 2020. All rights reserved.
Online Contents Available at: <http://www.bbrc.in/>

activities and the opportunity of conveying these molecules throughout the physiological organization. [6-10] Nickel and copper are two basic examples among the various transition metal ions, which have previously revealed promising capabilities in many of the above stated area.[11-15].

The ability of the serum albumin to bind towards the diverse endogenous and exogenous samples delivers this plasma protein to achieve an effective character for conceivable drug delivery. Thus, the study of the interaction of small molecules with different types of serum albumin is also crucial for the understanding of metallo-pharmaceutical pharmacokinetics and structure-activity relationships. The studies on the interactions between proteins and nickel and copper were not studied much.[16-21]. Thus, we account the synthesis and characterization of a new mononuclear nickel complex [Ni(L)].Cl [where, ligand, HL=1-Phenyl-3-(2-piperazin-1-yl-ethylimino)-but-1-en-1-ol]. The interaction of Ni(II) complex with bovine serum albumin (BSA) also have been considered that displayed favorable consequences with effectual affinity towards the protein.

Experimental Section

MATERIAL AND METHODS

We used the chemicals from Sigma aldrich and applied without additional refinement. A BRUKER TENSOR 27 instrument were recorded Infrared spectra (4000 to 500 cm^{-1}) with using KBr pellets. We used Bruker-Daltonics Mass spectrometric microTOF-Q II mass spectrometer for study. Moreover, the elemental analyses were done with a ThermoFlash 2000 elemental analyzer. Emission spectroscopy were measured in a Horiba JobinYvon (Model: FM-100) made Fluoromax-4p spectrofluorometer from using a 1 cm path length-based quartz cuvette. UV-Visible (200 to 800 nm) spectra were recorded in 0.1 cm path length cell (Hellma, Muellheim/Baden, Germany) using a scan rate of 20 nm min^{-1} and band width of 1 nm. The ligand, HL and corresponding complexes were characterized using various spectroscopic techniques, such as, FT-IR, ESI-MS, and elemental analysis.

Synthesis of 4-((2-(piperazin-1-yl)ethyl)imino)pent-2-en-2-ol (HL): 0.65 g of amino ethyl piperazine (5 mmol) poured into 25 mL of MeOH was mixed with a solution of 0.50 g (5 mmol) of acetyl acetone in 20 mL of methanol. The mixture was then refluxed for 4 h at 60 degree centigrade. A yellow oily compound is formed after evaporating the volatile solvent. Yield: 84%. Anal. Calcd (%): $\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}$: H, 8.48; C, 70.30; N, 15.37; O, 5.85. Found (%): C, 71.08; N, 15.88; O, 6.02. $[\text{C}_{16}\text{H}_{23}\text{N}_3\text{O} + \text{H}]^+$ (m/z) Calculated- 274.18 (m + H)⁺; obtained- 274.18 (m + H)⁺ (Figure 1).

Synthesis of complex, [Ni(L)].Cl: Ligand, HL (0.062 g, 0.25 mmol) and $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (0.58 g, 0.25 mmol) containing 15 mL of MeOH solution was stirred at refluxing condition for 4 h and the producing light green coloured solution. After that, the solution was concentrated by evaporating

the methanol solvent. Green needle shaped crystals were finally obtained after two or three days by keeping the reaction mixture to the slow evaporation at room temperature. Yield: 85%. Anal. Calcd (%): $\text{C}_{16}\text{H}_{21}\text{ClN}_3\text{NiO}_5$: H, 4.93; C, 44.74; N, 9.78. Found (%):H, 5.07; C, 44.64; N, 9.26. $[\text{C}_{16}\text{H}_{21}\text{N}_3\text{NiO}]^+$ (m/z in positive ESI-MS mode) calculated - 330.05 (m)⁺; obtained - 330.10 (m)⁺ for $[\text{Ni(L)}]^+$. Designated infra-red data on KBr pallets (v cm^{-1}): 1603-1604 (C=N), 2951 (N-H) (Figure 2).

Figure 1: ESI-MS of Schiff base ligand (HL)




Figure 2: Electrospray ionisation mass spectrometry data of [Ni(L)].Cl



RESULTS AND DISCUSSION


Synthesis of ligand and complex: The reaction of 1 : 1 molar ratio of amino ethyl piperazine with 1-phenyl-1,3-butanedione in methanol directed to the development of Schiff base ligand, HL (Scheme 1). The ligand can behave as a tetradentate ligand or as a tridentate ligand depending upon its conformation. That is why these types of ligand may be called as flexidentate ligand. Upon reaction of HL with nickel perchlorate in MeOH, a light green colour medium was found. The solution was then dried to the rotary evaporator and the solid compound was collected. A pinch of the sample compound was then dissolved in the acetone for get the nice crystals of [Ni(L)].Cl within 3-4 days at room temperature (Scheme 1).

Scheme 1. Formation of nickel complex with the piperazine moiety.

FTIR Spectroscopy: The IR spectra of nickel complex (Figure 3) have a prominent band around 1603 and 1604 cm^{-1} detectable to $\text{v}(\text{C}=\text{N})$ stretching mode respectively. Furthermore, Ni(II) complex shows intermediate intensity

bands in the range of 2951 to 3443 cm^{-1} because of $\nu(\text{N-H})$ stretching.

Figure 3: FTIR of complex, $[\text{Ni}(\text{L})]\cdot\text{Cl}$.




BSA binding study: Interaction of transition metal complexes with BSA protein are generally checked by the intrinsic fluorescence intensity. Generally, tryptophan, tyrosine, and phenylalanine residues are the main intrinsic component for showing emission intensity of a protein. The fluorescence spectrum of complex (Figure 4) with BSA shows that there is a sequential decrease in the fluorescence intensity. The Stern–Volmer equation is given in the following for determination of various parameters.

$$\frac{F_0}{F} = 1 + k_q \tau_0 [Q] = 1 + K_{SV} [Q]$$

where F and F_0 are the emission intensities in the presence and the absence of a quencher, k_q is the rate constant for the bimolecular quenching, τ_0 is the fluorophore's average life time when the quencher is not present and $[Q]$ is quencher (here complex) concentration. K_{SV} is the Stern-Volmer quenching constant in M^{-1} .

Figure 4: Quenching of the fluorescence of bovine serum albumin BSA by $\text{Ni}(\text{II})$ complex.



CONCLUSION

In summary, a new nickel complex $[\text{Ni}(\text{L})]\cdot\text{Cl}$ having square planer geometry with Schiff base ligand HL have been synthesized and characterized. All the synthesized

ligand and complexes are well characterized using several analytical techniques. The interaction of $\text{Ni}(\text{II})$ complex with BSA protein, which displayed strong interaction with high fluorescence quenching and that is responsible for the interplay between the metal complex and protein.

REFERENCES

- Barnett, S.M., Goldberg, K.I. and Mayer, J.M., 2012. A soluble copper–bipyridine water-oxidation electrocatalyst. *Nature chemistry*, 4(6), pp.498–502.
- Chohan, Z.H., Supuran, C.T. and Scozzafava, A., 2004. Metalloantibiotics: synthesis and antibacterial activity of cobalt (II), copper (II), nickel (II) and zinc (II) complexes of kefzol. *Journal of enzyme inhibition and medicinal chemistry*, 19(1), pp.79–84
- del Campo, R., Criado, J.J., Garcica, E., Hermosa, M.R., Jimenez-Sanchez, A., Manzano, J.L., Monte, E., Rodriguez-Fernández, E. and Sanz, F., 2002. Thiourea derivatives and their nickel (II) and platinum (II) complexes: antifungal activity. *Journal of Inorganic Biochemistry*, 89(1-2), pp.74–82.
- Enyedy, É.A., Horváth, L., Hetényi, A., Tuccinardi, T., Hartinger, C.G., Keppler, B.K. and Kiss, T., 2011. Interactions of the carrier ligands of antidiabetic metal complexes with human serum albumin: A combined spectroscopic and separation approach with molecular modeling studies. *Bioorganic & medicinal chemistry*, 19(14), pp.4202–4210.
- Kalaivani, P., Saranya, S., Poornima, P., Prabhakaran, R., Dallemer, F., Padma, V.V. and Natarajan, K., 2014. Biological evaluation of new nickel (II) metallates: Synthesis, DNA/protein binding and mitochondrial mediated apoptosis in human lung cancer cells (A549) via ROS hypergeneration and depletion of cellular antioxidant pool. *European Journal of Medicinal Chemistry*, 82, pp.584–599.
- Kasuga, N.C., Sekino, K., Koumo, C., Shimada, N., Ishikawa, M. and Nomiya, K., 2001. Synthesis, structural characterization and antimicrobial activities of 4- and 6-coordinate nickel (II) complexes with three thiosemicarbazones and semicarbazone ligands. *Journal of inorganic biochemistry*, 84(1-2), pp.55–65.
- Krishnamoorthy, P., Sathyadevi, P., Butorac, R.R., Cowley, A.H., Bhuvanesh, N.S. and Dharmaraj, N., 2012. Variation in the biomolecular interactions of nickel (II) hydrazone complexes upon tuning the hydrazide fragment. *Dalton Transactions*, 41(22), pp.6842–6854.
- Lauria, A., Bonsignore, R., Terenzi, A., Spinello, A., Giannici, F., Longo, A., Almerico, A.M. and Barone, G., 2014. Nickel (II), copper (II) and zinc (II) metallo-intercalators: structural details of the DNA-binding by a combined experimental and computational investigation. *Dalton Transactions*, 43(16), pp.6108–

6119.

- Leung, C.H., Zhong, H.J., Chan, D.S.H. and Ma, D.L., 2013. Bioactive iridium and rhodium complexes as therapeutic agents. *Coordination Chemistry Reviews*, 257(11-12), pp.1764-1776.
- Li, X., Gorle, A.K., Sundaraneedi, M.K., Keene, F.R. and Collins, J.G., 2018. Kinetically-inert polypyridylruthenium (II) complexes as therapeutic agents. *Coordination Chemistry Reviews*, 375, pp.134-147.
- Ma, D.L., Wang, M., Mao, Z., Yang, C., Ng, C.T. and Leung, C.H., 2016. Rhodium complexes as therapeutic agents. *Dalton Transactions*, 45(7), pp.2762-2771.
- Ma, D.L., Wu, C., Li, G., Yung, T.L. and Leung, C.H., 2020. Transition metal complexes as imaging or therapeutic agents for neurodegenerative diseases. *Journal of Materials Chemistry B*.
- Rafique, S., Idrees, M., Nasim, A., Akbar, H. and Athar, A., 2010. Transition metal complexes as potential therapeutic agents. *Biotechnology and Molecular Biology Reviews*, 5(2), pp.38-45.
- Ramachandran, E., Raja, D.S., Mike, J.L., Wagner, T.R., Zeller, M. and Natarajan, K., 2012. Evaluation on the role of terminal N-substitution in 6-methoxy-2-oxo-1, 2-dihydroquinoline-3-carbaldehyde thiosemicarbazones on the biological properties of new water-soluble nickel (II) complexes. *RSC advances*, 2(22), pp.8515-8525.
- Saeidifar, M., Mansouri-Torshizi, H. and Saboury, A.A., 2015. Biophysical study on the interaction between two palladium (II) complexes and human serum albumin by Multispectroscopic methods. *Journal of Luminescence*, 167, pp.391-398.
- Sathyadevi, P., Krishnamoorthy, P., Jayanthi, E., Butorac, R.R., Cowley, A.H. and Dharmaraj, N., 2012. Studies on the effect of metal ions of hydrazone complexes on interaction with nucleic acids, bovine serum albumin and antioxidant properties. *Inorganica Chimica Acta*, 384, pp.83-96.
- Shaik, S., 2010. Biomimetic chemistry: Iron opens up to high activity. *Nature Chemistry*, 2(5), p.347.
- Tietze, D., Tischler, M., Voigt, S., Imhof, D., Ohlenschläger, O., Görlach, M. and Buntkowsky, G., 2010. Development of a Functional cis-Prolyl Bond Biomimetic and Mechanistic Implications for Nickel Superoxide Dismutase. *Chemistry—A European Journal*, 16(25), pp.7572-7578.
- Topala, T., Bodoki, A., Oprean, L. and Oprean, R., 2014. Bovine serum albumin interactions with metal complexes. *Clujul medical*, 87(4), p.215.
- Trynda-Lemiesz, L., Karaczyn, A., Keppler, B.K. and Kozłowski, H., 2000. Studies on the interactions between human serum albumin and trans-indazolium (bisindazole) tetrachlororuthenate (III). *Journal of inorganic biochemistry*, 78(4), pp.341-346.
- Zhang, Y.P., Li, Y., Xu, G.C., Li, J.Y., Luo, H.Y., Li, J.Y., Zhang, L. and Jia, D.Z., 2019. Synthesis, crystal structure, DNA/bovine serum albumin binding and antitumor activity of two transition metal complexes with 4-acylpyrazolone derivative. *Applied Organometallic Chemistry*, 33(3), p.e4668.