

Assessment of Avian Road Kill Mortality in the State Highway Passing Through Agricultural Landscape

Amol S. Rawankar^{1*} and Gajanan A. Wagh²

¹Department of Zoology, Jagadamba Mahavidyalaya Achalpur, Amravati, Maharashtra, India

²Department of Zoology, Shri Shivaji Science College, Amravati, Maharashtra, India

ABSTRACT

Roads are becoming one of the greater threats to avian fauna. In India, very few studies have been carried out to assess the mortality of avian fauna and other vertebrates. The data on the road kill survey was collected from Jan 2015 to Dec 2017. For the survey of a road-killed avian fauna, the road passing through the agricultural landscape (from Amravati to Paratwada) was selected based on geographical location and availability of vegetation diversity. The survey found a total of 694 carcasses of 38 different species belonging to 25 families that were killed on Amravati to Paratwada Highway. The highest mortality of Greater Coucal (*Centropus sinensis*) was recorded on Amravati to Paratwada state highway. While studying the seasonal variations, it was observed that the maximum mortality was found from June to September and the minimum from February to May.

KEY WORDS: AVIAN MORTALITY, AGRICULTURAL LANDSCAPE, ROAD VEHICLE COLLISION, STATE HIGHWAY.

INTRODUCTION

In this era of industrialization and modernization, the activities of humans affect the population of avian fauna. There are approximately 80 million birds are killed due to collisions with road vehicles every year in the United States alone (Erickson et al. 2005). Highways through wildlife reservoirs affect the fauna seriously and the effects range from habitat loss and fragmentation, direct mortality through collision with vehicles (Foster and Humphrey 1995; Das et al. 2007; Row et al. 2007; Baskaran and Boominathan 2010).

For birds and other vertebrate groups, a direct threat of roads is death due to collisions with vehicles (Erritzoe et al. 2003). Increasing traffic speed and volume have been found to also increase mortality and the mortality rates are generally highest during summer and Spring (Case 1978; Loss et al. 2014b). Mortality rates have been found to increase with the increasing width of the road corridor (Oxley et al. 1974; Pintoa et al. 2020). Pintoa et al. (2020) studied the temporal patterns of bird mortality due to road traffic collisions in the Mediterranean region. This study provides evidence that bird mortality due to road vehicle collisions may change

between years and be highest during the breeding season (Pintoa et al. 2020).

Migratory species travel long distances and they may be even more at risk because they are presumably exposed to more road crossing events than non-migratory species (Harris and Scheck 1991). Higher mortality of adult and juvenile birds having roadside habitats can create sink populations that can only persist through immigration (Mumme et al. 2000; Bishop and Brogan 2013). Betleja et al. (2020) studied how high birds fly above roads, and how they use the road infrastructure such as bridges, lampposts, etc. This study found that species differed significantly in the height at which they crossed over roads, and 30 % of the birds crossing were at heights below 12 m, which is a potential collision height (Betleja et al. 2020). Roads are becoming one of the greater threats to animal and plant populations. In India, the highways are going through many protected areas and cause a severe impact on wildlife and their habitat (Vijayakumar et al. 2001). In India, very few studies have been carried out to assess the mortality of avian fauna and other vertebrates on the roads passing through reserve forests (Chhangani 2004; Das et al. 2007; Baskaran and Boominathan 2010; Selvan et al. 2012; Rawankar and Wagh 2018; Betleja et al. 2020).

These studies provide the data on various casualties due to the vehicular collision in birds and other vertebrate fauna

Article Information:*Corresponding Author: amol_rawankar@yahoo.co.in
Received 10/03/2022 Accepted after revision 26/06/2022
Published: 30th June 2022 Pp- 301-308
This is an open access article under Creative Commons License,
<https://creativecommons.org/licenses/by/4.0/>.
Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/15.2.6>

in the different parts of India and also suggest the urgent need to undertake such a survey to estimate the magnitude of this threat in different parts of India.

Figure 1: Map showing Amravati to Paratwada State Highway passing through agricultural land (Satellite image).




Figure 2: State Highway from Amravati to Paratwada passing through agricultural land (Actual Photograph)



MATERIAL AND METHODS

The Amravati to Paratwada State Highway Number – 6 is passing through the agricultural landscape and large numbers of vehicles are passing through this highway at very high speed every day. Hence this study was carried out to estimate the avian mortality due to road vehicle collisions on this state highway.

Road Vehicle collision Study: The data of the roadkill survey was collected from Jan 2015 to Dec 2017. For the survey of avian roadkill, Amravati to Paratwada, 50-kilometer state Highway Number – 6 was selected which is passing through the agricultural landscape. The survey was performed using a motorcycle. The motorcycle was ride at a speed of 10- 20 km/hour on the roads. The survey was performed in the morning hours from 7.00 to 9.00 am. The road, as well as 5 feet, fix area along the side of the road, was screened out for avian carcasses. When the dead bird was encountered, a photograph of the bird was taken. The survey was performed twice a week on this road. This survey method was followed by various researchers and found satisfactory in evaluating the road kills. (Das 2007; Baskaran 2010; Selvan 2012; Betleja et al. 2020).

Once the roadkill was encountered and photographed then it was removed from the road to prevent the double counting. Taxonomic keys were used to identify the road kills. Up to genus and species level identification was done. The road kills reported by locals, birders, and the author's occasional visits and road kills data were also recorded (Grimmat et al. 2009; Rasmussen and Anderton 2012; Wadatkar et al. 2016; Betleja et al. 2020).

The mortality estimation is subject to multiple biases and hence to calculate the annual mortality estimate accurately, the biases such as observer bias (probability of detection) and scavenging activity (persistence time) are considered. To calculate the mortality estimate, the formula provided by Gerow et al. was used (Tajera et al. 2018; Betleja et al. 2020).

$$M = (C / nS) * (I / T_p)$$

Where M is the estimated mortality; C is the number of carcasses found; nS is the number of surveys conducted (nS = 104), I is the number of days between surveys (I = 03); T is the persistence time of each taxon (T = 2.2 days), and p is the probability of detection (p = 67 %). While calculating the mortality per km (with or without bias), an average of three years was taken into consideration. Standard deviation was calculated for each road-killed bird species on Amravati to Paratwada State highway during the year 2015 to 2017 (Betleja et al. 2020).

RESULTS AND DISCUSSION

The survey found a total of 694 carcasses of 37 different avian species belonging to 25 families that were killed by road vehicles collision on the Amravati to Paratwada Highway. The highest mortality of Greater Coucal (*Centropus sinensis*) was encountered, and followed by Red - vented Bulbul (*Pycnonotus cafer*), Spotted Owlet (*Athene brama*), Common Myna (*Acridotheres tristis*), House Crow (*Acridotheres tristis*), Indian Nightjar (*Caprimulgus asiaticus*), Ashy Prinia (*Prinia socialis*), Indian Roller (*Coracias benghalensis*), Yellow-eyed Babbler (*Chrysomma sinense*), Laughing Dove (*Spilopelia senegalensis*), Jungle Babbler (*Turdoides striata*), Asian Koel (*Eudynamis scolopaceus*), Common Stonechat (*Saxicola torquatus*), Jungle Prinia (*Prinia sylvatica*), Indian Pond Heron (*Ardeola grayii*), Red-wattled Lapwing (*Vanellus indicus*), Barred buttonquail (*Turnix suscitator*), Indian Robin (*Copsychus fulicatus*), Baya Weaver (*Ploceus philippinus*), Indian Thick-knee (*Burhinus indicus*), Common Tailor Bird (*Orthotomus sutorius*), Pied Cuckoo (*Clamator jacobinus*), Rufous Treepie (*Dendrocitta vagabunda*), Coppersmith Barbet (*Psilopogon haemacephalus*), Barn Owl (*Tyto alba*), Large Grey Babbler (*Turdoides malcolmi*), Indian Scops Owl (*Otus bakkamoena*), Yellow Wagtail (*Motacilla flava*), Scaly-breasted Munia (*Lonchura punctulata*), Silver bill Munia (*Euodice malabarica*), Wire - tailed Swallow (*Hirundo smithii*), Red - rumped Swallow (*Hirundo daurica*), Common House Martin (*Delichon urbicum*), Long-Tailed Shrike (*Lanius schach*) and Tawny bellied Babbler (*Dumetia hyperythra*).

Table 1. Showing the species-wise mortality and its percentage to the total carcasses killed due to a road vehicle collision and standard deviation for each species, on Amravati to Paratwada State highway during the year 2015-2017

Scientific Name	Common Name	2015	2016	2017	Total (%)	SD
<i>Centropus sinensis</i>	Greater Coucal	29	65	79	173 (24.93)	25.79
<i>Athene brama</i>	Spotted Owlet	16	16	24	56 (8.07)	4.62
<i>Pycnonotus cafer</i>	Red-vented Bulbul	16	18	20	54 (7.78)	2.00
<i>Acridotheres tristis</i>	Common Myna	11	13	14	38(5.48)	1.53
<i>Corvus splendens</i>	House Crow	11	9	8	28(4.03)	1.53
<i>Caprimulgus asiaticus</i>	Indian Nightjar	12	9	6	27 (3.89)	3.00
<i>Coracias benghalensis</i>	Indian Roller	9	3	10	22 (3.17)	3.79
<i>Spilopelia senegalensis</i>	Laughing Dove	8	6	8	22 (3.17)	1.15
<i>Prinia socialis</i>	Ashy Prinia	10	4	7	21 (3.03)	3.00
<i>Chrysomma sinense</i>	Yellow-eyed Babbler	9	7	3	19 (2.74)	3.06
<i>Prinia sylvatica</i>	Jungle Prinia	6	6	4	16 (2.31)	1.15
<i>Turdoides striata</i>	Jungle Babbler	7	2	6	15 (2.16)	2.65
<i>Ardeola grayii</i>	Indian Pond Heron	6	4	5	15 (2.16)	1.00
<i>Burhinus indicus</i>	Indian Thick-knee	6	5	4	15 (2.16)	1.00
<i>Vanellus indicus</i>	Red-wattled Lapwing	6	4	2	12 (1.73)	2.00
<i>Psilopogon haemacephalus</i>	Coppersmith Barbet	5	5	2	12 (1.73)	1.73
<i>Tyto alba</i>	Barn Owl	4	2	6	12 (1.73)	2.00
<i>Eudynamys scolopaceus</i>	Asian Koel	7	2	2	11 (1.59)	2.89
<i>Saxicola torquatus</i>	Common Stonechat	7	4		11 (1.59)	2.12
<i>Orthotomus sutorius</i>	Common Tailor Bird	5	5	1	11 (1.59)	2.31
<i>Turdoides malcolmi</i>	Large Grey Babbler	4	5	1	10 (1.44)	2.08
<i>Copsychus fulvicatus</i>	Indian Robin	6	0	3	9 (1.3)	3.00
<i>Clamator jacobinus</i>	Pied Cuckoo	5	3	1	9 (1.3)	2.00
<i>Otus bakkamoena</i>	Indian Scops Owl	2	5	1	8 (1.15)	2.08
<i>Ploceus philippinus</i>	Baya Weaver	6	0	1	7 (1.01)	3.21
<i>Turnix suscitator</i>	Barred buttonquail	6	0	0	6 (0.86)	3.46
<i>Dendrocitta vagabunda</i>	Rufous Treepie	5	0	0	5 (0.72)	2.89
<i>Bubulcus ibis</i>	Cattle Egret	0	5	0	5 (0.72)	2.89
<i>Lonchura punctulata</i>	Scaly-breasted Munia	0	2	0	2 (0.29)	1.15
<i>Merops orientalis</i>	Green Bee-eater	0	2	0	2 (0.29)	1.15
<i>Euodice malabarica</i>	Silver bill Munia	0	1	0	1 (0.14)	0.58
<i>Motacilla flava</i>	Yellow Wagtail	0	0	1	1 (0.14)	0.58
<i>Hirundo smithii</i>	Wire-tailed Swallow	0	0	1	1 (0.14)	0.58
<i>Hirundo daurica</i>	Red-rumped Swallow	0	0	1	1 (0.14)	0.58
<i>Delichon urbicum</i>	Common House Martin	0	0	1	1 (0.14)	0.58
<i>Lanius schach</i>	Long Tailed Shrike	0	0	1	1 (0.14)	0.58
<i>Dumetia hyperythra</i>	Tawny-bellied Babbler	0	0	1	1 (0.14)	0.58
	Unidentified Birds	7	17	10	34(4.9)	5.13
	Total	231	229	234	694	


The Family wise mortality of avian species was also calculated. Highest mortality was encountered in the Centropodidae and then followed by Pycnonotidae, Strigidae, Sturnidae, Corvidae, Caprimulgidae, Cisticolidae, Coraciidae, Sylviidae, Columbidae, Leiotrichidae, Cuculidae, Muscicapidae, Ardeidae, Charadriidae, Turnicidae, Ploceidae, Burhinidae, Megalaimidae, Tytonidae, Estrildidae, Motacillidae, Hirundinidae, Laniidae, and Timaliidae. The year wise mortality of the

birds due to road vehicular collisions are shown in Table No. 1 and season wise details in Fig. 3.

Carcasses of Greater Coucal were found more frequently than other reported bird carcasses on the studied highway and perhaps the most dominant species on all the roads of Vidarbha and Maharashtra. This bird species cannot make pace with the speed of the vehicle because of its large size, low-level flight pattern, and ground-dwelling habit

and hence gets killed in collision with the vehicle. Indian Roller and Common Myna are the tree cavity-nesting birds. These birds prefer the holes of the Neem plant (*Azadirachta indica*) and Babul plant (*Acacia nilotica*) along the roadside. The frequency of road vehicle collisions of Indian Roller and Common Myna was observed maximum in summer because these birds' activities get increased near the nest site. This observation correlates with a previously published study (Erritzoe et al. 2003; Betleja et al. 2020).

Figure 3: Season wise Avian Mortality due to road vehicle collision from 2015 - 2017 on Amravati - Paratwada Highway




Owls, owlets (Spotted Owlet), and Nightjars are the nocturnal bird species. As Owls and Owlets feed on mice, rats, and other small animals, they used roadside trees for roosting. These birds are also tree cavity-nesting birds and they become active during the night, but due to high-intensity LED flashlights of vehicles they cannot make pace with the speed of the vehicle and get killed. Spotted Owlet and nightjars are the most frequently killed species of birds among the nocturnal birds (Betleja et al. 2020).

While studying the seasonal variations, it is observed that the maximum mortality was found during the months of June to September and the minimum in the months of Feb to May. The majority of the bird species have a breeding season during the rainy months from June to September. During the breeding season, the movements of the bird's increase, and hence the chances of collision also increase. Pintoa et al. (2020) studied the temporal patterns of bird mortality due to road traffic collisions in the Mediterranean region and their study also found that bird mortality due to road vehicle collisions was highest during the breeding season. Betleja et al. (2020) studied how high birds fly above roads, and how they use road infrastructure such as bridges, lampposts, etc. This study found that species differed significantly in the height at which they crossed over roads, and 30 % of the birds crossing were at heights below 12 m, which is a potential collision height (Betleja et al. 2020).

Due to rain, the wings of the birds get wetted, and hence the weight of the wings increases, and birds require more energy to fly. Due to this reason, birds find difficulty in flying due to wet body feathers. Again during monsoon, the frogs, toads, snakes, lizards, and small mammals come on roads

in more number, and the birds which feed on these animals also come on roads frequently to catch these animals. Due to this reason, the mortality of nocturnal raptors (Owls and Owlets) get increased during the monsoon season. On Amravati to Paratwada State Highway, the estimated mean annual mortality was found 4.62 birds/km/year, and when recalculated by considering the biases such as observer bias and scavenging activity it is found 12.44 birds/km/year (Betleja et al. 2020).

Photo Plate No 1: Photographs of Carcasses of the birds Killed Due to Road vehicle Collision



Various study results show that the most dominant species found dead due to road vehicle collision belong to the family passerines, Columbidae, nocturnal birds such as owlets and nightjars. On the contrary, the present study recorded the observations that the Greater Coucal was the most dominant species belonging to the family Centropodidae found dead on the roads due to the vehicular collision. This scientific report presents that the Greater Coucal is the most dominant species found dead due to the road vehicle collision on State highways passing through agricultural landscape (Dhindsa et al. 1988; Chhangani 2004; Erickson et al. 2005; Selvan et al. 2012; Betleja et al. 2020).

Photo Plate No2: Photographs of Carcasses of the birds Killed Due to Road vehicle Collision



Jungle Babbler
(*Turdoides striata*)




Plain Prinia
(*Prinia inornata*)



Indian Roller
(*Coracias benghalensis*)




Yellow-eyed Babbler
(*Chrysomma sinense*)



Asian Koel
(*Eudynamis scolopaceus*)



House Crow
(*Corvus splendens*)



Indian Scops Owl
(*Saxicola torquatus*)



Common Stonechat
(*Otus bakkamoena*)

Erritzoe et al. (2003) and Boves and Belthoff (2012) reported that the passerines and Owls are the most dominant among the road-killed species. The present study also recorded the significant mortality of Passerines and Owls due to road vehicle collisions. When the data of road vehicle collision was evaluated based on feeding habitat (Table No. 2), it is found that the maximum number of species were insectivorous (21 species) followed by Omnivorous (06 species), Carnivorous (4 species), Granivorous (3 Species), Frugivorous (2 Species) and Piscivorous (1 species). The insectivorous and omnivorous birds do come on the roads to feed on the small insects and other animals such as amphibians, snakes, crabs, etc., and hence the chance of

their collision with vehicles increases. The granivorous birds come on the roads to feed on the grains dispersed on the roads. Frugivorous birds such as Red Vented Bulbul, Brahmany Starling, and Coppersmith Barbet feed on the fruits of the roadside trees and collide with vehicles when trying to cross the roads. In the Piscivorous bird category, the study found only one species which is the Indian Pond Heron. This bird mainly feeds on fishes, frogs, aquatic insects, etc. The record of these species as roadkill may be a matter of chance only (Betleja et al. 2020).

When a comparison was made between diurnal and nocturnal birds (Table No.2), it is found that the majority of the species which got killed due to road vehicle collisions are diurnal (33 species) and only 4 species were nocturnal. The nocturnal bird species include the Owls (Barn Owl, Scops Owl), Owllet (Spotted Owllet), and Indian Nightjar. The mortality in diurnal birds occurred during the early hours of the day when the diurnal birds are more active and the speed of the vehicle is very high. The mortality in nocturnal birds such as Owls, owllets, and Nightjar occurred during the night-time (Betleja et al. 2020).

Following recommendations will be useful to minimize road vehicle collision and avian species conservation.

1. It is recommended that limiting the speed of the vehicle on the highways passing through the critical areas to 40 to 50 km/hour (Orlowski and Nowak, 2006), further speed breakers and signposts at the roadsides should be introduced in Wildlife conservation policies.
2. Structural elements can encourage birds to fly above traffic or below the road through bridges or culverts. Flight diversion works best for species with direct, rapid flight. Poles that produce an illusion of a solid barrier were effective in reducing bird road kills.
3. Vehicle LED lighting appears to blind Owls and increases the chances of collision with vehicles hence light pollution could be addressed as a part of the strategy.
4. The combined efforts and continuous monitoring of Foresters, Non-Government Organisations (NGO), and Birders are needed for the conservation of the Avian Diversity.

This study provides baseline data on the magnitude of avian mortality due to road vehicle collisions on the Amravati to Paratwada state highway in the Amravati District. Due to such loss of these species, the total avian population becomes severely affected. The number of mortality due to road vehicle collisions may be quite large because the avian carcasses get cleared immediately by scavengers after death and remain unnoticed; sometimes the collided birds have dashed far away from the road and cannot be noticed. Further, this was the first attempt to estimate the mortality of avian fauna due to road vehicle collision on State highway no.6 Amravati to Paratwada (50Km) in the Amravati district. The better estimation and evaluation of avian carcasses need long-term study (Betleja et al. 2020).

Table 2. Showing feeding habits and diurnal/ Nocturnal habit of birds killed by road vehicle collision Amravati - Paratwada State Highway

Sr. No	Scientific Name	Insectivorous	Granivores	Frugivorous	Carnivorous	Omnivorous	Piscivores	Diurnal	Nocturnal
1	<i>Prinia socialis</i>	+						+	
2	<i>Eudynamys scolopaceus</i>	+						+	
3	<i>Turnix suscitator</i>					+		+	
4	<i>Ploceus philippinus</i>		+					+	
5	<i>Bubulcus ibis</i>	+						+	
6	<i>Tyto alba</i>				+				+
7	<i>Delichon urbicum</i>	+						+	
8	<i>Acridotheres tristis</i>					+		+	
9	<i>Saxicola torquatus</i>	+						+	
10	<i>Orthotomus sutorius</i>	+						+	
11	<i>Psilopogon haemacephalus</i>			+				+	
12	<i>Centropus sinensis</i>	+						+	
13	<i>Merops orientalis</i>	+						+	
14	<i>Corvus splendens</i>					+		+	
15	<i>Caprimulgus asiaticus</i>	+							+
16	<i>Ardeola grayii</i>						+	+	
17	<i>Copsychus fulicatus</i>	+						+	
18	<i>Coracias benghalensis</i>	+						+	
19	<i>Otus bakkamoena</i>				+				+
20	<i>Burhinus indicus</i>				+			+	
21	<i>Turdoides striata</i>	+						+	
22	<i>Prinia sylvatica</i>	+						+	
23	<i>Turdoides malcolmi</i>	+						+	
24	<i>Spilopelia senegalensis</i>					+		+	
25	<i>Lanius schach</i>					+		+	
26	<i>Clamator jacobinus</i>	+						+	
27	<i>Pycnonotus cafer</i>			+				+	
28	<i>Cecropis daurica</i>	+						+	
29	<i>Vanellus indicus</i>	+						+	
30	<i>Dendrocitta vagabunda</i>					+		+	
31	<i>Lonchura punctulata</i>		+					+	
32	<i>Euodice malabarica</i>		+					+	
33	<i>Athene brama</i>				+				+
34	<i>Dumetia hyperythra</i>	+						+	
35	<i>Hirundo smithii</i>	+						+	
36	<i>Chrysomma sinense</i>	+						+	
37	<i>Motacilla flava</i>	+						+	

+ = Present

CONCLUSION

The findings of the present study recorded the avian mortality of 37 bird species due to road vehicle collision on Amravati to Paratwada state highway, which is passing through the agricultural landscape. Greater Coucal, Red-vented Bulbul, Spotted Owlet, Common Myna, Nightjars, Indian Roller, Dove, and Babblers were found the most affected agricultural habitats dependent bird species. But Greater Coucal was reported the most dominant species affected due to this road vehicle collision threat. The

maximum mortality of bird species was recorded during the rainy seasons and the minimum during the summer season. This report presents the first-ever detailed quantitative study on avian mortality due to road vehicle collisions on state highways passing through agricultural landscapes.

ACKNOWLEDGEMENTS

The research fellowship was provided by the Council of Scientific and Industrial Research (CSIR). The invaluable support was provided by Dr. J. S. Wadtkar, Mr. Jagdev

Iwane, Mr. Hayat Qureshi, Mr. Prathmesh Tiwari and all other Wildlife and Environment Conservation Society (WECS) volunteers.

Conflict of Interests: Authors declare no conflict of interests to disclose.

Data availability Statement: All data / results / information is available with the authors and can be shared on a reasonable request made to the corresponding author when required.

REFERENCES

- Baskaran N. and Boominathan D. (2010). Road kill of animals by highway traffic in the tropical forests of Mudumalai Tiger Reserve, southern India. *Journal of Threatened Taxa* 2(3): 753-759. DOI: <https://doi.org/10.11609/JoTT.o2101.753-9>
- Betleja J., Łukasz J., Sparks T. H. et al. (2020). Birds crossing over roads: species, flight heights and infrastructure use. *European Journal of Ecology*, 6(2).m <https://doi.org/10.17161/eurojocol.v6i2.14788>
- Bishop C.A. and Brogan J. M. (2013). Estimates of avian mortality attributed to vehicle collisions in Canada. *Avian Conservation and Ecology* 8(2):2. <http://dx.doi.org/10.5751/ACE-00604-080202>
- Boves T. J. and Belthoff J. R. (2012). Roadway mortality of Barn owls in Idaho, USA. *The Journal of Wildlife Management* 76:1381–1392. <https://doi.org/10.1002/jwmg.378>
- Case R. M. (1978). Interstate highway road-killed animals: a data source for biologists. *Wildlife Society B* 6: 8- 13.
- Chhangani A. K. (2004). Frequency of avian road – kills in Kumbhalgarh Wildlife Sanctuary, Rajasthan, India. *Forktail* 20: 110-111.
- Das A., Ahmed, M. F., Lahkar, B.P. et al. (2007). A pre-liminary report of reptilian mortality on road due to vehicular movement near Kaziranga National Park, Assam, India. *Zoos' Print Journal* 22(7): 2742–2744. <http://doi.org/10.11609/jott.3001.9.3.10004-10010>
- Dhindsa M.S., Sandhu, J. S., Sandhu, P. S. et al. (1988). Road- side birds in Punjab (India): relation to mortality from vehicles. *Environment Conservation* 15: 303–310
- Erickson W.P., Johnson, G.D. and Young, D.P. Jr. (2005). A summary and comparison of bird mortality from anthropogenic causes with an emphasis on collisions. *USDA Forest Serv. Gen. Tech. Rep. PSW-GTR* .191, 1029–1042.
- Erritzoe J., Mazgajski ,T. D., and Rejt, L. (2003). Bird casualties on European roads – a review. *Acta Ornithologica*, 38:77–93.
- Foster M.L. and Humphrey (1995). Use of highway under-passes by Florida Panthers and other Wildlife. *Wildlife Society Bulletin* 23(1): 95-100. <http://www.jstor.org/stable/3783202>
- Gerow K, Kline, N. C., Swann, D. E., et al. (2010). Estimating Annual Vertebrate Mortality On Roads At Saguaro National Park, Arizona. *Human–Wildlife Interactions*, 4(2):283–292
- Grimmet R., Inskipp, C. and Inskipp, T. (2009). *Birds of the Indian Subcontinent*. Oxford University Press. Edition 5, pp 1-384.
- Harris L. D. and Scheck J. (1991). From implications to applications: the dispersal corridor principle applied to conservation of biological diversity. D. A. Saunder and R. J. Hobbs, editors. *Nature conservation the role of corridors*. Surrey Beatty and Sons, Chipping Norton, Australia. 189-220 Pp..
- Loss S.R., Will T. and Marra, P. P (2014b). Estimation of annual bird mortality from vehicle collisions on roads in the United States. *Journal of Wildlife Management* 78:763–71. <https://doi.org/10.1002/jwmg.721>
- Mumme R. L., Schoech, S. J., Woolfenden, G. E. et al. (2000). Life and death in the fast lane: demographic consequences of road mortality in the Florida Scrub-Jay. *Conservation Biology*, 14:501–512.
- Orlowski G. and Nowak, L. (2006). Factors influencing mammal roadkills in the agricultural landscape of south-western Poland. *Polish Journal of Ecology* 54(2):283-294.
- Oxley D.J., Fenton, M.B., and Carmody, G.R. (1974). Effects of roads on populations of small mammals. *Journal of Applied Ecology* , 11 : 51–59.
- Pandora P., Rui L., António M. et al. (2020). Temporal patterns of bird mortality due to road traffic collisions in a Mediterranean region, *Bird Study*, 67 (1).DOI: 10.1080/00063657.2020.1779652
- Rasmussen P.C. and Anderton, J.C (2012). *Birds of South Asia*. The Ripley Guide. Vols. 1 and 2, 2nd edition. National Museum of natural history – Smithsonian Institution, Michigan State University and Lynx Edicions, Washington , D.C Michigan and Barcelona.
- Rawankar A. S and Wagh, G. A. (2018). Surveillance of Avian Mortality on the road passing through Pohra – Malkhed Reserve forest, Dist Amravati, Maharashtra. *International Journal of Zoology Studies* 3(2):135 - 139
- Row J. R., Blouin-Demers, G. and Wheatherhead, P.J. (2007). Demographic effect of road mortality in black rat snakes (*Elaphe obsoleta*). *Biological Conservation*, 137: 117-124.
- Selvan K. M., Sridharan, N. and John, S. (2012). Roadkill animals on national highways of Karnataka, India. *Journal of Ecology and the Natural Environment* ,4(14): 363-365

Tejera G., Rodriguez, B., Armas, C., et al. (2018). Wildlife-vehicle collisions in Lanzarote Biosphere Reserve, Canary Islands. PLoS ONE, 13(3): e0192731. <https://doi.org/10.1371/journal.pone.0192731>

Vijayakumar S.P., Vasudevan, K. and Ishwar, N.M. (2001).

Herpetofaunal mortality on the roads in the Anamalai Hills, southern Western Ghats. Hamadryad, 26(2): 265-272.

Wadatkar J., Kasambe, R., Wagh, G., et al. (2016). Checklist of Birds of Amravati District, Maharashtra WECS Amravati. PP.1-22