

Bacteriological and Physicochemical Evaluation of Selected Bottled Water Brands in Nasarawa, Kano State, Nigeria

Mustapha Abdulsalam,¹ Hafsa M. Ahmad², Olaitan L S³ and Adeyemi O A⁴

^{1,2}Department of Microbiology, Skyline University Nigeria

³Department of Microbiology, University of Gothenburg, Sweden

⁴Department of Microbiology, Ajayi Crowther University, Oyo.

ABSTRACT

Twenty-six different brands of bottled water were tested for bacteriological and physicochemical qualities in Nasarawa Local Government Area of Kano State, Nigeria. Bacteriological assessments have been used to outline the wide varieties of heterotrophic bacteria, overall coliform and faecal coliform. The chloride content, pH, temperature, conductivity, fluoride content, total hardness and suspended particles have been additionally investigated. The coliform count ranged from 0 to 1000 coliform/100ml of the sample while the heterotrophic bacterial count ranged from 0 to 118 cfu/ml. In the bottled water samples, there was no faecal coliform and fluoride contents. The pH, temperature, conductivity, Chloride content, total hardness, and total suspended particles were all within acceptable limits between 6.50 and 8.30, 19.8 and 21.00C, 63.10 and 86.92 μ S/cm, 3.58 and 4.64 mg/l, 1 and 86 mg/l and 1.0 x 10⁻³ to 3.0 x 10⁻³ g/100ml, respectively. *Staphylococcus*, *Aeromonas*, *Klebsiella*, *Micrococcus* and *Bacillus* were among the bacterial species found. For heterotrophic bacteria, total coliform and faecal coliform, 25 (96%), 24 (92%) and 0 (100%) of the 26 different bottled water samples met the standard respectively. For pH, temperature, conductivity, fluoride content, chloride content, total hardness and total suspended particles, all of the bottled water samples also met the criteria. According to the results of this study, 88.46 percent of the bottled waters tested were fit for consumption based on bacteriological and physicochemical factors. When producing bottled water, it is recommended that stringent criteria should be followed to ensure that the bottles are well capped, and that statistics including the producing date, expiration date and consignment number be imprinted on the bottles.

KEY WORDS: SUSPENDED PARTICLES, HETEROTROPHIC BACTERIA, PHYSICOCHEMICAL, BOTTLED WATER.

INTRODUCTION

Despite decades of government efforts, safe and clean water supply in Nigeria's urban areas remains objectionable. In most developing countries, the common method for delivering municipal-level, safe drinking water via treated pipe-borne water is woefully inadequate. (Othmani et al 2021), The importance of water quality to public health cannot be overstated. Water is responsible for the transmission of many infectious illnesses via the faecal-oral pathway. Drinking water-borne diseases claim the lives of approximately five million youngsters every years and sicken one-sixth of the world's population (Yang et al 2021 Amanda et al 2022).

The significance and contribution of neighborhood produce low-fee opportunity ingesting water schemes

to accomplishing sustainable get entry to in growing countries' rural and urban settings can't be over emphasized, particularly in the light of renewed global responsibilities to the Millennium Development Goals (MDGs) set for 2015. Bottled water is widely accessible, although it is out of reach for the majority of Nigerians. It is sold all over the country, and the best quality varies depending on the producer, raising concerns about the safety of certain of these items. The National Agency for Food, Drug Administration, and Control (NAFDAC) took the ingenuity to free Nigeria of soiled bottled and sachet water by mandating companies to chronicle with it so that it could monitor their activities, (Vanessa et al., 2021).

Given the lack of clean drinking water in Nigeria and the hot weather, sachet and bottled water sales persisted prominent in 2015 (Ugochukwu et al., 2017). In both urban and rural areas of Nigeria, a large percentage of households who do not have admittance to potable water. It is assumed that at least half of the populace purchases water on a day-to-day basis (Richard et al., 2021).

Article Information: *Corresponding Author: mustapha.abdulsalam@sun.edu.ng

Received 17/10/2022 Accepted after revision 25/11/2022

Published: Dec 2022 Pp- 507-512

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/15.4.4>

The increased availability of bottled water in the country necessitates an examination into the operations of the producers to evaluate the magnitude to which they adhere to the established standards. The Federal Government of Nigeria has spent a significant amount of money on basic health care in the last decade since many health workers have expressed concern about the increasing symmetry of dangerous water-borne infections like typhoid fever, diarrhea and measles (Moremi 2012). As a result, a small number of people have taken precautions by boiling and filtering their drinking water at home. In some families, however, drinking bottled water is a safer strategy to avoid contact with these infections.

The product was launched in Nigeria to give safe drinking water free of waterborne infections. However, with the growth of bottled water firms, the original purpose of supplying safe drinking water has been compromised. The purpose of this study was to scrutinize the quality of bottled water in light of the increasing demand for secured and drinkable potable water, as well as the economic expansion of small scale entrepreneurs in the bottled water industry. The study's goal is to assess the bacteriological and physicochemical strengths of bottled water, depict and recognize the bacterial isolates found in the bottled water, define the incidence of bacteria isolated in the bottled water, and develop research-based initiatives to ensure the quality of bottled water sold in the Nasarawa area of Kano State, Nigeria.

MATERIAL AND METHODS

Sample Collection: Twenty-six different brands of bottled water had been acquired from shops in Nasarawa and coded A to Z so that it will disguise the identification of the producers' identities. The samples had been gathered and transported to the laboratory in smooth coolers, wherein they had been preserved at ambient temperature until they were needed.

Bacteriological counts of bottled water samples: The number of heterotrophic bacteria was estimated using a standard plate count (SPC). Bacterial cultures were grown on nutrient agar (Nawzat 2021 & Karissa et al., 2021). Using the 3-3-3 multiple tube fermentation regime, the total coliform count was determined. The culture was done in MacConkey broth. The number of significant tubes with acid and gas generation was recorded after incubation, and the most probable number (MPN) of coliform per 100ml of sample taken was calculated using the MPN index table (Karissa et al., 2021). Using the spread plate approach, the faecal coliform (*E. coli*) of the water samples was determined by Eosin methylene blue (EMB) agar. The number of *E. coli* colonies was reckoned and expressed as colony forming units per milliliter (cfu/ml) (Aneja 2014).

Characterization and Identification of bacterial isolates:

The colony morphology, cellular features, and biochemical reactions of the bacteria were used to identify them (Veronica et al., 2020 & Shima et al., 2021). To acquire pure culture of the isolates, all colonies were sub-cultured from the mixed culture. The genuine isolates were then

conveyed to sterilize nutrient agar slants and stowed in the refrigerator as stock cultures (Karissa et al., 2021).

Determination of physicochemical parameters: The pH of the sampled bottled water was evaluated using a pre-calibrated Denver pH meter model 20 (Karissa et al., 2021). The total suspended particles were estimated by weighing a pyrex beaker after it had been dried up in an oven at 105°C for 60 seconds (Nawzat 2021).

Determination of Temperature: All temperature of the bottle water samples had been measured using a simple mercury-in-glass thermometer calibrated in degrees Celsius, as defined by (Siti et al., 2021, (Muhammad et al., 2021) and (Sheshe & Magashi 2014).

Table 1. Showing the Bacteriological Counts of Bottled Water Samples

Samples	FC (cfu/ml)	TC (MPN/100ml)	HBC (cfu/ml)
A	0	0	2
B	0	0	0
C	0	0	0
D	0	0	68
E	0	0	1
F	0	0	1
G	0	0	2
H	0	1000	118
I	0	0	3
J	0	0	10
K	0	0	3
L	0	0	7
M	0	0	0
N	0	0	0
O	0	0	0
P	0	220	7
Q	0	0	1
R	0	0	0
S	0	0	0
T	0	0	0
U	0	3	61
V	0	0	0
W	0	0	0
X	0	0	2
Y	0	0	1
Z	0	0	10

Key:FC= Faecal Coliform, TC = Total Coliform HBC= Heterotrophic Bacterial Count.

Determination of Conductivity: It was done using the conductivity meter. The conductivity meter was connected to a power source. 25ml of the sample was collected in a flat bottom flasks, followed by 25ml of deionized water in

a second flat bottom flasks. The conductivity of deionized water was read as the control, and then the conductivity probe was placed in the flasks containing the water sample, and its conductivity was read from the conductivity meter (Amanda et al., 2022).

Determination of Fluoride: In a dry square sample cell, 10 ml of each water sample were placed, and 2cm³ of SPADNS reagent was added and mixed. After 30 seconds, the absorbance of the samples was calculated with a spectrophotometer (Nor Azlida et al., 2020).

Determination of Total Chloride: The total chloride was determined using an argentometric titration, which entailed titrating 100 ml of distilled water with water was added of 5% potassium chromate as an indicator and 0.1 N AgNO₃ as a standard. At the end, the solution changed the colour from yellow to pale pink (Sule et al., 2016).

Table 2. Showing the Bacterial Isolates' Characterization and Identification

Isolated bacterial	Gram reaction	Cell shape	Cell arrangement	Motility	Oxidase	Catalase	Amylase	Starch hydrolysis	Inulin	Chitinase	Lactose	Urease	Sucrose	Glucose	Raffinose	Mannitol	VP	Nitrite	Protease	Starch	Cellulose	OXF	Probable identity
1	+	R	S	-	+	+	+	+	+	+	-	-	-	A	A	-	+	A	A	F	-	-	<i>Staphylococcus sp.</i>
2	-	R	CH	+	+	+	+	+	+	+	-	-	-	A	-	-	-	+	A	A	F	-	<i>Aeromonas sp.</i>
3	+	Co	CL	-	+	+	+	+	+	+	-	-	-	A	A	-	+	A	A	OX	-	-	<i>Micrococcus sp.</i>
4	-	R	CL	-	+	+	+	+	+	+	-	-	-	A	-	-	+	-	-	-	-	-	<i>Klebsiella sp.</i>
5	-	R	S	-	+	+	+	+	+	+	-	-	-	A	-	-	-	-	-	A	OX	-	<i>Bacillus sp.</i>

Key: + = Reaction; - = Non- reaction; Co= Cocci, CL = Cluster, CH = Chain; R= Rod, S= Single, F = Fermentative.; OX = Oxidative; OXF = Oxidation-Fermentation, A= Acid production; VP = Voges Proskauer

Determination of Total Hardness: In a complexometric titration, total hardness was determined by titrating 100 ml of water with 0.1 N EDTA using Erichrome black-T as an indicator. The water sample was treated with 10 drops of 25% ammonia before the titration began. The purple color changed to a faint blue color at the end (Mica et al., 2021).

RESULTS AND DISCUSSION

Bacterial Counts: The bottled water's total heterotrophic bacterial counts oscillated from 0 to 118 cfu/ml while total coliform counts ranged from 0 to 1000 coliform/100ml. There was no faecal coliform in all the bottled water samples (Table 1). In this investigation, 96.15 % of the bottled water samples had heterotrophic bacterial counts of fewer than 100 cfu/ml. In reality, 88.46% of the bottled water samples had heterotrophic bacterial counts of less than 10 cfu/ml.

Table 3. Showing the Bacterial Isolates' presence in the Bottled Water Samples

Sample	S1	S2	S3	S4	S5
A	+	+	-	+	-
B	-	-	-	-	-
C	-	-	-	-	-
D	-	-	-	+	-
E	+	-	-	+	-
F	-	+	-	-	-
G	-	+	-	+	-
H	-	+	-	+	-
I	+	+	+	+	-
J	-	+	+	-	+
K	+	-	+	-	-
L	-	+	-	+	-
M	+	-	+	-	+
N	+	+	+	+	-
O	-	-	-	-	-
P	-	-	-	-	-
Q	-	-	+	-	-
R	+	-	-	+	+
S	+	-	-	+	-
T	-	+	-	-	-
U	+	-	+	-	-
V	-	+	+	-	-
W	-	+	-	+	-
X	+	-	-	-	+
Y	-	+	-	-	+
Z	-	-	+	+	+

Key: + = Isolated; - = Not isolated; S1 = *Staphylococcus sp.*; S2 = *Proteus sp.*; S3 = *Micrococcus sp.*; S4 = *Klebsiella sp.*; S5 = *Aeromonas sp.*

Isolated Bacteria from Bottled Water Samples: The bacteria *Staphylococcus sp.*, *Bacillus sp.*, *Micrococcus sp.*, *Klebsiella sp* and *Aeromonas sp.* were identified and characterized in the bottled water samples (Table 2). Table 3 shows where they were found in the bottled water samples.

Bottled Water Samples' Physicochemical Properties: The Hydrogen ion concentration of the bottled water samples ranged from 6.50 to 8.30, and the total hardness ranged from 1 to 86 mg/l. The quantities of chloride and general suspended solids was ranged from 3.58 and 4.64 mg/l and 1.0×10^{-3} to 3.0×10^{-3} g/100ml, correspondingly. (Table 4).

In terms of public health, the significance of safe and quality drinking water cannot be overstated. Some water-borne diseases such as dysentery, cholera, typhoid etc, have

become far less common since the introduction of bottled water. All of the producers were successful in maintaining a faecal coliform count of zero in their goods. Furthermore, 24 bottled water brands (92.30 %) surpassed the maximum coliform limit for drinkable water of 10 per 100 mL of water WHO (2022). However, for 23 of the 26 varieties of bottled water, zero total coliform count was indorsed for emergency use, accounting for 88.46 percent of the entire coliform count (Bukar et al., 2015).

The bacteria identified in the bottled water might have come from the manufacturing machinery or faulty capping and

post-production contamination in bottled water distribution and sales. Many studies have observed that the many heterotrophic bacteria, total coliform and the number of bacterial species found in this study was substantially lower than that seen in sachet water. In five types of sachet water sold in Maiduguri, Bukar et al. (Sasikaran et al., 2012) found heterotrophic bacterial counts ranging from 15400 to 19600 cfu/ml. These figures are higher than the maximum heterotrophic bacterial count discovered in this study, which was 118 cfu/ml in a bottled water brand. Aerobic bacterial counts varying from 0–800 cfu/ml were observed in 22 brands of bottled water obtainable in the Jaffna Peninsula (Omalu et al., 2012).

Table 4. Showing the Physicochemical Potentials of Bottled Water Samples

Sample	pH	Temperature (°C)	TH(mg/l)	CC(mg/l)	Conductivity (µS/cm)	TSS(g/100ml)	FCo (mg/l)
A	6.70	20.4	33	4.20	63.10	0.003	0
B	6.60	20.3	18	4.64	66.20	0.001	0
C	8.30	19.6	10	3.85	63.47	0.002	0
D	7.50	20.1	14	4.64	73.13	0.001	0
E	6.80	20.6	19	3.91	72.40	0.003	0
F	6.50	20.8	7	4.16	72.40	0.001	0
G	6.70	20.4	15	3.91	80.40	0.001	0
H	7.50	20.3	86	4.64	81.33	0.002	0
I	7.50	20.1	8	4.64	86.41	0.003	0
J	6.70	20.2	1	3.93	63.95	0.003	0
K	7.40	20.4	7	4.19	64.00	0.001	0
L	6.70	20.6	9	3.91	64.30	0.001	0
M	7.50	20.1	4	4.44	74.79	0.002	0
N	7.50	19.9	9	3.91	70.20	0.002	0
O	7.90	21.0	11	4.62	79.60	0.003	0
P	8.00	20.0	15	3.91	84.30	0.001	0
Q	7.90	20.9	2	3.58	86.90	0.002	0
R	6.50	20.2	4	3.75	65.30	0.001	0
S	7.50	19.8	24	4.62	72.40	0.002	0
T	6.80	20.0	2	3.91	72.40	0.003	0
U	6.70	20.1	7	4.42	80.40	0.003	0
V	7.50	20.0	9	3.91	72.40	0.001	0
W	7.90	20.6	4	3.84	63.47	0.001	0
X	6.50	20.1	1	3.94	72.10	0.002	0
Y	7.60	20.3	8	3.91	80.40	0.001	0
Z	6.50	19.8	11	3.93	63.56	0.003	0

Key: TH = Total Hardness, CC = Chloride content, TSS = Total Suspended Solid, FCo = Flouride Content

Klebsiella, *Pseudomonas*, *Proteus* sp., and *Chromobacterium violaceum* reported found from sachet water marketed in Samaru, Nigeria, according to (Ugochukwu et al. 2017). Coliform, *Staphylococcus*, and *Pseudomonas* were found in many kinds of bottled water sold in Ile-Ife, South Western Nigeria, by Igbeneghu and Lamikanra (Bukar et al., 2015). Non-carbonated bottled drinking water sold in Sri Lanka

contained *Klebsiella pneumoniae*, *Enterobacter cloacae*, *Pseudomonas aeruginosa*, and *Pasteurella haemolyticus* (Igbeneghu and Lamikanra 2014). However, the three bacteria genera found in this study, *Staphylococcus*, *Micrococcus* and *Aeromonas*, suggest that bottled water is healthier than sachet water. *Staphylococci* and *Micrococci* are

commensals that can be found on human skin, soil and other surfaces. As a result, they conquered the bacterial species isolated from the bottled water samples used in this study. Both water and soil have been shown to contain *Aeromonas* species (Herath et al., 2012).

In order for bacteria to survive in water, pH is critical. The pH values, temperature and conductivity of the bottled water samples used in this investigation were between 6.5 and 8.5, 20-21°C, 5 µS/cm - 199 µS/cm which is the range allowed in potable water (WHO (2022 & Ojosipe 2007) kinds of bottled water sold in the Jaffna Peninsula have pH values ranging from 4.11 to 7.58 (Omalu et al., 2012).

The pH of all the samples, as well as their temperature, chloride content and turbidity was quite well below their limits, similar to those reported by (Sule et al., 2017). The suspended solids in bottled water were less than 30 mg/l (corresponding to 0.003 g/100ml in home water) (Willey et al., 2008). The pH of water is crucial for bacteria to live. The pH of the bottled water samples utilized in this research ranged from 6.5 to 8.5, which is the range permitted in potable water (Amanda et al., 2022). The pH of 22 types of bottled water offered on the Jaffna Peninsula ranges from 4.11 to 7.58 (Omalu et al., 2012). Total suspended solids in bottled water were less than 30 mg/l (0.003 g/100ml in household water) (Willey et al., 2008).

CONCLUSION

Data of the present study strongly suggest that the government should step up its efforts to monitor the fast growing water bottling industry in order to ensure that the public has access to safe and hygienic water. During the manufacturing process, manufacturers should strictly follow all good manufacturing practices (GMP) standards. Bottled water should also be explicitly labeled with an expiration date, authentic manufacturing date, batch number and bottles ought to be closed appropriately to avoid contagion. If any batch of brand fails to gratify regulatory criteria, regulatory organizations should then adequately notify the public. In such cases, the product should be withdrawn from the public. According to the findings of this study, In terms of bacteriological and physicochemical properties, twenty-four of the twenty-six diversities of bottled water samples assessed, 92% of the samples met the approved drinking water quality standards. As a result, the vast majority of bottled water sold in Nasarawa, Kano State is perfectly safe to drink.

Conflict of Interest statement: The authors have stated that there are no competing interest.

REFERENCES

Amanda, Fermiano da Cruz. , Alessandra, Oliveira de Abreu., Paula, Araujo de Souza., Bianca, Deveza., Caroline, Totolote Medeiros., Verônica, Santos Sousa., Bruna, Peres Sabagh., Maria, Helena., Simões, Villas Bôas. (2022). Adaptation and validation of a method for evaluating the bactericidal activity of ethyl alcohol in gel format 70% (w/w). *Journal of Microbiological Methods*,

193:106402

Amina, Othmani., Sara, Magdouli. P., Senthil, Kumar., Ashish, Kapoor., Padmanaban, Velayudhaperumal, Chellam., Ömür, Gökkuş. (2021). Agricultural waste materials for adsorptive removal of phenols, chromium (VI) and cadmium (II) from wastewater: A review. *Environmental Research*, 204(Pt A):111916.

Aneja, K.R. (2014). *Laboratory Manual of Microbiology and Biotechnology*. 1st ed. Published by MedTech Scientific International PVT. Ltd., New Delhi, India, pp. 120 – 121.

Bukar, A.M., Isa, M.A., Mustapha, A., Kyari M.Z. and Ibrahim, F.K. (2015). Bacteriological Analysis of Sachet Water in Maiduguri Metropolis. *The Journal of Applied Sciences Research*, Vol. 2, No. 1, pp. 20–25.

Emily Ying Yang, Chan., Kimberley, Hor Yee Tong., Caroline, Dubois., Kiara, Mc Donnell., Jean, H. Kim., Kevin, Kei Ching, Hung., Kin On. Kwok. (2021). Narrative Review of Primary Preventive Interventions against Water-Borne Diseases: Scientific Evidence of Health-EDRM in Contexts with Inadequate Safe Drinking Water. *International Journal of Environmental Research and Public Health*, 18(23): 12268.

Herath, A.T., Abayasekara, C.L., Chandrajith R. and Adikaram, N.K. (2012). Temporal Variation of Microbiological and Chemical Quality of Non-Carbonated Bottled Drinking Water Sold in Sri Lanka. *Journal of Food Science*, Vol. 77, No. 3, pp. 160 – 164.

Igbeneghu, O.A. and Lamikanra, A. (2014). The Bacteriological Quality of Different Brands of Bottled Water Available to Consumers in Ile-Ife, South Western, Nigeria. *BioMed Central Research Notes*, Vol. 7, pp. 847 – 859. in the Millennium Development Goals Era. *Public Health Nutrition*, 24(13): 4346-4357 *Journal of Microbiological Methods*, 193:106402

Karissa, Culbreath., Heather, Piwonka., John, Korver., Mir, Noorbakhsh. (2021). Benefits Derived from Full Laboratory Automation in Microbiology: A Tale of Four Laboratories. *Journal of Clinical Microbiology*, 59(3): e01969-20.

Mica, Cabrera., Faizah, Taher. Alendre, Llantada., Quyen, Do., Tyeshia, Sapp., Monika, Sommerhalter. (2021). Effect of Water Hardness on Catechin and Caffeine Content in Green Tea Infusions. *Molecules*. 26(12):3485.

Moremi, I.R. (2012). Challenges of Infectious Diseases in Africa. *Journal of Medical Sciences*, Vol. 6, No. 3, pp. 9-12.

Muhammad, Haya. Khan., Mohammad, Nafees., Nisar, Muhammad., Ubaid, Ullah., Rahib, Hussain. , Muhammad, Bilal. (2021). Assessment of Drinking Water Sources for Water Quality, Human Health Risks, and Pollution Sources: A Case Study of the District Bajaur, Pakistan. *Archives Environmental Contamination and Toxicology*,

80(1): 41-54.

Nawzat, Saadi., Khulud, Alotaibi., Laylan, Hassan., Quinshell, Smith. , Fumiya, Watanabe. , Ashraf, A. Khan., Tansel, Karabacak. (2021). Enhancing the antibacterial efficacy of aluminum foil by nanostructuring its surface using hot water treatment. *Nanotechnology*, 32(32).

Nor Azlida, Mohd Nor., Barbara, L. Chadwick., Damian, J. Farnell., Ivor, G. Chestnutt. (2020). The impact of stopping or reducing the level of fluoride in public water supplies on dental fluorosis: a systematic review. *Rev Environ Health*, 35(4): 419-426.

Ojosipe B.A. (2007). Parameters in Water and Wastewater Analysis, 6th Mandatory Training Workshop with Theme: Water. Institute of Public Analysts of Nigeria (IPAN), Lagos, pp. 29.

Omalu, I.C.J., Mohammed, A.Z., Olamide, P.I., Ayanwale, V.A., Adeniran, I. and Gbise, S. (2012). Bacteriological and Physicochemical Analysis of Sachet Water in North Central Nigeria. *Journal of Pharmaceutical and Biomedical Sciences*, Vol. 21, No. 2, pp. 1– 4.

Peres Sabagh., Maria, Helena., Simões, Villas Bôas. (2022). Adaptation and validation of a method for evaluating the bactericidal activity of ethyl alcohol in gel format 70% (w/w).

Richard, M. Mariita., Sébastien, A. Blumenstein., Christian, M. Beckert., Thomas, Gombas., Rajul, V. Randive. (2021). Disinfection Performance of a Drinking Water Bottle System with a UV Subtype C LED Cap Against Waterborne Pathogens and Heterotrophic Contaminants. *Front Microbiology*, 12: 719578.

Sasikaran, S., Sritharan, K., Balakumar, S. and Arasarahman, V. (2012). Physical, Chemical and Microbial Analysis of Bottled Drinking Water. *Ceylon Medical Journal*, Vol. 57, No. 3, pp. 111-116.

Sheshe, M.U. and Magashi A.M. (2014). Assessment of Physicochemical quality of sachet drinking water produced in selected local government areas of Kano metropolis, Kano state- Nigeria. *Bayero Journal of Pure and Applied Sciences*, 7(2): 31 – 35.

Shima, Aboutalebian., Kazem, Ahmadikia., Hamed, Fakhim., Javaher, Chabavizadeh. , Ahmadreza, Okhovat.

, Mahnaz, Nikaeen., Hossein, Mirhendi.(2021). Direct Detection and Identification of the Most Common Bacteria and Fungi Causing Otitis External by a Stepwise Multiplex PCR. *Frontier in Cellular and Infection Microbiology*, 11:644060.

Siti Nor, Fazillah. Abdullah., Azimah, Ismail., Hafizan, Juahir., Fathurrahman, Lananan., Nor Muzlinda, Hashim., Nadiana, Ariffin., Tengku Azman, Tengku Mohd. (2021). Estimation of rainwater harvesting by the reflectance of the purity index of rainfall. *Environmental Science of Pollution and Research International*, 28(27): 35613-35627.

Sule I. O., Agbabiaka T. O., Saliu B. K., Bello A. B., Adeboye A. B. (2017) Bacteriological and physicochemical assessment of selected brands of bottled water in Ilorin, Nigeria. *Al-Hikmah Journal of Pure & Applied Sciences*, 4:15-22.

Sule, I.O., Odebisi-Omokanye, M.B., Gambari-Ambali, R.O. and Okewale, T.A. (2016). “Effects of Disinfectant A” on the Physicochemical and Bacteriological Quality of some Well Water. *Journal of Science, Technology, Mathematics and Education*, Vol. 12, No. 1, pp.1 – 8.

Ugochukwu, S., Giwa, F. I and Giwa, A. (2017). Bacteriological evaluation of sampled sachet water sold in Samaru- Zaria, Kaduna State, Nigeria. *Nigerian Journal of Basic and Clinical Sciences*, Vol. 12, No.1, pp. 6-12.

Vanessa, Pike., Beverly, Bradley., Aviva, I. Rappaport., Stanley, Zlotkin. , Nandita, Perumal. (2021). A scoping review of research on policies to address child undernutrition

Veronica, Veses. , Maria, Del Mar Jovani-Sancho., Raquel, González-Martínez., Isidoro, Cortell-Ballester. , Chirag, C. Sheth. (2020). Raising awareness about microbial antibiotic resistance in undergraduate dental students: a research-based strategy for teaching non-laboratory elements of a microbiology curriculum. *BMC Medical Education*, 20(1): 47.

WHO (2022). Guidelines for Drinking Water Quality, 4th ed., Geneva, Switzerland. <http://www.who.int>.

Willey, J.M., Sherwood, L.M. and Woolverton, C.J. (2008). Prescott, Harley and Kleins, *Microbiology*, 7th ed., McGraw Hill, New York, pp. 471– 589.