

# Bioscience Biotechnology Research Communications

Volume-16 Number (2) Apr-May-Jun 2023

Print ISSN: 0974-6455

Online ISSN: 2321-4007

CODEN: BBRCBA

Website: [www.bbrc.in](http://www.bbrc.in)

An International Peer  
Reviewed Open Access Journal

Published By:

Society for Science & Nature (SSN)

Bhopal India

website: [www.ssnb.org.in](http://www.ssnb.org.in)

Online Content Available: Every 3 Months at

[www.bbrc.in](http://www.bbrc.in)



**SOCIETY**  
FOR SCIENCE AND NATURE

SCIENCE FOR LIFE

RESEARCH ARTICLES

- Predictive Modeling of COVID- 19 Confirmed Cases Using Regressive Objective Regression Methodology 68-72  
**Fernando Martínez Fernández, Ricardo Osés Rodríguez, Rigoberto Fimia Duarte, Claudia Osés Llanes, Melba Zayas González, Maribel I Noda Valledor and Dania Artiles Román**
- Women with Postpartum Depression in Kerala: A Survey 73-76  
Objective Regression Methodology  
**Pooja Prasad and Balakrishnan Kalamullathil**
- BMI-For-Age Percentile as Criteria for Assessment of 77-83  
Obesity in Children and Adolescents  
**Manju Dewan**
- Adsorption of Phenol And Resorcinol on Parthenium Based Activated 84-90  
Carbon (PAC) in Basal Salt Medium:  
**Khushi Gandhi, Shambhatee Srivastav, Huda Afreen and Ravi Kant Singh**
- Avifaunal Abundance of Luming Forest Reserve Area, Assam, India 91-99  
**Prof. Jashodeb Arjun and Rajesh Paul**
- Standardization and Evaluation of Buffers: A One Step DNA Extraction 100-103  
Protocol from Microbial Cultures  
**Poornima Shyam, Dr. Shyama Subramaniam, Dr. Shyam Subramaniam and Edwina Jospiene M**
- Didactic Strategy for the Use of Virtual Itineraries in the Training of 104-114  
Graduates in Education Geography  
**Oliday Aguilar Espinosa, Erich Rodríguez Vallejo, Rafael Armiñana García and Rigoberto Fimia Duarte**
- Localization of neuropeptide Y(NPY) Immunoreactivity in the Proximal and 115-119  
Distal Intestinal Region of Teleost Fish, Notopterus Notopterus  
**Tulika A Khadse, Yashashree A Gadhikar, Trupti S Khedkar and Sudhir G Chird**
- Ecology of Biofouling Phytoplankton in Chinnamuttom Harbour Waters Southeast Coast of India 120-125  
**P Nithya, B Dhanalakshmi and P Santhanam**
- Knowledge and Management Assessment of Temporomandibular Joint (TMJ) Disorders 126-131  
Among Dental interns at King Saud University  
**Hamad Albagieh, Abdulrahman Alghamdi, Musaad Almeyfayz, Abdulrahman Alshehri and Osama Alsaiady**

# Bioscience Biotechnology Research Communications

Open Access International Journal

## Editor-In-Chief

**Prof Sharique A Ali**

PhD FLS FRBS (UK)

**180 FULL RESEARCH PAPERS**

Ex- Cooperating Scientist USDA (PL-480)

Virginia State University Virginia USA

Professor of Physiology & Head Department of Biotechnology,

Saifia College, Barkatullah University Bhopal India

Publons Researcher ID [ADN-6124-2022](https://publons.com/author/1234567890/ADN-6124-2022)

Website: <http://www.drshariqali.com>

Scopus ID: 7403093928 Clarivate ID: E-2495-2019 [Google Scholar](https://scholar.google.com/citations?user=7403093928)

Orcid Id: <https://orcid.org/0000/0002/0378/7385>

---

## ASSOCIATE EDITORS

---

### **Dr S Mohanty PhD (IIT)**

Biomedical Sciences  
Senior Director Senseonics  
University of Minnesota  
Gaithersburg Maryland USA  
[Short Bio](#)

### **Prof A Denizli PhD**

Cancer Biologist Ankara,  
Turkey [Short Bio](#)

### **Dr Laxmi Thangavelu PhD (Toronto)**

Pharmacology & Biomedicine  
Saveetha Medical University  
Chennai, India  
[Short Bio](#)

### **Dr Lisetskiy Fedor PhD**

Environment and Resource Management  
Director Federal-Regional Centre Natural Resources  
Belgorod State University Belgorod  
Russia [Short Bio](#)

### **Dr M P Gashti PhD**

Biomaterial Technology  
British Columbia V1X 7Y5,  
Canada [Short Bio](#)

### **Dr I El Emary PhD**

Information Technology  
Professor & Dean of Computer Science  
King Abdulaziz University Jeddah,  
Saudi Arabia [Short Bio](#)

### **Dr A. Bath MD**

Immunology  
Kansas University Medical Center,  
Kansas USA [Short Bio](#)

### **Dr S. Salim PhD**

Clinical Medicine  
Research Scientist Clinical Diagnostics  
LGC Group Gaithersburg MD 20878 USA

ORCID Id: <https://orcid.org/0000-0001-6642-3450> [Short Bio](#)

### **Prof D K Belsare PhD DSc FNASc**

Biosciences  
Baylor College of Medicine  
Houston USA. & Barkatullah  
University Bhopal, India [Short Bio](#)

---

## ACADEMIC EDITORS

---

### **Dr M Bakr PhD MDS**

Dental Sciences  
Griffith University Gold Coast Campus:  
Southport, QLD, Australia  
[Short Bio](#)

### **Dr Qinglu Wang PhD**

Human Genetics & Biomedical Technology  
Department of Basic Medical Education  
University, Zibo, 255213, China

Orcid Id: <https://orcid.org/0000-0002-2891-9399> [Short Bio](#)

**Dr K Jasim PhD** Environmental  
Toxicology The University of Alabama  
at Birmingham (UAB) Alabama 35233  
USA [Short Bio](#)

**Dr P Muthuirulan PhD**  
Human Pathology  
Harvard University Cambridge  
MA USA [Short Bio](#)

**Dr Bashar Saad PhD**  
Biochemistry / Cell Biology  
Full Professor Arab American University  
Palestine [Short Bio](#)

**Dr FA Kabbinwar MD FACP**  
Ex-Professor of Oncology at UCLA  
San Diego California USA [Short Bio](#)

**Dr G Goyal PhD**  
DM Cardiology  
Director Cardiology  
QRG Super Specialty, Hospital  
Delhi NCR India [Short Bio](#)

**Dr M Maxime PhD**  
Physiology, Molecular and Cellular Biology  
American University of Raïs Al-Khaimah,  
United Arab Emirates  
[Short Bio](#)

**Dr W Thong-Asa PhD (Medical Physiology)**  
Department of Zoology Kasetsart University  
Bangkok, Thailand  
<https://www.researchgate.net/profile/Wachiryah-Thong-Asa>

**Dr Halison C. Golias PhD**  
Microbiology and Biotechnology  
Federal Technological  
University of Paraná Brazil [Short Bio](#)

**Dr SM Singh PhD**  
Tumor Immunology  
Professor of Animal Biotechnology  
School of Biotechnology  
Banaras Hindu University Varansi India  
[Short Bio](#)

**Dr Shaima Miraj PhD**  
Health Sciences  
Saudi Electronic University  
Riyadh Saudi Arabia [Short Bio](#)

---

#### INTERNATIONAL EDITORIAL AND ADVISORY BOARD

---

**Dr Kazutoshi Okuno PhD**  
Former Professor Plant Genetics  
and Breeding Science, University of Tsukuba  
Japan [Short Bio](#)

**Dr Alex Eberle PhD**  
Pathobiology  
Emeritus Professor Molecular Biology  
University of Basel Switzerland  
Oberer Batterieweg 71 CH-4059 Basel Switzerland [Short Bio](#)

**Dr Saurav Das PhD**  
Agriculture Sciences  
Horticulture University of Nebraska  
Lincoln, USA [Short Bio](#)

**Dr Ng Z Xiang PhD**  
Molecular Biochemistry  
School of Biosciences  
University of Nottingham  
Malaysia [Short Bio](#)

**Dr R Fimia Duarte PhD**  
Biomedical Sciences  
Department of Biology  
Central University Marta Abreu of Las Villas.  
Villa Clara Cuba Island  
[Short Bio](#)

**Dr AM Castrucci PhD**  
Cell Physiology  
Professor of Physiology & Biochemistry  
Sao Paulo University Brazil [Short Bio](#)

**Prof Monica Butnariu PhD**  
Nutritional Biochemistry  
Banat's University of Agricultural Sciences  
Timisoara, Romania  
Scopus Id: 15070536800 [Short Bio](#)

**Dr SK Pal PhD**  
Professor of Genetics  
Skyline University, Kano, Nigeria  
[Short Bio](#)

**D Kumar PT PhD**  
Biomedical Sciences  
Boston University College of Health  
& Rehabilitation Sciences: Sargent College  
Director, Movement and Applied Imaging Lab  
Boston MA USA [Short Bio](#)

**Prof S Shenuka PhD**  
Health Sciences / Dentistry  
University of Kwazulu  
Natal South Africa  
[Short Bio](#)

**Dr Absar Ahmad PhD**  
Chemical Sciences  
National Chemical Laboratory, CSIR  
Pune 411008, India [Short Bio](#)

**Dr P. Rattanachaikunsopon PhD**  
Biomedical Sciences  
Department of Biological Sciences, Faculty of Science,  
Ubon Ratchathani University, Warin Chamrap,  
Ubon Ratchathani 34190, Thailand [Biography](#)

**Dr M. Miglani MS (Ortho) AIIMS**  
Director (Orthopedics)  
Fortis Multispecialty Hospital New  
Delhi India [Short Bio](#)

**Dr Dilian G Georgiev PhD**  
Department of Ecology  
University of Plovdiv, Plovdiv  
Bulgaria Orcid Id: <https://orcid.org/0000-0003-2885-4895>

**Prof SKM Habeeb PhD**  
Applied Bioinformatics  
School of Bioengineering,  
SRM Institute of Science & Technology  
Kattankulathur 603203 Tamil Nadu India  
[Biography](#)

---

#### EDITORIAL TEAM

---

##### EXECUTIVE EDITOR

**Dr Ayesha PhD FSSN**  
Professor of Biochemistry Saifia College,  
Barkatullah University Bhopal India  
Orcid Id: <https://orcid.org/0000-0002-7924-3106>

##### MANAGING EDITOR

**Dr Mohd Miraj PhD**  
Director Health Sciences  
AIHMS Gautam Nagar  
New Delhi India [Short Bio](#)

---

#### STATISTICAL EDITORS

---

##### Dr Shahnawaz Anwer PhD

Polytechnic University  
Hongkong [Short Bio](#)

##### Dr Vinars Dawane PhD

Environmental Biotechnology,  
Dhar India [Short Bio](#)

---

#### HONORARY TECHNICAL CONSULTANTS

---

##### Dr LK Jakkala PhD

Clinical Medicine  
2nd floor, Quadrant 4 Cyber Towers  
HITECH City Hyderabad Telangana 500081, India

##### Er Faraz Ali BE

MBA IIM (Indore), First floor C52 HB Colony  
Koifiza Second  
Bhopal 462001 India

---

#### EDITORIAL TEAM MEMBERS

---

##### Dr J Peter PhD (Cell Biology)

Principal and Professor of Zoology  
Shashib College Bhopal 462036 India

##### Dr Arjun Deb PhD

Professor of Zoology & Biochemistry  
Lumding College Assam 782447 India

##### Dr R Ahamed MD

Community Medicine  
College of Medicine  
VC78+QMQ, Industrial Area,  
Al Majma'ah 15341, Saudi Arabia

##### Dr Naima Parveen PhD (Bioinformatics)

Department of Biotechnology  
MANF UGC Fellow Saifia College  
of Science Bhopal 462001 India

##### Dr Sushma Prasad PhD (Animal Sciences)

Zarifa Farm, Kachhwa Road,  
Karnal, Haryana 132001, India

##### Dr Ishrat Naaz PhD (Structure Biology)

Department of Biotechnology  
MANF UGC Fellow Saifia College  
of Science Bhopal 462001 India

##### Dr Kamal Zaidi PhD (Enzymology)

Department of Microbiology  
Peoples University  
Peoples Campus, Bhanpur,  
Bhopal, 462037 India

##### Dr Anjali Choudhary PhD (Toxicology)

Department of Biochemistry  
Opposite to Dussehra Maidan,  
BHEL Square, Sector A, Govindpura,  
Bhopal, 462023 India

##### Dr Raj Sharma PhD (Pharmacology)

Pharmaceutical Sciences  
Chhattisgarh Institute of Medical  
Sciences (CIMS), Bilaspur, CG, India

##### Dr Neelu Qayyumi PhD (Bioscience)

Professor and Head Life Sciences  
Mittal College Opposite to Bhopal Memorial  
Hospital Research Centre (BMHRC), Navi Bagh,  
Karond, Bhopal, 462008 India

## Editors Communique

Have we tamed the coronavirus? May be yes,  
as pandemics do not die, they can only be faded !

Science and technology has made it possible, in the shortest span of time, it has shown that with firm determination and international cooperation, we can win over the onslaughts of even the worst of the pandemics. COVID-19 is perhaps fading over now, due to our coordinated efforts worldwide. Though we have lost millions, in the two year period, partly due to the mishandling of the viral attacks and somewhat by our own follies and carelessness. Anyway lessons learnt from the past, always make us more stronger and determined. Let us now not relax and work on a better mode, as all is still not well yet. The almost taming of the virus and its cousins have indicated some of the concealed failures, on which we have to focus now. We have to be more vigilant, and even a bit of laxity can spoil the good work done. On societal and governmental parts, utmost care and caution is required on a long term basis.

On behalf of Bioscience Biotechnology Research Communications, we falter at words to express our deep sense of solitude and grief on the catastrophic events of the world wide pandemic, spanning over two years now. We pray for the strength to bear this universal calamity and come up with long lasting fortitude to eradicate it soon.

*Biosc Biotech Res Comm* is an open-access international platform for publication of original research articles, exciting meta-reviews, case histories, novel perspectives and opinions in applied areas of biomedical sciences. It aims to promote global scientific research and development, via interactive and productive communications in these areas, helping scholars to present their cherished fruits of research grown on toiled and tilled trees of hard work in life sciences. Being the publication of a non-profit academic Society for Science and Nature, Bhopal India, since 2008, *Biosc Biotech Res Comm* strongly believes in maintaining high standards of ethical and quality publication.

Quality publication is one of the ways to keep science alive, and good journals have a leading role to play in shaping science for humanity! As teachers, we have great responsibilities, we have to advocate our students to accomplish and show them the path to test their mettle in hard times to excel, especially in the post COVID 19 era. Science and its advocates will rise more to the occasion and will soon provide succor to the already grief stricken humanity.

Sharique A. Ali, PhD  
Editor-in-Chief

Bioscience Biotechnology  
Research Communications  
Journals Website: <http://www.bbrc.in>  
Editors Website: <http://www.drshariqali.com>



# Bioscience Biotechnology Research Communications

Open Access International Journal

## About Us

*Biosc Biotech Res Comm* is an official publication of an academic, non-profit Society for Science and Nature, Bhopal India, since 2008. It is a peer reviewed journal that publishes original research articles pertaining to exciting applied areas of biology including biomedical sciences. The aim of *Biosc Biotech Res Comm* is to promote quality scientific research in fundamental and applied areas of biological and biomedical sciences and their allied branches.

It publishes scholarly articles demonstrating academic quality and scientific rigor for disseminating current scientific information on research and development for human welfare. *Biosc Biotech Res Comm* audiences its large number of readers from diverse regions of Asia, Africa, Europe and other developing nations across the world. It is an open access means of scientific communication, that provides contribution to the existing knowledge as well as envisages scientific temper specially in the young minds, pursuing science as a career.

Articles aimed for publication in *Biosc Biotech Res Comm* must have new experimental methods of biotechnological significance, novel laboratory obtained results, interesting interpretation of data pertaining to providing practical solutions to human-welfare problems of substantial scientific significance. The publishers of *Biosc Biotech Res Comm* believe in best of publication standards, hence a single journal is being published by them since 2008, to focus on its high academic standards, selecting quality papers for a timely schedule of publication. *Biosc Biotech Res Comm* strives hard to maintain high quality and follows best practices of publication, particularly in prioritizing originality and quality, hence it has a tough rate of article selection. Less than 50 percent of submitted manuscripts are accepted, and reluctantly, a large number of articles are returned by us.

Articles are selected for possible publication, keeping in view the novelty of the work, originality (plagiarism / similarity levels are checked), word count, explicit English language using quality writing, lucid presentation and interpretation of data, along with conclusive data based statements showing contribution to the existing knowledge. Before final acceptance each article undergoes several rounds of unbiased anonymized revisions, strictly complying the reviewers comments and their satisfaction.

*Biosc Biotech Res Comm* categorizes articles into exciting analytical systematic data based reviews, novel case reports, original research articles, rapid communications and letters to the editor, including lively correspondence and perspectives. Each type of article has a special format and should strictly comply with the up-dated instructions for authors, which are published in all issues of *Biosc Biotech Res Comm* as well as are on the official website of the journal.

## Aims and Scope

*Biosc Biotech Res Comm* is an open access means of scientific communication that provides contribution to the existing knowledge as well as envisages scientific temper in the young minds, pursuing science as a career. It publishes scholarly articles following scientific rigor for disseminating current information on research and development in applied biology and biomedical sciences. Articles may include new experimental methods of bio-medical significance, new laboratory obtained results, novel interpretation of existing data pertaining to providing practical solutions to human welfare problems of substantial scientific significance.

*Biosc Biotech Res Comm* has a special task of helping researchers from developing countries to present their cherished fruits of research grown on toiled and tilled trees of hard work. Such scholars are encouraged with significant waivers in publication charges. All articles under submission to *Biosc Biotech Res Comm* should aim for the development of technological concepts, rather than merely recording the facts, showing evidence of scholarly publication.

Articles submitted to *Biosc Biotech Res Comm* are evaluated according to their intellectual merit, without regard to the race, gender, sexual orientation, religious beliefs, ethnic origin, citizenship, political philosophy, or institutional affiliation of the author (s). Editorial decisions on manuscripts submitted to our journal are based on independent, anonymized peer review reports. The journal is committed to an editorial process that is not compromised by financial or political influence, thereby actively seeking and encouraging submissions from underrepresented segments of the global scholarly communication ecosystem.

Incomplete studies and manuscripts not in strict compliance with the journals policies will be strongly discouraged and rejected. Each type of article has a special format and should comply with the updated *Biosc Biotech Res Comm* [Instructions for authors / submission check List](#), published in its issues. All articles in *Biosc Biotech Res Comm* are published under a [Creative Commons License, International Attribution 4.0 BY-CC](#), meaning thereby a free unlimited use of the articles for academic purposes without any embargo. We are particular in demonstrating conformance with established industry guidelines and best practices promoted by professional scholarly publishing organizations such as: [Committee on Publication Ethics \(COPE\)](#) and [Principles of Transparency and Best Practice in Scholarly Publishing](#).

Biosc Biotech Res Comm strives hard to promote quality scientific research in fundamental and applied areas of biotechnology, bioscience and biomedical sciences via interactive communication among biologists, biotechnologists, health science personnel and biomedical experts from Asia and other regions of the world. It audiences its large number of authors from diverse regions such as Europe, Asia, South East Asia, Russian Federation, the Asia Pacific including several developing nations, because of its quality and timely schedule of publication. The journal is read by a large community of scholars, scientists and students from many continents

## **Journal Polices of Bioscience Biotechnology Research Communications**

(Author Ethical Statement / Copyright forms / Plagiarism Check Report)

### **Authors**

Authors are specifically those who have made:

Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; and / or drafting the work or revising it critically for important intellectual content; and / or final approval of the version to be published.

The corresponding author's specific responsibilities include:

- Manuscript correction and proof reading. Handling the revisions and re-submission of revised manuscripts up to the acceptance of the manuscripts. Agreeing to and signing the Author Publishing Copyright / Ethical Statement/ Plagiarism Level Check Certificate Forms on behalf of relevant co-authors.
- Arranging for payment of an APC (article processing charge) where one is required. The affiliation of the corresponding author is used to determine eligibility for discounted or waived APCs under our journals Waiver Policies.
- Acting on behalf of all co-authors in responding to queries from all sources postpublication, including questions relating to publishing ethics, reuse of content, or the availability of data, materials, resources etc.
- Acknowledgments section in their publication with permission, for example to recognize the contributions of anyone who provided research or writing assistance.
- We integrate with established and emerging industry standards to increase transparency in authorship (for example, ORCID).

**Author Affiliations:** Any article affiliations should represent the institution(s) at which the research presented was conducted and/or supported and/or approved. For non-research content, any affiliations should represent the institution(s) with which each author is currently affiliated.

**Acknowledgements of funds / grants** etc received for the submitted work must be mentioned before the section of references: This work was supported by ——— Name of Agency, department / Grant number ——— Year to ——— ( Name of Author (s)).

Where no specific funding has been provided for the research, we ask that corresponding authors use the following sentence: The author(s) received no financial support for the research, authorship, and / or publication of this article.

### **Plagiarism**

Plagiarism is defined as using some ones else's ideas, words, data, or other material produced by them without acknowledgement. It is the unauthorized use or close imitation of the language and thoughts of another author and representing them as one's own original work and *Biosc Biotech Res Comm* condemns all forms of plagiarism, following a very strict and vigilant policy of removing this malady. Within the academia, it is considered dishonesty or fraud and offenders are subject to academic censure.

Plagiarism can be unintentional or intentional, reproducing academic material without appropriate credit to the original authors (Citations / References). Similarly self-plagiarism is the re-use of significant, identical or near identical portions of one's own work without citing the original work. This is also known as recycling fraud. Worst form of plagiarism is to steal the whole article from some journal and publish it under one's own name in another journal.

Plagiarism, fabrication, unethical or redundant publication grossly violates the editorial policies of *Biosc Biotech Res Comm*, which follows best practice guidelines given by the [International Committee of Medical Journal Editors \(ICMJE\)](#) and [Committee on Publication Ethics \(COPE\)](#), as mentioned in the Instructions for Authors *Biosc Biotech Res Comm*.

**All authors submitting their MS to *Biosc Biotech Res Comm* must complete and sign the ethical statement form and append the Plagiarism Check Certificate of their MS along with copy-right form ([www.bbrc.in](http://www.bbrc.in)) failing which, their MS will not be processed further.**

The Editorial Committee of *Biosc Biotech Res Comm* will blacklist any author found to be guilty of plagiarism or exceeding the standard limits of similarity levels of text matter in their MS. The name of author(s) committing plagiarism or using similar text without appropriate citations will also be disseminated to concerned authorities.

We do not tolerate plagiarism in any of our publications, and we reserve the right to check all submissions through appropriate plagiarism checking tools. Submissions containing suspected plagiarism, in whole or part, will be rejected. If plagiarism is discovered post publication, we will follow our guidance outlined in the Retractions, Corrections and Expressions of Concern section of these guidelines. We expect our readers, reviewers and editors to raise any suspicions of plagiarism, either by contacting the relevant editor or by emailing at [editor@bbrc.in](mailto:editor@bbrc.in).



## Complaint Policy of Biosc.Biotech.Res.Comm

**Genuine complaints in Publication:** Complaint or expression of dissatisfaction made in honest intention of improvisation are always welcome, as they provide an opportunity and instant moment of attaining quality. The editorial team of Bioscience Biotechnology Research Communications shall strive hard to establish, along with the publisher, a transparent mechanism for appeal against editorial decisions or any related matter of publication. If still there are any genuine complaints related to ethical publishing, we are always open to them for the sake of maintaining quality and ethics of publication.

Please write your complaint with Journal title, Vol No/ Issue No /Year /Page numbers, full title of the MS and necessary author details along with type of complaint. The complaint must be about something that is within the jurisdiction of Bioscience Biotechnology Research Communications, its contents or process such as authorship, plagiarism, copy right violation, multiple, duplicate, or concurrent publications/simultaneous submissions etc. Similarly, undisclosed conflicts of interest, reviewer bias or competitive harmful acts by reviewers or any bias of apparent discontentment, backed by logic and judicial discretion will be immediately looked into without any bias and discrimination.

If the Editor receives a complaint that any contribution to the Journal breaks intellectual property rights or contains material inaccuracies or otherwise unlawful materials, a detailed investigation may be requested into, with the parties involved, substantiating their materialistic claims in writing, following the law of natural justice. We assure that we will make a good faith determination to remove the allegedly wrongful material or take actions as per law. All the investigations and decisions are to be documented to the Journal.

Our aim is to ensure that Bioscience Biotechnology Research Communications follows best practices in publication and is of the highest quality, free from errors. However, we accept that occasionally mistakes might happen, which are inadvertently made or beyond human control, giving opportunity to all parties to decide the best to rectify.

**Editorial Complaints Policy:** The Managing Editor and staff of Bioscience Biotechnology Research Communications will make every efforts to put matters right as soon as possible in the most appropriate way, offering right of reply where necessary. As far as possible, we will investigate complaints in a blame-free manner, looking to see how systems can be improved to prevent mistakes occurring.

**How to Make a Complaint:** Complaints about editorial content should be made as soon as possible after publication, preferably in writing by email to [editor@bbrc.in](mailto:editor@bbrc.in) or by on-line submission at [www.bbrc.in](http://www.bbrc.in)

## Peer Review Policy

Unbiased, independent, critical assessment is an intrinsic part of all scholarly work, including the scientific process. Peer review is the critical assessment of manuscripts submitted to journals by experts who are not part of the editorial staff, and is, therefore, an important extension of the scientific process. Each article submitted to Biosc. Biotech. Res. Comm for publication is reviewed by at least two specialist reviewers of the concerned area. The dual review process is strictly followed and in certain controversial cases, the opinion of a 3rd reviewer can also be sought.

## Manuscript Processing

Upon on-line submission of the manuscript, the author will be acknowledged with a MS number, via e-mail. Initially an article will be reviewed by the Editorial team to judge the academic quality, scientific rigor and format of the manuscript, in particular its relevance to the scope of the journal, compliance with instructions to authors check list and levels of similarity / accidental plagiarism.

Article submissions must consist of academic material that is unique and original, meaning that articles must engage cited material through critical thought. Articles must follow conventions of the English language in regard to proper grammar, punctuation, and typical writing practices. All factual statements must be supported by cited sources or research evidence. Authors must ensure the accuracy of citations, quotations, diagrams, tables, and maps.

Articles written in poor English language with confusing or illogical statements, or not conforming to instructions to authors of Biosc.Biotech.Res. Comm will either be rejected or returned to the authors for reformatting. Manuscripts deemed proper only will be forwarded to at least two subject experts to work as anonymized reviewers in a time bound frame of 4 to 5 weeks, to provide their unbiased input on the overall quality of the reviewed manuscript as per standard international norms.

Acceptable manuscripts will be checked for data analysis and verification of references before the author is notified about the status of the paper with any suggestions for modifications strictly as reviewers comments and revisions asked. Editors will check at every step for full compliance and revision of all such articles in press. Finally accepted articles will then be forwarded for typesetting and formatting, and the galley proof will be sent to the authors for proof reading, before final publication in a time bound period. For detailed process of manuscript, please see the flow chart of MS processing in Biosc.Biotech.Res.Comm.

## Guidelines for Reviewers

An unpublished manuscript is a privileged document. Please protect it from any form of exploitation. Don't cite a manuscript or refer to the work it describes before it has been published and don't use the information that it contains for the advancement of your own research or in discussions with colleagues. Adopt a positive, impartial attitude toward the manuscript under review, with the aim of promoting effective and constructive scientific communication.

If you believe that you cannot judge a given article impartially, please return it immediately to the editor. Reviews must be completed within 4 to 5 weeks. If you know that you cannot finish the review within that time, immediately return the manuscript to the editor.

In your review, consider the following aspects of the manuscript: –Adherence to style of the MS as set forth in Instructions to Authors of *Biosc Biotech Res Comm*.

- Adequacy of title, abstract and its contents. Explicit language and clear expression of findings in the manuscript.
- Significance of objectives, questions or subjects studied, with a clear justification or rationale.
- Originality of work: It should be checked through standard plagiarism software only.
- Appropriateness of approach or methodology and adequacy of experimental techniques with citations, so that the work can be easily replicated.
- Appropriateness of clear images, figures and or tables and length of article, word count etc..
- Experimental data its lucid presentation and critical interpretation.
- Soundness of conclusion based on data, and interpretation and relevance of discussion of the manuscript.
- Appropriate literature citations as per Harvard Style of References with updated references.
- All sources must be cited in the reference list and in the main text. References with non-English titles must include a translation. All in-text citations must be cited in the reference list and all sources in the reference list must be cited within the article. Sources accessed online must include a DOI or URL.

If you wish to mark the text of the manuscript, use a pencil or make a photocopy, mark it, and return it together with the original. You can be particularly helpful in pointing out unnecessary illustrations and data that are presented in both tabular (and graphic) form and in detail in the text. Such redundancies are a waste of space and readers time.

A significant number of authors have not learnt how to organize data and will benefit from your guidance. Do not discuss the paper with its authors. In your comments intended for transmission to the author, do not make any specific statement about the acceptability of a paper. Suggested revision should be stated as such and not expressed as conditions of acceptance. Present criticism dispassionately and avoid offensive remarks.

Organize your review so that an introductory paragraph summarizes the major findings of the article, gives your overall impression of the paper and highlights the major shortcomings. This paragraph should be followed by specific numbered comments which if appropriate may be subdivided into major and minor points. Confidential remarks directed to the editor should be typed (or handwritten) on a separate sheet, not on the review form. You might want to distinguish between revisions considered essential and those judged merely desirable.

Your criticisms, arguments and suggestions concerning the paper will be most useful to the editor and to the author if they are carefully documented. Do not make dogmatic, dismissive statements, particularly about the novelty of work. Substantiate your statements.

Reviewer's recommendations are gratefully received by the editor. However, since editorial decisions are usually based on evaluations derived from several sources, reviewers should not expect the editor to honor every recommendation.

## Conflict of Interest

Conflict of interest exists when as author (or the author's institution), reviewer, or editor has financial or personal relationships that inappropriately influence (bias) his or her actions (such relationship are also known as dual commitments, competing interests, or competing loyalties). However, conflicts can also occur for other reasons, such as personal relationships, academic competition, and intellectual passion. Increasingly, individual studies receive funding from commercial firms, private foundations, and the government. The conditions of this funding have the potential to bias and otherwise discredit the research.

When authors submit a manuscript, they are required to disclose all financial and personal relationships that might bias their work. To prevent ambiguity, authors must state explicitly whether potential conflicts do or do not exist. It is the discretion of editorial committee of *Biosc BiotechRes. Comm* to resolve any conflict of interest between the author(s) and reviewers. Editors may choose not to consider an article for publication if they feel that the research is biased by the sponsors funding the research project.

## Duplicate and Redundant Publication

Duplicate or redundant publication, or self-plagiarism, occurs when a work, or substantial parts of a work, is published more than once by the author (s) of the work without appropriate cross-referencing or justification for the overlap.

We expect our readers, reviewers and editors to raise any suspicions of duplicate or redundant publication, either by contacting the relevant editor or by emailing at [editor@bbrc.in](mailto:editor@bbrc.in). When authors submit manuscripts to our journals, these manuscripts should not be under consideration, accepted for publication or in press within a different journal, book or similar entity, unless a journal is explicit that it does not have an exclusive submission policy.

## Retractions

Editors will consider retractions, corrections or expressions of concern in line with COPE's Retraction Guidelines. If an author is found to have made an error, the journal will issue a corrigendum. If the journal is found to have made an error, they will issue an erratum. Retractions are usually reserved for articles that are so seriously flawed that their findings or conclusions should not be relied upon, or that contain substantial plagiarism or life-endangering content. Journals that publish Accepted Manuscripts may make minor changes such as those which would likely occur during copyediting, typesetting or proofreading, but any substantive corrections will be carried out in line with [COPE's Retraction Guidelines](#).

## Ethical Issues

### 1. Animal and Human Studies

Ethical declarations in research form an integral part during the submission process of a manuscript to a journal. Bioscience Biotechnology Research Communications requires that the experimental conditions under which animal and human assays and tests are performed are as per standard protocols used worldwide. Authors must make it clear in writing that the procedures they used were as humane as possible and have been compiled with the guidelines for animal care of their institutions or with national / international guidelines. Studies on animals must comply with the prevailing standards of animal welfare according to Indian Council of Medical Research Guidelines or Central Committee of Animal Ethics in India and likewise following similar conditions elsewhere, (Ethical Approval Committees/ Institutional Review Board with Approval Number is necessary). For details of animal studies please see : ARRIVE and Guide for the Care and Use of Laboratory Animals

Studies involving human subjects / patients / and also if the manuscript includes case reports / case series, authors need to provide the following: Name of the Ethical Committees /Institutional review Board, they have obtained consent from along with approval number /ID. Authors should specifically mention that the study was in accordance with the Helsinki Declaration of 1975 ( Human research: Helsinki Declaration as revised in 2013).

### Human Studies: Ethical Standards and Informed Consent

++For studies involving human subjects and volunteers, please indicate in the manuscript, in a section preceding the References, the following statement or an analogous statement that applies to your situation: "All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975 Human research: Helsinki Declaration as revised in 2013.

Informed consent should be obtained from all patients for being included in the study." If any identifying information about participants is included in the article, the following sentence should also be included: "Additional informed consent was obtained from all individuals for whom identifying information is included in this article." If you have not included or cannot include this statement in your manuscript, please provide the reason or an alternative statement here and in the manuscript.

### 2. Disclosure of Interest

Authors must provide details of any financial or personal relationships that might bias the work being submitted.

In a section of text preceding the References, please provide relevant information for each author(s) with regard to any conflicts of interest. All submissions must include disclosure of all relationships that could be viewed as presenting a potential conflict of interest.

### 3. Acknowledgement of sources:

Proper acknowledgment of the work of others must always be given. Funding acknowledgement must be properly made with grant details, number etc.

**Data access and retention:** Authors may be asked to provide the raw data in connection with a paper for editorial review, and should be prepared to provide public access to such data.

## Open Access Policy Statement

Bioscience Biotechnology Research Communications is an open access journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author for any used content, however all freely used content must be properly cited with details. This is in accordance with the BOAI definition of open access. The full text of all content of Bioscience Biotechnology Research Communications is available for free and has open access without delay with no embargo period.

All research articles published in our journal are fully open access: immediately freely available to read, download and share. Articles are published under the terms of a Creative Commons license which permits use, distribution and reproduction in any medium, provided the original work is properly cited. The author(s) and copyright holder(s) grant(s) to all users a free, irrevocable, worldwide, perpetual right of access to, and a license to copy, use, distribute, transmit and display the work publicly and to make and distribute derivative works, in any digital medium for any responsible purpose, subject to proper attribution of authorship, as well as the right to make small numbers of printed copies for their personal use.

A complete version of the work and all supplemental materials, including a copy of the permission as stated above, in a suitable standard electronic format is deposited immediately upon initial publication in at least one online repository that is supported by an academic institution, scholarly society, government agency, or other well-established organization that seeks to enable open access, unrestricted distribution, interoperability, and long-term archiving.

Open access is a property of individual works, not necessarily journals or publishers. Community standards, rather than copyright law, will continue to provide the mechanism for enforcement of proper attribution and responsible use of the published work, as they do now.

### **Retractions/ Corrections / Withdrawal**

Submission of an article to Biosc. Biotech. Res.Comm implies that the work has NOT been published or submitted elsewhere, therefore, the journal is strongly against unethical withdrawal of an article from the publication process after submission. Once the article is submitted, it is the absolute right of the editorial board to decide on article withdrawals. For genuine withdrawal, the corresponding author should submit a request which must be signed by all co-authors explaining the explicit reasons of withdrawing the manuscript.

Accepted articles in final stages of publication if are withdrawn, will entail withdrawal fees. The request will be processed by the editorial board and only serious genuine reasons will be considered if possible. The decision of the editorial board will be final and not negotiable. Unethical withdrawal or no response from the authors to editorial board communication will be subjected to sanction a ban to all authors, and their institute will also be notified.

It is a general principle of scholarly communications, that the editor of a journal is solely and independently responsible for deciding which articles submitted to the journal shall be published. In making this decision the editor is guided by policies of the journal's editorial board and constrained by such legal requirements in force regarding libel, copyright infringement and plagiarism. An outcome of this principle is the importance of the scholarly archive as a permanent, historic record of the transactions of scholarship.

Articles that have been published shall remain extant, exact and unaltered as far as is possible. However, very occasionally circumstances may arise where an article is published that must later be retracted or even removed. Such actions must not be undertaken lightly and can only occur under exceptional circumstances. In all cases, official archives of our journal will retain all article versions, including retracted or otherwise removed articles.

This policy has been designed to address these concerns and to take into account current best practice in the scholarly and library communities. As standards evolve and change, we will revisit this issue and welcome the input of scholarly and library communities. See also the National Library of Medicine's policy on retractions and the recommendations of the International Committee of Medical Journal Editors (ICMJE) concerning corrections and retractions.

### **Article withdrawal**

Only used for Articles in Press which represent early versions of articles and sometimes contain errors, or may have been accidentally submitted twice. Occasionally, but less frequently, the articles may represent infringements of professional ethical codes, such as multiple submission, bogus claims of authorship, plagiarism, fraudulent use of data or the like. Articles in Press (articles that have been accepted for publication but which have not been formally published and will not yet have the complete volume/issue/page information) that include errors, or are discovered to be accidental duplicates of other published article(s), or are determined to violate our journal publishing ethics guidelines in the view of the editors (such as multiple submission, bogus claims of authorship, plagiarism, fraudulent use of data or the like), may be withdrawn.

Withdrawn means that the article content (HTML and PDF) is removed and replaced with a HTML page and PDF simply stating that the article has been withdrawn according to the Policies on Article in Press Withdrawal with a link to the current policy document.

### **Article Retraction**

Infringements of professional ethical codes, such as multiple submission, bogus claims of authorship, plagiarism, fraudulent use of data or the like. Occasionally a retraction will be used to correct errors in submission or publication. The retraction of an article by its authors or the editor under the advice of members of the scholarly community has long been an occasional feature of the learned world. Standards for dealing with retractions have been developed by a number of library and scholarly bodies, and this best practice is adopted for article retraction by us. A retraction note titled "Retraction: [article title]" signed by the authors and/or the editor is published in the paginated part of a subsequent issue of the journal and listed in the contents list. In the electronic version, a link is made to the original article. The online article is preceded by a screen containing the retraction

or where the article, if acted upon, might pose a serious health risk. In these circumstances, while the metadata (Title and Authors) will be retained, the text will be replaced with a screen indicating the article has been removed for legal reasons.

## Article Replacement

In cases where the article, if acted upon, might pose a serious health risk, the authors of the original article may wish to retract the flawed original and replace it with a corrected version. In these circumstances the procedures for retraction will be followed with the difference that the database retraction notice will publish a link to the corrected re-published article and a history of the document.

## Licensing and Copyright Terms

### Copyright

*Biosc Biotech Res Comm* has a policy of copy right, where all the published content of its scholarly articles by its authors can be used for immediate free access to the work and permitting any user to read, download, copy, distribute, print, search, or link to the full texts of articles, crawl them for indexing, pass them as data to software, or use them for any other lawful purpose.

All articles published by Biosc Biotech Res Comm will be distributed Freely under the terms and conditions of the Creative Commons Attribution License (CC-BY) <https://creativecommons.org/licenses/by/4.0/>.

Thus, any one is freely allowed to copy, distribute, and transmit the article on condition that the original article and source is correctly cited.

### Licensing Policy

*Biosc Biotech Res Comm* has a policy of licensing for use and re- use of the published content without any embargo period, following policy that its authors are copyright holders of their scholarly work, granting usage rights to others using [Creative Commons](#) licenses for this purpose.

### Privacy Statement

The names and email addresses entered in the journal site will be used exclusively for the stated purposes of the journal and will not be made available for any other purpose and will not be shared to any other party.

### Guidelines for Reviewers

An unpublished manuscript is a privileged document. Please protect it from any form of exploitation. Don't cite a manuscript or refer to the work it describes before it has been published and don't use the information that it contains for the advancement of your own research or in discussions with colleagues. Adopt a positive, impartial attitude toward the manuscript under review, with the aim of promoting effective and constructive scientific communication.

If you believe that you cannot judge a given article impartially, please return it immediately to the editor. Reviews must be completed within 3 weeks. If you know that you cannot finish the review within that time, immediately return the manuscript to the editor.

In your review, consider the following aspects of the manuscript: –Adherence to style of the MS as set forth in Instructions to Authors of Biosc Biotech Res Comm

- Adequacy of title, abstract and its contents. Language and expression of findings in the manuscript.
- Significance of research questions or subject studied.
- Originality of work: It should be checked through standard plagiarism software only.
- Appropriateness of approach or methodology and adequacy of experimental techniques.
- Appropriateness of figures and or tables and length of article.
- Experimental data its presentation and interpretation.
- Soundness of conclusions and interpretation and relevance of discussion of the manuscript.
- Appropriate literature citations as per Harvard Style of References with updated references.
- Any help you can give in clarifying meaning in the manuscript will be appreciated. We prefer reviewers to use the manuscript comment review system, enabling the authors to make the necessary changes as suggested by the reviewers, which can be later checked for compliance.

If you wish to mark the text of the manuscript, use a pencil or make a photocopy, mark it, and return it together with the original. You can be particularly helpful in pointing out unnecessary illustrations and data that are presented in both tabular (and graphic) form and in detail in the text. Such redundancies are a waste of space and readers time.

A significant number of authors have not learnt how to organize data and will be benefit from your guidance. Do not discuss the paper with its authors. In your comments intended for transmission to the author, do not make any specific statement about the acceptability of a paper. Suggested revision should be stated as such and not expressed as conditions of acceptance. Present criticism dispassionately and avoid offensive remarks.

Organize your review so that an introductory paragraph summarizes the major findings of the article, gives your overall impression of the paper and highlights the major shortcomings. This paragraph should be followed by specific numbered comments which if appropriate may be subdivided into major and minor points. Confidential remarks directed to the editor should be typed (or handwritten) on a separate sheet, not on the review form. You might want to distinguish between revisions considered essential and those judged merely desirable.

Your criticisms, arguments and suggestions concerning the paper will be most useful to the editor and to the author if they are carefully documented. Do not make dogmatic, dismissive statements, particularly about the novelty of work. Substantiate your statements. Reviewer's recommendations are gratefully received by the editor. However, since editorial decisions are usually based on evaluations derived from several sources, reviewers should not expect the editor to honour every recommendation.

### **Editorial Committee of Bioscience Biotechnology Research Communications**

The Editorial committee consisting of the Editor-in-Chief, Executive Editor, Associate Editors, Assistant Editor (s), Journal Managers and the Editorial Secretaries meet frequently to expedite the business of the journal. The editorial committee strictly follows the guidelines provided for international quality and transparent publication.

We strive to follow COPE's Principles of Transparency and Best Practice in Scholarly Publishing <https://publicationethics.org/resources/guidelines-new/principles-transparency-and-best-practice-scholarly-publishing> and encourage our publishing partners to uphold these same principles in general and International Committee of Medical Journal Editors in Uniform Requirements for Manuscripts Submitted to Biomedical Journals: Writing and Editing for Biomedical Publication which can be downloaded from <http://www.icmje.org/>

### **Advisory Board**

An advisory board comprising of members with significant professional experience in different fields of biological and biomedical sciences helps the Editorial Committee of *Bioisc Biotech Res Comm* in all policy matters when needed. Senior advisory board members from India as well as abroad are members of the journal. Each member has been selected due to the expertise and experience in the field of their specializations.

Journal Cover [www.bbrc.in](http://www.bbrc.in)



# CONTENTS

RESEARCH ARTICLES

Biosc.Biotech.Res.Comm. Volume Volume 16 • Number 2 • Apr-May-Jun (2023)

## BIOMEDICAL COMMUNICATION

Predictive Modeling of COVID- 19 Confirmed Cases Using Regressive Objective Regression Methodology 68-72

**Fernando Martínez Fernández, Ricardo Osés Rodríguez, Rigoberto Fimia Duarte, Claudia Osés Llanes, Melba Zayas González, Maribel I Noda Valledor and Dania Artilles Román**

## BIOTECHNOLOGICAL COMMUNICATION

Women with Postpartum Depression in Kerala: A Survey Objective Regression Methodology 73-76

**Pooja Prasad and Balakrishnan Kalamullathil**

## BIOTECHNOLOGICAL COMMUNICATION

BMI-For-Age Percentile as Criteria for Assessment of Obesity in Children and Adolescents 77-83

**Manju Dewan**

## BIOTECHNOLOGICAL COMMUNICATION

Adsorption of Phenol And Resorcinol on Parthenium Based Activated Carbon (PAC) in Basal Salt Medium: 84-90

**Khushi Gandhi, Shambhavee Srivastav, Huda Afreen and Ravi Kant Singh**

## ZOOLOGICAL COMMUNICATION

Avifaunal Abundance of Lumding Forest Reserve Area, Assam, India 91-99

**Prof. Jashodeb Arjun and Rajesh Paul**

## BIOTECHNOLOGICAL COMMUNICATION

Standardization and Evaluation of Buffers: A One Step DNA Extraction Protocol from Microbial Cultures 100-103

**Poornima Shyam, Dr. Shyama Subramaniam, Dr. Shyam Subramaniam and Edwina Jospine M**

## BIOTECHNOLOGICAL COMMUNICATION

Didactic Strategy for the Use of Virtual Itineraries in the Training of Graduates in Education Geography 104-114

**Oliday Aguilar Espinosa, Erich Rodríguez Vallejo, Rafael Armiñana García and Rigoberto Fimia Duarte**

## ZOOLOGICAL COMMUNICATION

Localization of neuropeptide Y(NPY) Immunoreactivity in the Proximal and Distal Intestinal Region of Teleost Fish, Notopterus Notopterus 115-119

**Tulika A Khadse, Yashashree A Gadhikar, Trupti S Khedkar and Sudhir G Chird**

## ZOOLOGICAL COMMUNICATION

Ecology of Biofouling Phytoplankton in Chinnamuttom Harbour Waters Southeast Coast of India 120-125

**P Nithya, B Dhanalakshmi and P Santhanam**

## BIOMEDICAL COMMUNICATION

Knowledge and Management Assessment of Temporomandibular Joint (TMJ) Disorders Among Dental interns at King Saud University 126-131

**Hamad Albagieh, Abdulrahman Alghamdi, Musaad Almeyfayaz, Abdulrahman Alshehri and Osama Alsaiady**

# Predictive Modeling of COVID-19 Confirmed Cases Using Regressive Objective Regression Methodology

Fernando Martínez Fernández<sup>1</sup>, Ricardo Osés Rodríguez<sup>2</sup>, Rigoberto Fimia Duarte<sup>3,4\*</sup>, Claudia Osés Llanes<sup>5</sup>, Melba Zayas González<sup>6</sup>, Maribel I. Noda Valledor<sup>3</sup> and Dania Artilles Román<sup>1</sup>

<sup>1</sup>Faculty of Medicine, University of Medical Sciences of Villa Clara (UCM-VC).

Polyclinic "Marta Abreu" Santa Clara, Villa Clara, 50100 Cuba.

<sup>2</sup>Provincial Meteorological Center of Villa Clara, 50100 Cuba.

<sup>3</sup>Faculty of Health Technology and Nursing (FTSE), University of Medical Sciences of Villa Clara (UCM-VC), 50200 Cuba.

<sup>4</sup>Central University "Marta Abreu" of Las Villas, Villa Clara, 50100 Cuba

<sup>5</sup>Specialist in Hygiene and Epidemiology. Assistant Director of Hygiene and Epidemiology, XX Anniversary Polyclinic, Villa Clara, 50100 Cuba.

## ABSTRACT

The use of predictive models for the evolution of the pandemic is of great help in decision-making by the authorities. The fundamental objective of this work was to obtain through the Regressive Objective Regression, predictions of confirmed cases of COVID-19 in the Marta Abreu Teaching Polyclinic of the city of Santa Clara. In short-term modeling the model was significant at 19.7% with an error of 0.12. Variables dichotomies, saw tooth and saw tooth inverted and risk returned in 1.3, and 12 cases the trend is negative and not significant. We can conclude that a perfect result was obtained in the long term with the ROR methodology. The short-term ROR model depends on the cases of COVID-19 in the previous case, 3 cases back and 12 cases back without significant trend. The long-term model is perfect and depends on the cases of COVID-19 in 12 cases ago, with a negative trend.

**KEY WORDS:** CORONAVIRUS; COVID-19; MATHEMATICAL MODELING; ROR.

## INTRODUCTION

The world has been involved in an epidemiological crisis caused by the new SARS-CoV-2 coronavirus that gave rise to the contagious and deadly disease COVID-19 (Cortellis, 2020; WHO, 2020). Since the first cases were diagnosed in the city of Wuhan, capital of Hubei province, China, the number of infected and dead people has been increasing, which led the World Health Organization (WHO) to declare a pandemic state for COVID-19 on March 11, 2020 (Sun et al., 2020; Wang et al., 2020; WHO, 2020).

According to the Research Group of Mathematical Models in Science and Technology: Development, Analysis, Numerical Simulation and Control (MOMAT) of the Institute of Interdisciplinary Mathematics of the Complutense University of Madrid, Spain, the application of the Be-

CoDiS (Between-Countries Disease Spread) model in the analysis of the COVID-19 pandemic numerically projects that this viral phenomenon will be present for a prolonged period of time in the world (Ivorra & Ramos, 2020).

Several models and methodologies have been applied in the study, analysis and modeling of COVID-19 in the world, where they stand out: Ordinary Differential Equation of First Order (EDOPO), linear type; Simple linear Regression model; Generalized Logistic Growth Model (GLM); Structured Susceptible-Exposed-Infected-Removed (SEIR)/SEIR model; the Bayesian Probability Mathematical Model; SIRD model (also employed in Cuba); Conceptual Model and the Simulation Model, among many others (Rue et al., 2017; Simpson et al., 2017; Osés et al., 2021a,b).

The main objective of the present research was to obtain, by means of the Objective Regressive Regression (ORR) methodology, predictions of confirmed cases of COVID-19

**Article Information:**\*Corresponding Author: [rigoberto.fimia66@gmail.com](mailto:rigoberto.fimia66@gmail.com)

Received 25/11/2022 Accepted after revision 25/05/2023

Published: June 2023 Pp- 68-72

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <http://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.1>

in the "Marta Abreu" Teaching Polyclinic in the city of Santa Clara, Villa Clara province, Cuba.

## MATERIAL AND METHODS

In the research, the daily data from January to March corresponding to the year 2021 of the number of positive cases to COVID-19 in the Teaching Polyclinic "Marta Abreu" of Villa Clara Cuba were used, from a total of 3 294 cases evaluated in consultation, 58 were positive.

The prognosis was carried out with the use of the methodology of Regressive Objective Regression (ROR) which has been implemented in different variables such as viruses and bacteria circulating in Villa Clara province. Objective Regressive Regression (ORR) modeling is based on a combination of Dummy variables with modeling (ARIMA), where two Dummy variables are created and the trend of the series is obtained; it requires few cases to be used and also allows the use of exogenous variables that make it possible to model and forecast in the long term, depending on the exogenous variable (Osés et al., 2020).

In the ROR methodology, in a first step, dichotomous variables DS, DI and NoC are created, where: NoC: Number of cases of the base, DS = 1, if NoC is odd; DI = 0, if NoC is even, when DI=1, DS=0 and vice versa. Subsequently, the module corresponding to the Regression analysis of the statistical package SPSS version 19.0 (IBM Company) is executed, specifically the ENTER method where the

predicted variable and the ERROR are obtained. Then the autocorrelograms of the ERROR variable will be obtained, paying attention to the maximums of the significant partial autocorrelations PACF. The new variables are then calculated according to the significant Lag of the PACF. Finally, these regressed variables are included in the new regression in a process of successive approximations until a white noise in the regression errors is obtained.

**Ethical aspects:** The research was subject to ethical standards, where all the information collected and provided was used only for the stated purpose. It did not involve physical or psychological affectations, in order to be able to generate new knowledge without violating the ethical principles established for these cases. On the other hand, all authors involved in the research, publication and dissemination of the results are responsible for the reliability and accuracy of the results shown (DHAMM, 2013).

## RESULTS AND DISCUSSION

The great development acquired by different sciences, such as epidemiology, mathematics, statistics and computer science, has led to the generation of new techniques in emerging and recurrent procedures in the field of biostatistics and bioinformatics (spatial statistics, neural networks and functional data analysis, among others) to describe the importance of mathematical models in infectious diseases (Martí & Caylà, 2009).

**Table 1. Descriptive statistics of confirmed patients at the "Marta Abreu" Polyclinic. Year 2021**

	Descriptive statistic				
	N	Minimum	Maximum	Half	Typ. desv.
Risk COVID-19	3294	.00	1.00	.0176	.13154
N valid (according to list)	3294				
Positives MA	3	7	31	19.33	12.014
N valid (by list)	3				

**Figure 1: Distribution of patients according to age. January-March 2021**




Table 1 calculated the descriptive statistics of the risk of contracting COVID-19, the value of zero was assigned if the PCR was negative, and if the PCR was positive, the value

of Risk 1; the mean value of this variable corresponds to 19.33 cases with a standard deviation of 12.01 cases.

The distribution according to age was irregular, with the highest value corresponding to patients aged 1 to 18 years, followed by those aged 19 to 29 years; children under 1 year of age were also present in the study (Figure 1).

COVID-19 cases were modeled in the short term using the ROR methodology. The model was significant at 19.7% with an error of 0.12 (Table 2). As can be seen in the table the Durbin-Watson statistic is close to 2, so we are in the presence of an adequate explanation of the variance using this model.

1. Predictor variables: Lag12Risk, Lag3Risk, Lag1Risk, DI, SD, NoC.
2. For regression through the origin (the model with no

intersection term), R-squared measures the proportion of the variability of the dependent variable explained by regression through the origin. The above CANNOT be compared with R-squared for models that include an intersection.

3. Dependent variable: RiskCOVID19
4. Linear regression through the origin

The ROR modeling of predictions obtained yielded very significant results for the study of the COVID-19 pandemic in the "Marta Abreu" Teaching Polyclinic, with a Fisher's

F of 22 significant at 100% (Table 3), thus proving that the sample variances are not equal.

Table 3. Analysis of variance of the model for COVID-19

1. Predictor variables: Lag12Risk, Lag3Risk, Lag1Risk, DI, SD, NoC.
2. This total sum of squares has not been corrected for the constant because the constant is zero for regression through the origin.
3. Dependent variable: COVID-19 risk.
4. Linear regression through the origin

Table 2. Summary of the model using Objective Regressive Regression

Model	Summary of model <sup>c,d</sup>				
	R	R squared <sup>b</sup>	R-squared corrected	Standard error of estimation	Durbin-Watson
1	.197a	.039	.037	.12352	2.009

Table 3. Analysis of variance of the model for COVID-19

Model	ANOVA <sup>c,d</sup>				
	Sum of squares	gl	Quadratic mean	F	Sig.
Regression	2.019	6	.336	22.052	.000a
Residual	49.981	3276	.015		
Total	52.000b	3282			

1. Dependent variable: COVID-19 risk.
2. Linear regression through the origin

Next, the model was run only with Lag12 Risk in order to have a long-term valuation by entering the Step POSITIVE variable in the model, obtaining a model that explains 100% of the variance of the model (Table 5).

1. Predictor variables: Step2133, Step1656, Step1655, Step1652, Step1644, Step1336, Step75, Step74, Step14, Step POSITIVE, DS, DI, Lag12Risk, NoC.
2. For regression through the origin (the model with no intersection term), R-squared measures the proportion of the variability of the dependent variable explained by regression through the origin. The above CANNOT be compared to R-squared for models that include an intersection.

The ROR model in question is made up of the following variables, DI and DS which are dichotomous variables, saw tooth DS and inverted saw tooth DI and of the regressed risk in 1,3 and 12 cases (Lag1 Risk; Lag3 Risk and Lag12 Risk). The trend is negative and not significant (Table 4).

Table 4. Results of the application of the ROR model plus trend

Model	Coeficientes <sup>a,b</sup>					
	Unstandardized coefficients	Typified coefficients				Sig.
	B	Error tip.	Beta	t		
	DS	.016	.005	.091	3.316	.001
	DI	.016	.005	.091	3.315	.001
	Tendency	-2.591E-6	.000	-.039	-1.136	.256
	Lag1Risk	.116	.017	.116	6.719	.000
	Lag3Risk	.057	.017	.057	3.326	.001
	Lag12Risk	.065	.016	.069	3.988	.000

1. Dependent variable: COVID-19 risk.
2. Linear regression through the origin

we are dealing with a perfect model, i.e., the result of the variables is equal.

Table 6 shows the analysis of variances using Fisher's test (F). In the results, Fisher's F can't be seen, which infers that

1. Predictor variables: Step2133, Step1656, Step1655, Step1652, Step1644, Step1336, Step75, Step74,

- Step14, Step POSITIVE, DS, DI, Lag12Risk, NoC.
- 2. This total sum of squares has not been corrected for the constant because the constant is zero for regression through the origin.
- 3. Dependent variable: COVID-19 risk.
- 4. Linear regression through the origin

Epidemic modeling is historical and has been refined with the advancement of science and technologies, demonstrating the essential role of mathematics in indicating where the disease may move and giving suggestions on how to decide (Guinovart, 2020). COVID-19 disease has been recognized as a global threat, in which predictive models for the epidemiological trend of its prevalence and incidence have

been being used worldwide (Osés et al., 2018; Prades, 2020; Osés et al., 2021c).

In the present research article, predictions were exposed for the cases confirmed to COVID-19 in the "Marta Abreu" Teaching Polyclinic. The increase in the number of children and adolescents infected by the new coronavirus in the mentioned health area showed that they are as susceptible to the disease as the rest of the population. Children and older adults should follow the same biosecurity measures as those prescribed by the Ministry of Public Health for the entire population. Coinciding with the daily reports of MINSAP, the highest number of cases was reported in females (MINSAP, 2021).

**Table 5. Model run with Lag12 in the long term**

Modelo	Summary of model <sup>c,d</sup>				
	R	R squared <sup>b</sup>	R-squared corrected	Standard error of estimation	Durbin-Watson
1	1.000 <sup>a</sup>	1.000	1.000	.0000	1.696

**Table 6. Analysis of Variance of the model for COVID-19**

Model	ANOVA <sup>c,d</sup>				
	Sum of squares	gl	Quadratic mean	F	Sig.
Regression	52.000	14	3.714	.	. <sup>a</sup>
Residual	.000	3268	.000		
Total	52.000 <sup>b</sup>	3282			

The results of the present investigation coincide with those obtained in a similar study conducted in the province of Cienfuegos by Medina (2020); where logistic and exponential models (specifically the Gompertz growth model) were used to obtain estimates. They state that the statistical prediction models obtained gave very significant results for the study of the COVID-19 pandemic in Cuba, which also agrees with the results obtained by other authors in this regard (Medina et al., 2020).

## CONCLUSION

The application of ROR modeling provides Public Health authorities and decision makers with information on the short- and medium-term behavior of variables of great interest for understanding the spread of SARS-CoV2. With the use of the ROR methodology, it was possible to predict the future number of positive cases in the "Marta Abreu" health area, making this model a valuable practical tool for correct and timely decision making.

## REFERENCES

Cortellis, T.M. (2020). Disease Briefing: Coronaviruses. A Clarivate Analytics solution. 3.23.2020. 51 pp. In: [https://clarivates.com/wp-content/uploads/dlm\\_uploads/2020/01/](https://clarivates.com/wp-content/uploads/dlm_uploads/2020/01/)

## CORONAVIRUS-REPORT-23.3.2020.pdf

DHAMM (Declaración de Helsinki de la AMM). (2013). Principios éticos para las investigaciones médicas en seres humanos. 64<sup>a</sup> Asamblea General, Fortaleza, Brazil, octubre. World Medical Association, Inc. – All Rights reserved. 9 pp.

Guinovart, D.R. (2020). Intervención en la mesa redonda. Sitio de la Presidencia de Cuba. <http://www.cubadebate.cu/noticias/2021/04/27/lo-afirman-los-expertos-pronosticos-de-la-covid-19-en-cuba-son-desfavorables/>

Ivorra, B., Ramos, A.M. (2020). Validation of the forecasts for the international spread of the coronavirus disease 2019 (COVID-19) done with the Be-CoDiS mathematical model. Reporte Técnico Grupo MOMAT, Universidad Complutense de Madrid, 1–14. DOI: <http://www.doi.org/10.13140/RG.2.2.31460.94081>

Martí, C.K.G., Caylà, J.A. (2009). Modelos matemáticos utilizados en el estudio de las enfermedades transmisibles. Rev Esp Salud Pública 83 (5). Disponible en: [https://www.mscbs.gob.es/biblioPublic/publicaciones/recursos\\_propios/resp/revista\\_cdrom/vol83/vol83\\_5/RS835C\\_689.pdf](https://www.mscbs.gob.es/biblioPublic/publicaciones/recursos_propios/resp/revista_cdrom/vol83/vol83_5/RS835C_689.pdf)

Medina, M.J., Cortés, C.M., Cortés, I.M., Pérez, F.A., Manzano, C.M. (2020). Estudio sobre modelos predictivos para la COVID-19 en Cuba. Medisur 18: 431-442. Disponible en: <http://medisur.sld.cu/index.php/medisur/article/view/4703>

MINSAP. (2021). Para prevenir y enfrentar el nuevo coronavirus, la participación popular es esencial. Disponible en: <https://salud.msp.gob.cu/para-prevenir-y-enfrentar-el-nuevo-coronavirus-la-participacion-popular-es-esencial/>

Osés, R.R., Carmenate, R.A., Pedraza, M.A.F., Fimia, D.R. (2018). Prediction of latitude and longitude of earthquakes



at global level using the Regressive Objective Regression method. *Advances in Theoretical & Computational Physics (Adv Theo Comp Phy)* 1: 1-5.

Osés-Rodríguez, R., Osés-Llanes, C., Fimia-Duarte, R., González-Meneses, A., & Iannacone, J. (2020). Pronostico de la COVID-19 por medio de la metodología de Regresión Objetiva Regresiva en Villa Clara y Cuba. *The Biologist* 18 (2). Disponible en: <https://doi.org/10.24039/rtb2020182754>

Osés, L.C., Osés, R.R., Fimia, D.R., Zambrano, G.M.P., Wilford, G.F.M. (2021a) Comparison of lineal ROR vs Nonlinear Weibull model for COVID-19 in Iraq. *Himalayan Journal of Applied Medical Sciences and Research* 2: 88-96.

Osés, R.R., Osés, C.L., Fimia, D.R., González, M.A., Iannacone, J., Bruna, S.T., Wilford, G.F.M. (2021b). Age Prediction for COVID-19 Suspects and Contacts in Villa Clara Province, Cuba. *EC Veterinary Science* 6: 41-51.

Osés, R.R., Fimia, D.R., Osés, L.C., Zambrano, G.M.P., Santos, Z.T.B., González, M.A. (2021c). Forecast of New and Deceased Cases of COVID-19 in Cuba with an Advance of 105 Days. *Acta Scientific Veterinary Sciences* 3: 31-36.

Prades, E.E., Martin, S.D. (2020). Modelos estadísticos

para las predicciones de la COVID-19 en Cuba. *Revista Cubana De Higiene y Epidemiología* 57. Disponible en: <http://www.revepidemiologia.sld.cu/index.php/hie/article/view/303/275>

Rue, H., Riebler, A., Sarbye, H., Illian, J.B., Simpson, D.P., Lindgren, F.K. (2017). Bayesian computing with INLA: a Review. *Annual Reviews of Statistics and its Applications* 4: 395-421.

Simpson, D.P., Rue, H., Martins, T.G., Riebler, A., Sarbye, S.H. (2017). Penalising model component complexity: a principled, practical approach to constructing priors. *Statistics Sciences* 32: 1-46.

Sun, Z., Thilakavathy, K.T., Kumar, S.S., He, G., Liu, S.V. (2020). Potential Factors Influencing Repeated SARS Outbreaks in China. *International Journal of Environmental Research and Public Health* 17: 16-33.

Wang, L.S., Wang, Y.R., Ye, D.W., Liu, Q.Q. (2020). A review of the 2019 Novel Coronavirus (COVID-19) based on current evidence. *International Journal of Antimicrobial Agents*. Doi: 10.1016/j.ijantimicag.2020.105948

WHO. (2020). Conferencia de la OMS. <https://www.who.int/docs/default-source/coronaviruse/transcripts/who-audio-emergencies-coronavirus-press-conference-full-and-final-11mar2020.pdf?sfvrsn=cb432bb32>



# Women with Postpartum Depression in Kerala: A Survey

Pooja Prasad\* and Balakrishnan Kalamullathil

Department of English and Languages, Amrita Vishwa Vidyapeetham, Kochi Campus, Kerala, India.

## ABSTRACT

The increase in number of Postpartum Depression (PPD) cases in Kerala is increasing day by day. So, a study on the awareness of PPD has become necessary in this scenario. This study tries to quantify the awareness among women in Kerala about PPD and covers how they tackled the issue through various methods used by them to cope with the issue. The study is conducted using an online survey method. A prepared questionnaire was circulated online among 150 young mothers who were born and brought up in Kerala. The questionnaire consisted of eight questions about PPD and baby blues. Each question was provided with options from which the participants had to choose one. The data thus collected were studied and analyzed. The results of the survey were analyzed to arrive at a conclusion. There was active participation from the participants' side. The results showed that the percentage of participants who were well aware of PPD was very low. The study brought to light that, much attention and activities are needed to solve PPD cases in Kerala. Most women who suffer from PPD are reluctant to seek medical help. This situation can change through proper campaigns and other related activities.

**KEY WORDS:** ANXIETY, AWARENESS, COPING MECHANISM, POSTPARTUM DEPRESSION, TREATMENT.

## INTRODUCTION

Postpartum depression is a serious issue faced by women in present-day society. Postpartum depression (PPD) is a medical condition that can be cured with proper care and treatment. In women, depression can occur during and after pregnancy. Depression after delivery can occur as 'baby blues' that last only for one or two weeks after childbirth. It has mild symptoms like mood swings, anxiety, and insomnia. A more severe condition is postpartum depression. It is long-lasting than baby blues and shows intense symptoms including anxiety and panic attacks, sadness, irritability, severe mood swings, appetite problems, difficulty with bonding with your baby, thoughts about harming baby or yourself and severe anger (Jayarajan 2021).

The lack of awareness about Postpartum Depression increases the depth of the problem. Studies show that the lack of early detection of PPD also worsens the condition. The awareness of the real problem, that PPD is a serious issue and can affect the female's later life will bring some changes in the present situation (Zauderer 2009). The purpose of the study is to bring the problem of PPD to the forefront and thus reduce the risk women face due to PPD nowadays. The outcome of this study will help where people are unaware of the seriousness of PPD and where no proper care is given

to women suffering from PPD. The study tries to quantify women awareness of the issue (Jayarajan 2021).

The relevance of the study is high as no such studies have ever been conducted in Kerala in recent times. Moreover, a study like this can contribute to the area of mental health, the overall well-being of a society and the development of the health sector of Kerala.

## MATERIAL AND METHODS

An in-depth analysis of Postpartum Depression was conducted using a survey. The study was conducted by circulating the prepared questionnaire among 150 young mothers. The sample of the study was selected after much research on the topic. The ages of the participants were in the range of 25 – 40. Mothers who gave birth in the last 10 years, and who were born and brought up in Kerala were considered for this study.

This was done to examine the recent developments in the area, especially in Kerala. The research subjects were verified to be cognizant of English and all the questions were in English. The participants were well informed about the intention of the survey. They were informed to read the instructions clearly and answer the questions. It was also assured that their details will be kept confidential. The survey mainly aimed at checking the knowledge/awareness women in Kerala had about PPD. Moreover, the study checked how they came to know about PPD and how far they were aware

**Article Information:**\*Corresponding Author: [prasadpooja710@gmail.com](mailto:prasadpooja710@gmail.com).

Received 25/03/2023 Accepted after revision 22/06/2023

Published: June 2023 Pp- 73-76

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.2>

of the issue. It also helped to collect information about their personal experience of PPD, and how they overcame it. The problem of bonding with the baby was also included as one question.

The questionnaire consisted of 8 questions about PPD and baby blues. These were intended to collect information about the level of awareness women had about PPD, how they came to know about it, have they attended any awareness programs etc. The survey also tried to collect information about the coping mechanism the participants chose to overcome the situation. Responses to this particular question revealed how worse were the condition and what percentage of women seeks medical help.

There were questions about the symptoms they had to suffer and about the time they experienced it. Each of the questions was given options from which the participants can choose one. The number of the options varied from question to question, that is, from 2 – 5. The research subjects were verified to be cognizant of English and all the questions were in English (Jayarajan 2021). The present study has been approved by the Institutional Ethics Committee of Amrita Vishwa Vidyapeetham, Coimbatore. All due permissions have been taken by the concerned authorities including consent etc.

## RESULTS AND DISCUSSION

The intensity of the psychic problems faced by women suffering from PPD needs to be discussed in detail, and a long-lasting solution has to be found. An important part of reducing the percentage of patients can be done by giving proper awareness. Not only females but also males and people of other genders have to be educated about the mental condition of women during and after pregnancy. Medical professionals suggest providing proper awareness can reduce the problem in a considerable way. This can contribute a lot to the healthy development of the baby and mother (Zauderer 2009).

**Figure 1: Pie chart showing the awareness of participants about Baby Blues**






Figure one shows that only about 53.9% of the participants heard about baby blues. This revealed the intensity of the situation. The participants included in the other section, that was, the 46.1%, might be the ones who have gone through

the same. But unawareness of the real problem worsens the condition. As baby blues last only for two or three weeks it is not a dangerous problem as PPD. But even in the case of PPD, a considerable percentage of people who were unaware of PPD was found. Figure 2 shows that 14.8 % of the participants were unaware of postpartum depression. And the results show that people were not much aware of serious problems like PPD as they are of Baby Blues. This condition has to be changed through continuous practices to make the general public aware of PPD and its symptoms (Jayarajan 2021).

**Figure 2: Pie chart showing the awareness of participants about PPD**



**Figure 3: Pie chart showing the response of participants to whether they faced symptoms of PPD**




There exists a social stigma in India, especially in Kerala, to consult a psychiatrist or a psychologist (The Hindu 2018). This is another reason which prevents the cure of PPD through proper treatment. People are reluctant to admit the fact that they are facing some mental problems. The same kind of reluctance is there in the case of PPD also. They fear a kind of 'othering' from society and its roots can be traced back to the stigma toward a madness that existed in the middle ages. This tendency limits the possibility to take proper medical care. This can also be controlled by conducting awareness programs and thus normalizing mental problems and depression (Foucault 1988; Kuriakose et al. 2020; Jayarajan 2021).

Figure three and four brings out the seriousness of the situation. In Figure three, one can see that about 77.3% of participants faced the symptoms of PPD. This is proof of the problem faced by women after delivery. Figure four gives

the period for which they faced these difficulties. 19.2% faced symptoms of PPD up to one year after delivery/C-section. For 28.3% it lasted for six months, and for 24.2% it lasted up to one month. Only 28.3% experienced it for two weeks after childbirth (Kuriakose et al. 2020).

**Figure 4: Pie chart showing how much time did the participants felt the symptoms**



**Figure 5: Pie chart showing the level of difficulty the participants faced in bonding with the baby**





Figure 5 shows that 8.6% of the participants found it difficult to create a bond with the baby and 27.3% of them felt the same for a short period. This is a period that is highly dangerous and may lead up to harming a child and oneself. Recent news reports give evidence of this were mothers in their depressive state harmed babies and some violent acts ended up in their death (Jayarajan 2021). The results of the survey indicate that most of the participants have gone through severe Postpartum Depression and all of them needed medical help. But Figure 6 shows the real scenario of Kerala's treatment of a serious mental issue like PPD (Jayachandran 2021).

**Figure 6: Pie chart showing how participants with PPD cope up with it**



Even though most of the participants faced severe PPD-related problems only 0.8% of them sought medical help and only 1.6% of them were ready to seek the help of a psychologist or psychiatrist. This reveals the pathetic condition of Kerala concerning PPD. This situation can be

changed by giving proper awareness about the issue. The collected data shows that only 7% of the participants had attended any awareness programs related to PPD. Most of them learned about this issue through social media. And some of them learned through articles and only 1.7% of them got informed through the newspapers. This shows that the government, health department, as well as the public, have to play a significant role in reducing the difficulties faced by mothers and newborns (Kuriakose et al. 2020).

The study, 'Prevalence of depression among middle-aged women in the rural area of Kerala', conducted by Archana P S, Soumitra Das and Sairu Philip et al., tries to determine the prevalence of depression among women aged 40 – 60 years in the rural area of Kerala using Patient Health Questionnaire 9 (PHQ – 9). The setting of the study is Ambalappuzha, a rural area in the district of Alappuzha. (Archana et al. 2017; Kuriakose et al. 2020). In contrast, this study tried to include young women from all over Kerala who gave birth in the last 10 years. Moreover, this study concentrates only on depression that comes as a result of pregnancy and/or delivery.

## CONCLUSION

The findings of the present study quantify the awareness young mothers in Kerala have about PPD. As this study deals with a health-related issue which is related to the well-being of society, it has significance in the present scenario. This study prompts studies in the future that may lead to change in the current situation. Proper awareness about the issue can be given through newspapers, news channels, social media and other media. Constant active participation from the public and initiatives from the government is needed to improve the situation.

## ACKNOWLEDGEMENTS

The study was supported by the Department of English Language and Literature faculty in Amrita School of Arts and Sciences, Kochi, Kerala, India.

**Conflict of Interests:** Authors declare no conflict of interests to disclose.

## REFERENCES

- Archana, PS, Das, S, Sairu, P et al. (2017). Prevalence of depression among middle aged women in the rural area of Kerala. *Asian Journal of Psychiatry*, volume 29. <https://doi.org/10.1016/j.ajp.2017.05.016>.
- Foucault, M (1988). *Madness and civilization: a history of insanity in the Age of Reason*. London, Routledge Classics, pp. 57-64. <https://jamanetwork.com/journals/jama/article-abstract/194624> (Accessed: 10 November)
- Jayarajan, S (2021). Why Kerala needs to address postpartum depression and psychosis in new mothers. *The News Minute*. 11 January [online]. Available at: <https://www.thenewsminute.com/article/why-kerala-needs-address-postpartum-depression-and-psychosis-new-mothers-141245> (Accessed: 1 October 2021)
- Kuriakose, S, Vinaychandran, S, Narayanan, D et al.

(2020). Postpartum depression and its association with social support: a cross-sectional study at a maternity hospital in Kerala.

Kerala Journal of Psychiatry, volume 33(2) [online] Available at: [https://www.researchgate.net/publication/350018707\\_Postpartum\\_Depression\\_And\\_Its\\_Association\\_With\\_Social\\_Support\\_A\\_Cross,Sectional\\_Study\\_At\\_A\\_Maternity\\_Hospital\\_In\\_Kerala](https://www.researchgate.net/publication/350018707_Postpartum_Depression_And_Its_Association_With_Social_Support_A_Cross,Sectional_Study_At_A_Maternity_Hospital_In_Kerala) (Accessed: 14 November 2021)

Miller, LJ (2002). Postpartum depression. *Jama*, 287(6), pp.762-765. <https://jamanetwork.com/journals/jama/>

[article-abstract/194624](https://www.researchgate.net/publication/350018707_Postpartum_Depression_And_Its_Association_With_Social_Support_A_Cross,Sectional_Study_At_A_Maternity_Hospital_In_Kerala)

The Hindu (2018). Remove social stigma of mental illness: expert, *The Hindu*, 8 June [Online] Available at: <https://www.thehindu.com/news/national/kerala/remove-social-stigma-of-mental-illness-expert/article24116979.ece> (Accessed: 3 October 2021) .

Zauderer, C (2009). Postpartum depression: how childbirth educators can help break the silence. *The Journal of Perinatal Education*, volume (18) [online] Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2684038/> (Accessed: 6 October 2021).



# BMI-For-Age Percentile as Criteria for Assessment of Obesity in Children and Adolescents

Manju Dewan

Associate Professor, DAV College, Sector 10, Chandigarh, India

## ABSTRACT

The most useful population-level measure of overweight and obesity is BMI. This measure is same for men and women of all ages. But it should not be thought to be as final factor to calculate the same body fat percentage in different individuals. Various proposed reference values for overweight and obesity exist for BMI, but the combined effect of BMI and BMI for age percentile recommendations is scarce. A total 2048 children participated in the epidemiological survey. According to Body Mass Index in the study, 66.26% subjects were found to be malnourished in the total population. The degree of obesity had been observed as Pre-obese 3.81% and Obese-I 3.81% and Obese-II were 0.88%. Pre-obese, obese-I and obese-II boys were more than the girls. According to BMI-for-age percentile, overall, 6.20% of children and adolescents were at risk of overweight and 4.39% were overweight during health survey while 24.80% are underweight.

**KEY WORDS:** BMI, BMI-FOR-AGE PERCENTILE, OBESITY, OVERWEIGHT.

## INTRODUCTION

Overweight and obese children can develop obesity in adulthood and can develop diabetes, hypertension and cardiovascular diseases at an early age (Krushnapriya et al 2015). The highest prevalence rates of childhood obesity have been observed in developed countries; however, its prevalence is increasing in developing countries as well (Popkin and Doak, 1998). Females are more likely to be obese as compared to males, owing to inherent hormonal differences (Gupta, 2009). It is difficult to create one parameter for the measurement of overweight and obesity in children and adolescents because there are number of physiological changes in children and adolescents.

Depending on the age, different methods to measure a body's healthy weight are available. BMI fails to distinguish between fat and fat-free mass (muscle and bone) and may exaggerate obesity in large muscular children. Body mass index (BMI) is a recommended index of obesity by World Health Organization (1996) and is related to various disease risks (Stevens et al, 1998). BMI is the most useful population-level measure of overweight and obesity as it is same for men and women, and for adults of all ages. However, it should not be considered as final parameter to calculate the same body

fat percentage in different individuals. Various proposed reference values for overweight and obesity exist for BMI, but the combined effect of BMI and BMI for age percentile recommendations are scarce. Therefore, the primary aim of this study is to compare the validity of BMI and BMI for age percentile as diagnostic tests for sample of 10-19 years old children and adolescents.

## MATERIAL AND METHODS

A total 2048 children participated in the epidemiological survey. Children who had participated in the study also underwent the health examination. There were numerous students who remained absent on the test date and on all subsequent make-up sessions. Many parents requested to excuse their children from the test. For interested children, consent form was signed by the parents before starting the survey work. Children those completed all steps of the survey were included and those who missed even a single step excluded. The World Health Organization, U.S. Centers for Disease Control and Prevention and International Obesity Task Force provide separate definitions of overweight and obesity in children and adolescent. Anthropometric measurements including height, weight, and waist and hip circumferences were taken to explore the incidence of obesity and the various factors associated with it.

**Body Mass Index (BMI):** This measure was used to assess body weight relative to height. Body mass index is a useful

**Article Information:**\*Corresponding Author: [manjudewan72@gmail.com](mailto:manjudewan72@gmail.com)

Received 04/04/2023 Accepted after revision 25/06/2023

Published: June 2023 Pp- 77-83

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <http://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.3>

tool in both clinical and public health practice for assessing nutritional status of the population.

Underweight	BMI less than 18.5
Ideal	BMI 18.5-22.9
Overweight	BMI 23-24.9
Obese – should lose weight	BMI 25.0-29.9
Very obese – lose weight now	BMI greater than 30.0

**BMI-for-Age Percentile:** The Centers for Disease Control and Prevention (CDC),2000 suggested the levels of concern for children based on the BMI-for-age percentile as follows:-

- Below the 5th percentile : **Underweight**
- 5th percentile < 85th percentile : **Healthy weight**
- 85th percentile to < 95th percentile : **At risk of overweight**
- 95th percentile and above : **Overweight**

A total 2048 children had participated in the epidemiological survey. They had also undergone the health examination. There were numerous students who remained absent on the test date and on all subsequent make-up sessions. Many parents requested to excuse their children from the test. For interested children, a consent form was signed by the parents before starting the survey work. Children those completed all steps of the survey were included and those who missed even a single step were excluded. Anthropometric measurements including height, weight and waist and hip circumferences were taken to explore the incidence of obesity and the various factors associated with obesity.

These measures were used to assess body weight relative to height. Body Mass Index is a useful tool in both clinical

and public health practice for assessing nutritional status of the population. Underweight: BMI less than 18.5; Ideal BMI: 18.5-22.9; Overweight: BMI 23-24.9; Obese: BMI 25.0-29.9; Very obese: BMI greater than 30.0.

The Centers for Disease Control and Prevention (CDC) 2000 suggested the levels of concern for children based on the BMI-For-Age Percentile as Below the 5th percentile: Underweight; 5th percentile < 85th percentile: Healthy weight; 85th percentile to < 95th percentile: At risk of overweight; 95th percentile and above : Overweight. BMI allows comparison between children of the same sex and age (Connor and Arif ; 2021).

## RESULTS AND DISCUSSION

A total 2048 children participated in the epidemiological survey. Out of these 2048 children, 1017 were from urban population and 1031 were from rural population. According to body mass index in the study, 66.26% subjects were found to be malnourished in the total population. The degree of obesity had been observed as Pre-obese 3.81% and Obese-I 3.81% and Obese-II were 0.88%. (Table-1 Graph-1).

According to BMI-for-age percentile, overall, 6.20% of children and adolescents were at risk of overweight and 4.39% were overweight during health survey while 24.80% are underweight.


Overweight children were highest among urban (6.39%) than rural (2.42%). (Table-2 Graph-2).

Obesity and overweight result from an imbalance in the energy expenditure equation such that either food intake is in excess; energy expenditure is deficit or both. Obesity presently has been properly documented as an ailment in its own right which is preventable through changes in life style especially diet. Obesity is a major determinant of many non-communicable diseases and induces diabetes mellitus and coronary heart disease.

**Table 1. Prevalence of overweight and obesity according to BMI Categories**

Group/Sub Group	Below 18.5		18.5-22.9		23.0-24.9		25.0-29.9		Above 30	
	N	%	N	%	N	%	N	%	N	%
All Data	1357	66.26	517	25.24	78	3.81	78	3.81	18	0.88

**Figure 1**



Physiologically, the body weight is constantly changing. Obesity is the result of excessive expansion of adipose tissue mass. This disorder is often accompanied by abnormalities in systemic carbohydrate and lipid metabolism and the secretion and action of insulin alterations thought to reflect diabetogenic effect of obesity. There remain questions about the best diagnostic criteria for this age group.

According to Dietz and Bellizzi, 1999, overweight, during childhood and adolescence, is characterized by a body mass index for age (BMI/age) above the 85th percentile and obesity by a BMI/age above the 95th percentile, based on the National Health and Nutrition Examination Survey




(NHANES II-III) standards. Other authors say it is possible to adopt absolute BMI limits for specific age groups, during the growth period, without considerably compromising diagnostic specificity or sensitivity (Monteiro et al, 1999).

An expert committee in 1997 declared BMI for age to be the principal measure of obesity in childhood. The adoption of regional BMI standards has also been proposed (Anjos et al, 1998).

**Table 2. Prevalence of overweight and obesity according to BMI for age percentile Categories**

Group/Sub Group	Under Wt.		Healthy Wt		Risk Ov.Wt		Over Wt.	
	N	%	N	%	N	%	N	%
All Data	508	24.80	1323	64.60	127	6.20	90	4.39

**Figure 2**



Diagnostic criteria should be simple, low cost, reproducible and reliable and offer elevated sensitivity and specificity, thus reducing false positive and negative diagnoses. Based on a pool of studies of BMI profiles by age in a number of different countries, including Brazil, proposed limits for overweight and obesity to be applied to the 2 to 20 age group internationally (Cole et al., 2000).

In this study, the prevalence of obesity was more in boys than in girls. It is possible that the Western feminine self-image and fear of obesity were more marked in girls than in boys. Girls as young as 5 years of age care more about their self-image and associate it to weight status. The same has been reported in other studies where boys had higher prevalence of obesity than girls at all ages (Brown et al, 2002 and Huang et al, 2003).

WHO in 1996 stated that BMI values are not dependent on age or gender. But, they may not narrate equally to levels of obesity in diverse populations (certain racial groups in Africa and Asia) as a result of differing body-proportions. Also, the BMI index may not accurately reflect the fatness levels of people with builds falling far outside the average. As mentioned earlier active persons like sportsmen and sportswomen, will tend to have BMI levels of "overweight" or even 'obese', due to their higher levels of muscle mass. BMI categories also fail to account for varying proportions of fat, bone, cartilage, water weight and more.

Centers for Disease Control and Prevention (CDC), 2000 stated that "overweight" is defined differently for children and teenagers than for adults. Due to growth and differing

maturation rates, a modified BMI compares heights and weights against growth charts which also consider gender. This index is known as the BMI-for-age percentile (for children 2-19 years of age), and reflects BMI as compared with boys and girls of the same age. Because of that increase, the American Diabetes Association and the American Academy of Pediatrics recommend routine screening of children at higher risk for type 2 diabetes. It is essential to identify the best anthropometric index in any population to predict chronic disease risk.

In the present study, according to BMI-for-age percentile, 24.80% children were underweight in of 10-19 age groups. In the present study 6.20% of children and adolescents were at risk of overweight and 4.39% were overweight during health survey. The results of the present study also shows that it is appropriate to choose BMI / age, based on the international standards, as a diagnostic indicator for overweight and obese children.

Worldwide one in 10 children, aged 5-17 year old, is overweight, a total of 155 million, of which around 30-45 million are obese (Lobstein et al, 2004). The prevalence rate of overweight in children under 5 years of age in developing countries is 3.3% or 17.6 million (Executive Board, 2001). Countries with the highest prevalence of overweight are located mainly in the Middle East, North Africa, and Latin America (de Onis et al, 2000). In the United States, the prevalence of childhood overweight tripled between 1980 and 2000 (Ogden et al, 2002). In Australia, the prevalence of childhood overweight almost doubled between 1985 and 1995 (Magarey et al, 2001). In Europe there are 14 million children who are overweight and 3 million of them are obese. In 2002 nearly a quarter of children within the European Union were overweight, higher than the predicted peak for 2010. (Lobstein et al, 2004). In India prevalence of overweight/obesity was:

- Delhi(North) Overweight, 22%; obesity 6%.(Sharma et al, 2007);
- Uttar Pradesh (North) Obesity, 7.6% (Gupta et al, 1999).

In the present study, as per the BMI for age percentile, 6.20% of children were at risk of overweight and 4.39% were overweight which less than the other studies. But the necessary steps must be taken at this stage to prevent the

agony of obesity and its related complications. Many factors were found to be responsible for the obesity epidemic in children such as increased consumption of energy-dense food, decreasing physical activity and the increasingly easy accessibility of food which can be controlled to prevent it.

Worldwide the urbanization of society is reducing children's physical activity opportunities (Hossain et al, 2007). Increasing calorie consumption is not matched by increased levels of physical activity in children; in fact, children globally are becoming more physically inactive (Anderson and Butcher, 2006). Schools are offering a less healthy environment where children are not protected from bad diets and not encouraged for physically active lifestyles (Story et al, 2006). Because of the link between food advertising and childhood obesity authorities in the UK have banned the advertising of high fat, salt and sugar products in or around programmes made for children ([www.ofcom.org.uk/consult/condocs/foodads](http://www.ofcom.org.uk/consult/condocs/foodads)).

The growing epidemic of childhood overweight and obesity is a major public health concern. Currently 15% of US youth are overweight, a prevalence nearly twice as high in children and three times as high in adolescents compared to 1980 prevalence rates (US Department of Health and Human Services, 2001). Almost two-thirds (60%) of overweight children have at least one cardiovascular risk factor (e.g., hypertension, hyperlipidemia) and the prevalence of type 2 diabetes mellitus is increasing in youth (American Diabetes

Association,2000).These trends may seriously compromise the future health and productivity of the US population and add to health care costs.

A total 2048 children participated in the epidemiological survey. Out of these 2048 children, 1017 were from urban population and 1031 were from rural population.

According to Body Mass Index in the study, 66.26% subjects were found to be malnourished in the total population. The degree of obesity had been observed as Pre-obese 3.81% and Obese-I 3.81% and Obese-II were 0.88%. (Table-I Graph-I).

According to BMI-For-Age Percentile, overall, 6.20% of children and adolescents were at risk of overweight and 4.39% were overweight during health survey while 24.80% are underweight.

Overweight children were highest among urban (6.39%) than rural (2.42%). (Table-II Graph-II).

Obesity and overweight result from an imbalance in the energy expenditure equation such that either food intake is in excess; energy expenditure is deficit or both. Obesity presently has been properly documented as an ailment in its own right which is preventable through changes in life style especially diet. Obesity is a major determinant of many non-communicable diseases and induces diabetes mellitus and coronary heart disease.

**Table 1a- Prevalence of overweight and obesity according to BMI Categories**

Group/Sub Group	Below 18.5		18.5-22.9		23.0-24.9		25.0-29.9		Above 30	
	N	%	N	%	N	%	N	%	N	%
<b>All Data</b>	1357	66.26	517	25.24	78	3.81	78	3.81	18	0.88

**Table 2a. Prevalence of overweight and obesity according to BMI-For-Age Percentile Categories**

Group/Sub Group	Underweight		Healthy Weight		Risk Overweight		Overweight	
	N	%	N	%	N	%	N	%
<b>All Data</b>	508	24.80	1323	64.60	127	6.20	90	4.39

Physiologically, the body weight is constantly changing. Obesity is the result of excessive expansion of adipose tissue mass. This disorder is often accompanied by abnormalities in systemic carbohydrate and lipid metabolism and the secretion and action of insulin alterations thought to reflect diabetogenic effect of obesity. There remain questions about the best diagnostic criteria for this age group.

According to Dietz and Bellizzi, 1999, overweight, during childhood and adolescence, is characterized by a body mass index for age (BMI/age) above the 85th percentile and obesity by a BMI/age above the 95th percentile, based on the National Health and Nutrition Examination Survey (NHANES II-III) standards. Other authors say it is possible to adopt absolute BMI limits for specific age groups, during the growth period, without considerably compromising diagnostic specificity or sensitivity (Monteiro et al, 1999). An expert committee in 1997 declared BMI for age to be the principal measure of obesity in childhood. The adoption of regional BMI standards has also been proposed (Anjos et al, 1998).

Diagnostic criteria should be simple, low cost, reproducible and reliable and offer elevated sensitivity and specificity, thus reducing false positive and negative diagnoses. Based on a pool of studies of BMI profiles by age in a number of different countries, including Brazil, proposed limits for overweight and obesity to be applied to the 2 to 20 age group internationally (Cole et al., 2000).

In this study, the prevalence of obesity was more in boys than in girls. It is possible that the Western feminine self-image and fear of obesity were more marked in girls than in boys. Girls as young as 5 years of age care more about their self-image and associate it to weight status. The same has been reported in other studies where boys had higher prevalence of obesity than girls at all ages (Brown et al, 2002 and Huang et al, 2003).

WHO in 1996 stated that BMI values are not dependent on age or gender of human. But, they may not narrate equally to levels of obesity in diverse populations (certain racial groups in Africa and Asia) as a result of differing body-proportions. Also, the BMI index may not accurately reflect the fatness levels of people with builds falling far outside the average. As mentioned earlier, active persons like sportsmen and sportswomen, will tend to have BMI levels of "overweight" or even 'obese', due to their higher levels of muscle mass. BMI categories also fail to account for varying proportions of fat, bone, cartilage, water weight and more.

Centers for Disease Control and Prevention (CDC), 2000 stated that "overweight" is defined differently for children and teenagers than for adults. Due to growth and differing maturation rates, a modified BMI compares heights and weights against growth charts which also consider gender. This index is known as the BMI-For-Age Percentile (for children 2-19 years of age), and reflects BMI as compared with boys and girls of the same age. Because of that increase, the American Diabetes Association and

the American Academy of Pediatrics recommend routine screening of children at higher risk for Type-2 diabetes. It is essential to identify the best anthropometric index in any population to predict chronic disease risk.

In the present study, according to BMI-For-Age Percentile, 24.80% children were underweight in of 10-19 age groups. In the present study 6.20% of children and adolescents were at risk of overweight and 4.39% were overweight during health survey. The results of the present study also shows that it is appropriate to choose BMI / age, based on the international standards, as a diagnostic indicator for overweight and obese children.

Worldwide one in 10 children, aged 5-17 year old, is overweight, a total of 155 million, of which around 30-45 million are obese (Lobstein et al, 2004). The prevalence rate of overweight in children under 5 years of age in developing countries is 3.3% or 17.6 million (Executive Board, 2001). Countries with the highest prevalence of overweight are located mainly in the Middle East, North Africa, and Latin America (de Onis et al, 2000). In the United States, the prevalence of childhood overweight tripled between 1980 and 2000 (Ogden et al, 2002).

In Australia, the prevalence of childhood overweight almost doubled between 1985 and 1995 (Magarey et al, 2001). In Europe there are 14 million children who are overweight and 3 million of them are obese. In 2002 nearly a quarter of children within the European Union were overweight, higher than the predicted peak for 2010. (Lobstein et al, 2004). In India prevalence of overweight/obesity was: Delhi(North) Overweight, 22%; obesity 6%.(Sharma et al, 2007);

Uttar Pradesh (North) Obesity, 7.6% (Gupta et al, 1999). In the present study, as per the BMI for age percentile, 6.20% of children were at risk of overweight and 4.39% were overweight which less than the other studies. But the necessary steps must be taken at this stage to prevent the agony of obesity and its related complications. Many factors were found to be responsible for the obesity epidemic in children such as increased consumption of energy-dense food, decreasing physical activity and the increasingly easy accessibility of food which can be controlled to prevent it. Overweight and obese children can develop obesity in adulthood and can develop diabetes, hypertension and cardiovascular diseases at an early age (Krushnapriya et al 2015).

Worldwide the urbanization of society is reducing children's physical activity opportunities (Hossain et al, 2007). Obesity in childhood is the most challenging public health issue in the twenty-first century. It has emerged as a pandemic health problem worldwide (Palanikumar and Sunil , 2021) Data from 2015 demonstrates that approximately 108 million children and 604 million adults globally had a BMI of 30 to classify them as obese; signifies an increase in the prevalence of obesity in almost all countries since 1980 and a doubling in prevalence in 70 countries during that period (NCD Risk Factor Collaboration, 2016).



Increasing calorie consumption is not matched by increased levels of physical activity in children; in fact, children globally are becoming more physically inactive (Anderson and Butcher, 2006). Schools are offering a less healthy environment where children are not protected from bad diets and not encouraged for physically active lifestyles (Story et al, 2006). Because of the link between food advertising and childhood obesity authorities in the UK have banned the advertising of high fat, salt and sugar products in or around programmes made for children ([www.ofcom.org.uk/consult/condocs/foodads](http://www.ofcom.org.uk/consult/condocs/foodads)).

The growing epidemic of childhood overweight and obesity is a major public health concern. Currently 15% of US youth are overweight, a prevalence nearly twice as high in children and three times as high in adolescents compared to 1980 prevalence rates (US Department of Health and Human Services, 2001). Almost two-thirds (60%) of overweight children have at least one cardiovascular risk factor (e.g., hypertension, hyperlipidemia) and the prevalence of type 2 diabetes mellitus is increasing in youth (American Diabetes Association, 2000). These trends may seriously compromise the future health and productivity of the US population and add to health care costs. A multidisciplinary approach to promote dietary and physical activity changes in the entire family should be used for the treatment and prevention of overweight and obesity in early childhood (Karel et al 2020).

## CONCLUSION

BMI categories also fail to account for varying proportions of fat, bone, cartilage, water weight and more. BMI index may not accurately reflect the fatness levels of people with builds falling far outside the average. The boys and girls of the same age have differing growth and maturation rates. Thus a modified BMI compares heights and weights against growth charts which also consider gender specific. For children 2-19 years of age, BMI-for-age percentile is the measure and it reflects BMI as compared with same aged boys and girls. Hence it is essential to identify the best anthropometric index in any population to predict chronic disease risk.

## ACKNOWLEDGEMENTS

The author is highly thankful to UGC, New Delhi for providing grant for undertaking the research project.

**Conflict of Interests:** I declare no conflicts of interest to disclose. This manuscript has not been submitted to, nor is under review at, another journal. The authors have no affiliation with any organization with a direct or indirect financial interest in the subject matter discussed in the manuscript.

## REFERENCES

- American Diabetes Association Type 2 diabetes in children and adolescents. *Pediatrics and adolescents*. 2000; 105: 671–680.
- American Diabetes Association. Type 2 diabetes in

- children and adolescents. *Diabetes Care* 2000; 23: 381–389.
- Anderson PM, Butcher KE. Childhood obesity: trends and potential causes. *Future Child*. 2006; 16(1):19-45.
- Anjos LA, Veiga, GV, Castro IRR. Distribuição dos valores do índice de massa corporal da população brasileira até 20 anos. *Rev Panamer Saude Publica*. 1998; 3:42-50.
- Brown M, Sibille K, Phelps L, McFarlane K. Obesity in children and adolescents. *Clinics in family practice* 2002; 4:1–9.
- Center for Disease Control (CDC). Body mass Index: BMI for children and teens. 2000. [apps.nccd.cdc.gov/dnpabmi](http://apps.nccd.cdc.gov/dnpabmi).
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000; 320:1240-3.
- De Onis M, Blossner M. Prevalence and trends of overweight among preschool children in developing countries. *Am J Clin Nutr*. 2000; 72 :1032-9.
- Dietz WH, Bellizzi MC. Introduction: the use of body mass index to assess obesity in children. *Am J Clin Nutr*. 1999; 70(1):1235-55.
- Executive Board Eb109/11 109th Session 11 Provisional agenda item 3.8. Report by the Secretariat. Childhood nutrition and progress in implementing the International Code of Marketing of Breast-milk Substitutes. World Health Organization. Geneva, 2001.
- Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics*. 1999; 103: 1175–1182.
- Gupta Prabha S. Nutrition: an Indian prospective. *Ind. Jnl. of Public Health*, 1999; XXXXIII (1): 11-16.
- Gupta RK. Nutrition and the Diseases of Lifestyle. In: Bhalwar RJ, editor. *Text Book of Public health and Community Medicine*. 1st ed. Pune: Department of community medicine AFMC, New Delhi: Pune in Collaboration with WHO India Office; 2009. p. 1199.
- Gardner K and Bird J Prevalence of overweight, obesity and underweight among 5-year-old children in Saint Lucia by three methods of classification and a comparison with historical rates. 2011. *Child Care Health and Development* 37(1):143-9 DOI:10.1111/j.1365-2214.2010.01154.x
- Hossain P, Kawar B, El Nahas M. Obesity and diabetes in the developing world—a growing challenge. *N Engl J Med*. 2007;356(3):213-5.

- Huang YC, Wu JY, Yang MJ. Weight-for-height reference and the prevalence of obesity for school children and adolescents in Taiwan and Fuchien areas. *J Chin Med Assoc* 2003;66:599–606.
- Krushnapriya Sahoo, Bishnupriya Sahoo, Ashok Kumar Choudhury, Nighat Yasin Sofi, Raman Kumar, and Ajeet Singh Bhadoria Childhood obesity: causes and consequences *J Family Med Prim Care*. 2015 Apr-Jun; 4(2): 187–192.
- Lobstein, T, Baur, L, Uauy R. IASO International Obesity Task Force. Obesity in children and young people: a crisis in public health *Obesity Reviews* 2004; 5:4-104
- Magarey AM, Daniels LA, Boulton TJ. Prevalence of overweight and obesity in Australian children and adolescents: reassessment of 1985 and 1995 data against new standard international definitions. *Med J Aust*. 2001; 174:561–564.
- Monteiro POA, Victora CG, Barros FC, Tomasi E. Diagnóstico de sobrepeso em adolescentes: estudo de diferentes critérios para o índice de massa corporal. *Rev Saude Publica*. 1999; 34:506-13.
- Ogden CL, Kuczmarski RJ, Flegal KM, Mei Z, Guo S, Wei R, Grummer-Strawn LM, Curtin LR, Roche AF, Johnson CL. Centers for Disease Control and Prevention 2000 growth charts for the United States: improvements to the 1977 National Center for Health Statistics version. *Pediatrics*. 2002; 109:45– 60.
- Popkin BM, Doak CM. The obesity epidemic is a worldwide phenomenon. *Nutr Rev*. 1998; 56:106–14.
- Sharma A, Sharma K, Mathur KP. Growth pattern and prevalence of obesity in affluent schoolchildren of Delhi. *Public Health Nutr*. 2007;10:485-91.
- Stevens J, Cai J, Pamuk ER, Williamson DF, Thun MJ, Wood JL. The effect of age on the association between body mass index and mortality. *N Engl J Med* 1998; 338: 1
- Story M, Kaphingst KM, French S. The role of schools in obesity prevention. *Future Child*. 2006 Spring; 16(1):109-42.
- U.S. Department of Health and Human Services: Diabetes in the United States: A strategy for prevention. U.S. department of health and human services, Public Health Service Centres for disease control and National Center for chronic disease prevention and Health promotion. Division of Diabetes Translation, 1993.
- US Department of Health and Human Services . The Surgeon General's call to action to prevent and decrease overweight and obesity. Rockville, MD, US Department of Health and Human Services, Public Health Service, Office of the Surgeon General; 2001.
- World Health Organization Expert Committee on Physical Status.(The World Health Organization Expert Committee on Physical Status. The Use and Interpretation of Anthropometry. Physical Status: Report of a WHO Expert Committee: WHO Technical Report Series 854, WHO, Geneva, 1996).
- [www.ofcom.org.uk/consult/condocs/foodads](http://www.ofcom.org.uk/consult/condocs/foodads)

# Adsorption of Phenol And Resorcinol on Parthenium Based Activated Carbon (PAC) in Basal Salt Medium: Equilibrium and Kinetics

Khushi Gandhi<sup>1</sup>, Shambhatee Srivastav<sup>1</sup>, Huda Afreen<sup>\*2</sup>, Ravi Kant Singh<sup>\*1</sup>

<sup>1</sup>Amity Institute of Biotechnology, Amity University, Noida, U.P., India.

<sup>2</sup>Department of Biotechnology, Dr. D. Y. Patil Biotechnology and Bioinformatics Institute Dr. D. Y. Patil Vidyapeeth, Pune, Maharashtra, India.

## ABSTRACT

The adsorption studies were carried out for the removal of Phenol and Resorcinol on *Parthenium* Based Activated Carbon (PAC) at 301 K and neutral pH ~7. To determine the adsorption equilibrium isotherms, the experiments were performed at the initial concentration of adsorbates with adsorbent doses from 1 g/l to 10 g/l. The experimental data were fitted to six isothermal models, and a nonlinear least square method has been used to estimate model parameters. It has been observed that for adsorption on PAC, Freundlich model was best fitted for the study and data was more closely related for the removal of both the pollutants from industrial wastewater. In this study, the adsorption kinetics were also studied in three distinct phases-rapid, medium & amp: slow for the time period till 12 hours. The parameter values found in this study shall be useful in designing and activated carbon adsorber for treatment of industrial wastewater at large scale. The present work shows that *Parthenium* based activated carbon (PAC) was an effective method for the adsorption of phenol and resorcinol and p-cresol from aqueous solution. The adsorption of phenol and its derivatives are in the following order: Phenol is adsorbed in maximum amount Resorcinol is comparatively adsorbed lesser than phenol.

**KEY WORDS:** PHENOL, RESORCINOL, ACTIVATED CARBON, ADSORPTION EQUILIBRIUM ISOTHERM, ADSORPTION KINETICS, ADSORPTION STUDIES.

## INTRODUCTION

Adsorption is considered as one of the most preferred and cost-effective methods for treatment of waste water contaminant constituents such Phenol and resorcinol the pollution of industrial wastewater, which can reduce the concentration of pollutants discharged into the plant (Agency for Toxic Substances and Disease Registry (ATSDR), 1990; Daifullah and Girgis, 1998; Mourão, Carrott and Ribeiro Carrott, 2006). Naturally, this will inevitably lead to a decline in water quality, especially causing scarcity of drinking water. These organic compounds in water pollution are mainly caused by heavy industrialization (Tanada et al., 1990; Wu and Tseng, 2006) most scientific research is moving towards an innovative process of implementing environmentally friendly treatments. These organic pollutants are non-biodegradable or poorly biodegradable. Like most pollutants, phenol is harmful, and it has been found in the wastewater of many industrial units (D'Ari

and Barker, 1985; Rathi and Puranik, 2002, Mourão and Ribeiro Carrott, 2006).

Phenol has an unpleasant odor and taste and is toxic to aquatic organisms, plants, and humans at concentrations of 5 µg/L-1 Long-term ingestion of phenol at concentrations between 10 and 240 mg L-1 can cause oral irritation, blurred vision, and diarrhea (World Health Organization (WHO), 1971; Fawell and Hunt, 1988). Therefore, this pollutant often appears in rivers, seawater, industrial waste, urban wastewater, and even groundwater. Among all the methods of treatment of wastewater loaded with organic compounds, the adsorption of phenol on solid supports such as activated carbon, clay, and transition metal oxides allows their removal from water with a high removal rate without the addition of chemicals.

*Parthenium hysterophorus* (Linn), sometimes known as carrot grass, is an invasive species that was introduced to India through a planned import from the United States (PAC). PAC is made by chemical activation using concentrated H<sub>2</sub>SO<sub>4</sub>. The produced activated carbon was examined and determined to be a useful adsorbent (Singh et al., 2008). This

**Article Information:**\*Corresponding Author: [rksingh1@amity.edu](mailto:rksingh1@amity.edu), [huda](mailto:huda).

Received 15/04/2023 Accepted after revision 23/06/2023

Published: June 2023 Pp- 84-90

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <http://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.4>



paper highlights the use of parthenium activated carbon's adsorption characteristics for the elimination of pollutants like phenol and others. The work being presented aims to investigate the adsorption of phenol and resorcinol on activated carbon made of parthenium.

## MATERIAL AND METHODS

**Preparation of adsorbent:** Activated carbon act as an effective adsorbent material because of its highly porous nature and provides a large surface area for adsorption (Kadirvelu, Sivasankari, et al., 2002; Singh et al., 2008). There are two preferred ways for the preparation of activated carbon first one is called as physical method and a second method called as chemical method. During the preparation of activated carbon there are several parameters which is required and needed to keep in mind such as temperature and heat duration and pressure. Dried activated carbon is pulverized and sieved to get the particles then the next step for the process of preparation of activated carbon is carbonization also called as pyrolysis which is carried out under heat treatment in the absence of air, Dried activated carbon is then pulverized and sieved to get the particles This leads to the initialization of porosity formation in activated carbons (Ajmal et al., 2006; Singh et al., 2008).

**Characterisation of Activated carbon:** A high porosity and this microporosity enable great adsorption and selectivity, making activated carbon the most effective adsorbate. Both PAC and GAC forms of activated carbon are offered. Basal zone, heterogeneous groups (mostly oxygen-containing surface groups), and inorganic ash are the typical three zones that make up the AC surface (Kumar et al 2003). Adsorption is only permitted in the mesoporous region when the adsorbent possesses liquid

adsorbate, leading to a greater parthenium activated carbon concentration in the ash. The basal zone contains the majority of the adsorption sites for aromatic chemicals (Li et al., 2002). The chemical properties of the carbon surface are defined by heterogeneous groups, who are more active (Kadirvelu, Sivasankari, et al., 2002). Physical, chemical, and electrochemical processes can change the nature of the surface groups (Pandey & Singh, 2014).

**Adsorbate solution:** For the preparation of synthetic adsorbate solution of phenol and resorcinol. Phenol and Resorcinol were used each with known concentration of 500 mg/l. And for the stock solution preparation and storage of (1% w/v) for both 10g of phenol and resorcinol were dissolved in 90 ml of water separately to make up the volume to 100 ml each. This solution is stored in cool and well- ventilated area and same way phenol is stored. For the experiment, the solution of desired concentrations was prepared by successive dilution of the stock.

**Determination of phenol and resorcinol in a solution :**The determination of phenol and resorcinol in a solution was done with the help of UV-vis spectrophotometer onto carbonaceous adsorbent at maximum wavelength of 217nm for phenol adsorption and 273nm wavelength for resorcinol adsorption (model Lambda 35; Perkin Elmer, Massachusetts 02451, U.S.A) (Kumar, Kumar and Kumar, 2003; Ajmal et al., 2006).

**Experiment procedure:**The experimental studies were carried out to observe the adsorption capacity. The process was carried out as follows: stems of parthenium plants were collected from fellow lands near Amity University Noida campus and then the batch experiment was carried out for the study of adsorption potential of *Parthenium* activated carbon.

**Table 1. Major Characteristics of the PAC**

S. No.	Properties	Parthenium based activated carbon PAC
1	BET surface area, m <sup>2</sup> /g	350
2	pH (1% solution)	6.8
3	Moisture content (%)	2.6
4	Ash content (%)	7.2
5	Bulk density (g/ml)	0.62
6	Average particle size (mm)	0.512
7	Average pore diameter (Ao)	36.74
8	Water soluble matter (%)	3.2
9	Acid soluble matter (%)	3.8

In this adsorption experiment, 10g of resorcinol aqueous solution sample (whose initial concentration and pH were known) was poured to a known adsorbent dose of 100ml in a 500ml stoppered conical flask and shaken a bit then for a constant time period of 24 hr and at varied temperature adsorption was studied with the help of different isotherm models, where adsorption kinetics were studied by keeping

the temperature constant at 301K, and according to the residual concentration of resorcinol and phenol their percent removal were calculated (Kumar et al 2003; Singh et al., 2008).

The waste effluents concentration varies at high range from 1gm to 1000 gm/l and sometimes more, so the experimental

studies the concentration resorcinol taken was 500 gm/l. The adsorbent dose concentrations in each 500 ml flask used ranges from 1gm then 2gm then 3gm and so on to 10 gm at an initial temperature of 301K with initially maintained at neutral pH 7. The samples were taken at some specific time intervals and filtered this study were carried out on Parthenium activated carbon. The quantity of adsorped phenol and resorcinol by 1 gram of adsorbent,  $q_t$  (mg/g) at any given time (t) was calculated by (Pandey and Singh, 2014).

$$q_t = [(C_o - C_t) V] / m \quad (1)$$

where  $C_o$  is the concentration of phenol and resorcinol at the initial time while  $C_t$  is the concentration at time t, respectively, V is the volume of adsorbate solution, and m is the mass of adsorbent dose used in the run. At equilibrium condition, the amount of adsorption per gm of adsorbent,  $q_e$  (mg/g), was calculated by:

$$q_e = [(C_o - C_e) V] / m \quad (2)$$

**Adsorption equilibrium isotherms:** For the study of adsorption different equilibrium isotherms are used. This equilibrium is generally expressed in terms of concentration as of adsorbate ( $C_e$ ) which is industrial wastewater in liquid state and adsorbent ( $q_e$ ) on which the process of adsorption occurs at equilibrium conditions which are keeping temperature and pressure constant. These isotherm models help us to study about interaction between adsorbate and adsorbent. In orders to correlate the data six different isotherm models are used which are: Freundlich isotherm model, Langmuir isotherm model, Radke isotherm model, Toth isotherm model, Redlich isotherm models and Fritz isotherm models (Abuzaid & Nakhla, 1996, Li et al., 2001; Nouri, Haghsresht and Lu, 2002).

The Langmuir model equation is expressed as:

$$q_e = (Q_0 b C_e) / (1 + b C_e) \quad (3)$$

here b is the equilibrium adsorption constant. It is a measurement of adsorption affinity and is associated with the adsorption free energy. The quantity of phenol or resorcinol required to completely create a monolayer on the surface of a gram of activated carbon is known as  $Q_0$ , which represents the adsorption capacity (Abuzaid and Nakhla, 1996). This model assumes that all of the activated carbon's surface has the same level of activity. Moreover, it is constrained to a particular concentration range.

The Freundlich, an empirical model, can be expressed in the equational form as:

$$q_e = K_F C_e^{1/n} \quad (4)$$

$K_F$  is the measure of the adsorption capacity and n is a measure of the adsorption intensity This model is applicable to the system involving heterogeneous surface and organic compound (e.g. activated carbon—phenolic compound). The Radke–Prausnitz, Toth, and Fritz–Schlunder models

are not common in use to describe adsorption of phenolic compounds on activated carbon.

**Kinetic study of adsorption:** Following the rate of adsorption and ultimate equilibrium conditions, the fluctuation in the concentration of unadsorbed or remaining phenol and resorcinol in the solution is almost minimal. Pseudo-first order and pseudo-second order kinetic models are the most used ones for phenolic chemical adsorption on activated carbon (Moreno-Castilla et al., 1995; Abuzaid and Nakhla, 1996). The ability of these two categories of kinetic models to predict the adsorption of phenol and resorcinol on PAC is investigated in this work.

As written, the pseudo-first order kinetic model is:

$$dq_t/dt = k_1(q_e - q_t) \quad (5)$$

where  $k_1$  is pseudo-first order adsorption rate constant ( $\text{min}^{-1}$ ). The model with pseudo-second order kinetics can be expressed as:

$$dq_t/dt = k_2(q_e - q_t)^2 \quad (6)$$

where  $k_2$  is the pseudo-second order adsorption rate constant ( $\text{g mg}^{-1} \text{min}^{-1}$ ). When it is integrated with boundary condition,  $q_t(t=0) = 0$  becomes:

$$\ln(q_e - q_t) = \ln q_e - k_1 t \quad (7)$$

Here,  $q_e$  and  $q_t$  represents the amounts of phenolic compounds studied which are phenol and resorcinol adsorbed per unit mass of adsorbent at equilibrium and at time t, respectively, and can be calculated by using the above equations. Based on the above equations order of kinetics is calculated which shows which order of reaction is followed by the studied interactions of adsorption (Moreno-Castilla et al., 1995; Abuzaid and Nakhla, 1996).

## RESULTS AND DISCUSSION

**Effect of adsorbent's dosage:** In order to know the effect of dose of adsorbent on the adsorption of resorcinol and phenol a number of adsorption experiments were carried out with varying amounts of PAC (1–10 g/l) were contacted with *phenol* and *resorcinol* Solutions. The analysis of this experimental series demonstrates that a greater adsorbent dosage increases the amount of available active sites for adsorption, leading to an increase in removal efficiency (Lata, Garg and Gupta, 2007). The % elimination changes very little incrementally as the dosage of PAC is increased further. Adsorbent dosage is not very effective after 99% removal, and the majority of active sites are left vacant. This causes a decrease in the amount of phenol and resorcinol adsorption decreases with increase in adsorbent dose. This research approximates the ideal adsorbent dosage needed to remove phenol and resorcinol under current operating circumstances when employing activated carbons (Kumar, Kumar and Kumar, 2003).

It takes 10 g/l of PAC dosage. The ideal dose of adsorbent for PAC is then required, which is 10 mg/l, to treat 500 mg/l

of phenol and resorcinol solution since the removal rate is so low. The final percentage of removal is represented in the below mentioned data the final removal percentage of phenol is estimated as 93.6% by 10g/l PAC dose, and the final percentage removal of resorcinol is estimated as 87.2% by 10g/l PAC dose (Pandey and Singh, 2014).

Here,  $q_e$  and  $q_t$  represents the amounts of phenolic compounds studied which are phenol and resorcinol adsorbed per unit mass of adsorbent at equilibrium and at time  $t$ , respectively, and can be calculated by using the above equations. Based on the above equations order of kinetics is calculated which shows which order of reaction is followed by the studied interactions of adsorption (Moreno-Castilla et al., 1995; Abuzaid and Nakhla, 1996). and Kumar, 2003). It takes 10 g/l of PAC dosage. The ideal dose of adsorbent for PAC is then required, which is 10 mg/l, to treat 500 mg/l of phenol and resorcinol solution since the removal rate is so low. The final percentage of removal is represented in the below mentioned data the final removal percentage of phenol is estimated as 93.6% by 10g/l PAC dose, and the final percentage removal of resorcinol is estimated as 87.2% by 10g/l PAC dose (Pandey and Singh, 2014).

**Table 2. Deviation Percentage for Phenol and Resorcinol in different isotherm models**

% Deviation ( $R^2$ )		
Isotherm Models	Phenol	Resorcinol
Freundlich $q_e = KFC^{1/n}$	13.5	11.64
Langmuir $q_e = (q_0 bCe)/(1+bCe)$	50.75	11.09
Redlich–Peterson $q_e = (K1Ce)/(1+K2C^be)$	13.45	11.67
Toth $q_e = (q^{\infty}eCe)/[a+Cne]^{1/n}$	14.17	11.92
Radke–Prausnitz $1/q_e = 1/KC_e + 1/KC_e^{1/n}$	14.93	9.27
Fritz–Schlunder $q_e = \alpha 1C_e^{\beta} 1/1 + \alpha_2 C_e^{\beta_2}$	14.45	11.62

**Isotherm studies:** By creating a dynamic phase equilibrium between the solid surface of the adsorbent and the liquid adsorbate, equilibrium is studied. This equilibrium is often stated in terms of the adsorbate's concentration and the amount of adsorbate that has been put onto the adsorbent. Adsorbate loading on the adsorbent is shown as ( $q_e$ ) in adsorption equilibrium isotherms, and the liquid phase concentration of adsorbate is shown as ( $C_e$ ) at equilibrium while maintaining temperature constant. Adsorption isotherm, then, depicts how adsorbate and adsorbent interact. Many models used to characterise the experimental isotherm data are empirical equations based on the form of the isotherm curve to correlate the isotherm data of this sort, six isotherm models are selected (Li et al., 2002).

The data mentioned below shows a comparative result of all the isotherm models studies. The data mentioned below shows that Freundlich isotherm model is best suited for Phenol PAC system, p-cresol PAC (Moreno-Castilla et al., 1995) system and Resorcinol PAC system whereas Langmuir isotherm model is least for Phenol PAC system and Resorcinol PAC system. To represent the whole concentration range with a single isotherm model, six isotherm models were tested. The correlation coefficient and greatest variation from the anticipated value served as the selection criteria for the isotherm model (Kadirvelu, Sivasankari, et al., 2002).

On this basis, Langmuir model could not fit the data for phenol with the maximum deviation percentage of 50.75 %. but it suited for resorcinol with the maximum deviation percentage of 11.09 %. The results of the Toth and Radke-Prausnitz are nearly similar. The only four parameter isotherm model studied is by Fritz Schlunder. This model gave the similar result to three parameter models with maximum deviation percentage for phenol of 14.45% and for resorcinol of 11.62%.

**Kinetic study:** Kinetics studies were performed in simple glass flask experiments. The calculated amounts of adsorbate were poured into 500 ml stoppered conical glass flask and activated carbon at a range of 1mg/l to 10mg/l variably was added to each flask. These flasks were kept untouched at 301K. The pre-determined amount of stock solution was added to each flask to obtain desired initial phenol and resorcinol concentration with working solution. The flasks were taken out and filtered one at a time at certain time intervals (Kadirvelu, Sivasankari, et al., 2002). The filtrate was used for phenol analysis. These experiments were done for 24 hours. There was very gradual decrease in concentration after 24 hours keeping the adsorbent dosage at 10 g/l, the temperature at 301 K, and the pH of the solution constant at 7. Using Kinetic equation, the values of ( $q_t$ ) corresponding to each expected concentration of phenol and resorcinol at various time intervals were determined.

According to the experiments carried on each set of trials, each activated carbon and phenolic interaction, which is a phenol-PAC system, followed a different pace of reaction. Pseudo first order kinetics was followed by pseudo second order kinetics and the resorcinol-PAC system. The estimation of time for the maximum removal of phenolic compounds were with the help of studying diffusion coefficient as the diffusion coefficient determines the time it takes a solute to diffuse a given distance in a medium (Pandey and Singh, 2014). The maximum time taken for phenol-PAC system was 4hrs with the diffusion coefficient  $R^2 = 0.975\%$ , whereas for resorcinol-PAC system it estimated to be 10hrs, with diffusion coefficient  $R^2 = 0.9926\%$ .

**Effect of Temperature:** When the temperature rises, phenol and its derivatives' ability to adsorb substances increases. It was well understood that rising temperature would speed up the rate of adsorbate molecule diffusion through the exterior boundary layer and into the pores of the adsorbent particles, resulting in a reduction in the solution's viscosity (Li et al., 2002). The equilibrium capacity of the adsorbate

for a certain adsorbate will also fluctuate with temperature. The following equations were used to determine the thermodynamic parameters such as free energy change (G), entropy (S) and enthalpy change (H):

$$\Delta G^0 = - R T \ln K_c$$

$$\Delta G = \Delta H - (T \Delta S)$$

$$\ln K_c = \Delta H/RT + \Delta S/R$$

$$\Delta G = + Ve$$

i.e.  $\Delta G \propto 1/\text{Temperature}$

$K_c$  = equilibrium constant.

$K_c = C_{as}/C_e$

$C_{as}$  = equilibrium const. (mg/L) of adsorbate on the solution.

$C_e$  = equilibrium const. (mg/L) of adsorbate on the solution.

This relation with free energy changes along with temperature pointed towards the direct proportionality of adsorption capacity with temperature. The positive value of ( $\Delta S$ ) shows the affinity of PAC for phenol and its derivatives (Phenol and Resorcinol). During the adsorption process, some structural changes occur that result in increased randomness at the interface of the solid-solution.

**Effect of pH of the solution:** The surface charge distribution, which results from the interaction of the surface with adsorbate ions, plays a significant role in the interaction between the adsorbent and the adsorbate. This demonstrates how pH has an impact on the adsorbent's surface charge and the amount of ionisation of the adsorbate, which has an impact on the adsorption phenomena. The initial pH of an aqueous solution, which is defined as the pH at which the surface of activated carbon has zero charge and is thus neutral (Moreno-Castilla et al., 1995; Lata, Garg and Gupta, 2007), is one of the parameters that is frequently examined. As the pH rises over its original pH, the surface of the activated carbon becomes negatively charged and the positive charges are diffused, allowing the adsorption to take place.

**Table 3. Effect of Temperature on Phenol, p-Cresol and Resorcinol**

S.No.	Adsorbate	Temperature (0K)	$\Delta G$ (KJ/mole)	$\Delta H$ (KJ/mole)	$\Delta S$ (J.K-1 min-1)	R2
1	Phenol	288	2.42	13.24	40.86	0.98
		298	2.04			
		308	1.68			
		318	0.82			
2	p-Cresol	288	3.18	14.02	30.62	0.97
		298	2.67			
		308	1.82			
		318	1.02			
3	Resorcinol	288	4.06	14.46	22.46	0.99
		298	3.46			
		308	2.72			
		318	1.88			

The surface of the activated carbon becomes negatively charged as the pH drops below the starting pH, and the positive ions in the solution are attracted to the surface via electrostatic attraction. Diffusive is now used to help in the adsorption process. The fluctuation in pH has little impact on the effects of pH on the adsorption of phenol and its derivatives. Adsorption isotherm were studied at different pH values. The observed results do not show significant change in the adsorption process at different pH values (AYRANCI and DUMAN, 2005). At low and high pH, their hydrophilic behaviour increases and makes them more surface-active (pH 4 – 9) studies. Preferred pH ~ 7 neutral due to significance difference in adsorption of pollutant.

**Effect of Particle Size:** The effect of particle size for a const. weight of PAC on the adsorption of Phenol and its

derivatives were carried out. The result reveals that the adsorption is higher on smaller particle size of the adsorbent (Kadirvelu, Senthilkumar, et al., 2002; Rajeshwarisivaraj and Subburam, 2002). The reason for the same is that adsorption is a surface phenomenal, so the greater surface area for adsorption lead to the greater accessibility to pores and hence increased the uptake of smaller particles.

## CONCLUSION

The present work shows that Parthenium based activated carbon (PAC) was an effective method for the adsorption of phenol and resorcinol and p-cresol from aqueous solution. The adsorption of phenol and its derivatives are in the following order. Shows the best adsorption of phenol then resorcinol. The generalised Freundlich model



was demonstrated to provide the best correlation for the adsorption for the adsorption of Phenol, Resorcinol and P-cresol onto PAC. The Langmuir isotherm model was found the least corelative for the adsorption studies on the PAC-phenol and PAC- Resorcinol system. As a result, it was stated that the phenol-PAC system, P-cresol-PAC system, and resorcinol-PAC system may all be employed with the Freundlich isotherm. Since activated carbon is a costly substance, repension is crucial.

In contrast, PAC is chosen since it is less costly and made from waste or harmful materials (Rajeshwarisivaraj and Subburam, 2002). In India, parthenium is a widespread weed that is inexpensive or free to purchase in the countryside. According to the aforementioned findings, it can be said that the PAC has outstanding adsorptive properties and may be successfully used to remove phenol and its derivative from industrial effluent (Kadirvelu, Sivasankari, et al., 2002; Rajeshwarisivaraj and Subburam, 2002; Singh et al., 2008). The adsorption rate forms from these experiments may be helpful in developing techniques for phenol and its derivatives' elimination from wastewater, that are both affordable and long-lasting (Singh, Vats and Tyagi, 2011; Singh et al., 2013).

## ACKNOWLEDGEMENTS

We would like to acknowledge Amity Institute of Biotechnology and Amity University, Noida campus for providing us with facilities and infrastructure for working on this study.

**Conflict of Interests:** The authors declare no conflict of interests.

## REFERENCES

- Abuzaid, N.S. And Nakhla, G.F. (1996) 'Effect Of Solution Ph On The Kinetics Of Phenolics Uptake On Granular Activated Carbon', *Journal Of Hazardous Materials*, 49(2-3), Pp. 217-230. Available At: [https://Doi.Org/10.1016/0304-3894\(96\)01760-8](https://Doi.Org/10.1016/0304-3894(96)01760-8).
- Agenc For Toxic Substances And Disease Registry (Atsdr) (1990) Toxicological Profile For Cresols (Draft). Atlanta.
- Ajmal, M. Et Al. (2006) 'Adsorption Studies On Parthenium Hysterophorous Weed: Removal And Recovery Of Cd(Ii) From Wastewater', *Journal Of Hazardous Materials*, 135(1-3), Pp. 242-248. Available At: <https://Doi.Org/10.1016/J.Jhazmat.2005.11.054>.
- Ayranci, E. And Duman, O. (2005) 'Adsorption Behaviors Of Some Phenolic Compounds Onto High Specific Area Activated Carbon Cloth', *Journal Of Hazardous Materials*, 124(1-3), Pp. 125-132. Available At: <https://Doi.Org/10.1016/J.Jhazmat.2005.04.020>.
- D'ari, L. And Barker, H.A. (1985) 'P-Cresol Formation By Cell-Free Extracts Of Clostridium Difficile', *Archives Of Microbiology*, 143(3), Pp. 311-312. Available At: <https://Doi.Org/10.1007/Bf00411256>.
- Daifullah, A.A.M. And Girgis, B.S. (1998) 'Removal Of Some Substituted Phenols By Activated Carbon Obtained From Agricultural Waste', *Water Research*, 32(4), Pp. 1169-1177. Available At: [https://Doi.Org/10.1016/S0043-1354\(97\)00310-2](https://Doi.Org/10.1016/S0043-1354(97)00310-2).
- Fawell, J.K. And Hunt, S. (1988) *Environmental Toxicology: Organic Pollutants*. New York: Halsted Press, John Wiley & Sons.
- Kadirvelu, K., Senthilkumar, P., Et Al. (2002) 'Activated Carbon Prepared From Biomass As Adsorbent: Elimination Of Ni(Ii) From Aqueous Solution', *Bioresource Technology*, 81(1), Pp. 87-90. Available At: [https://Doi.Org/10.1016/S0960-8524\(01\)00093-1](https://Doi.Org/10.1016/S0960-8524(01)00093-1).
- Kadirvelu, K., Sivasankari, C., Et Al. (2002) 'Activated Carbon From Parthenium As Adsorbent: Adsorption Of Hg (Ii) From Aqueous Solution', *Indian Journal Of Chemical Technology*, 9, Pp. 499-503.
- Kumar, A., Kumar, Shashi And Kumar, Surendra (2003) 'Adsorption Of Resorcinol And Catechol On Granular Activated Carbon: Equilibrium And Kinetics', *Carbon*, 41(15), Pp. 3015-3025. Available At: [https://Doi.Org/10.1016/S0008-6223\(03\)00431-7](https://Doi.Org/10.1016/S0008-6223(03)00431-7).
- Lata, H., Garg, V.K. And Gupta, R.K. (2007) 'Removal Of A Basic Dye From Aqueous Solution By Adsorption Using Parthenium Hysterophorus: An Agricultural Waste', *Dyes And Pigments*, 74(3), Pp. 653-658. Available At: <https://Doi.Org/10.1016/J.Dyepig.2006.04.007>.
- Li, A. Et Al. (2001) 'Adsorption Of Phenolic Compounds On Amberlite Xad-4 And Its Acetylated Derivative Mx-4', *Reactive And Functional Polymers*, 49(3), Pp. 225-233. Available At: [https://Doi.Org/10.1016/S1381-5148\(01\)00080-3](https://Doi.Org/10.1016/S1381-5148(01)00080-3).
- Li, A. Et Al. (2002) 'Adsorption Of Phenolic Compounds From Aqueous Solutions By A Water-Compatible Hypercrosslinked Polymeric Adsorbent', *Chemosphere*, 47(9), Pp. 981-989. Available At: [https://Doi.Org/10.1016/S0045-6535\(01\)00222-3](https://Doi.Org/10.1016/S0045-6535(01)00222-3).
- Moreno-Castilla, C. Et Al. (1995) 'Adsorption Of Some Substituted Phenols On Activated Carbons From A Bituminous Coal', *Carbon*, 33(6), Pp. 845-851. Available At: [https://Doi.Org/10.1016/0008-6223\(94\)00182-Y](https://Doi.Org/10.1016/0008-6223(94)00182-Y).
- Mourão, P.A.M., Carrott, P.J.M. And Ribeiro Carrott, M.M.L. (2006) 'Application Of Different Equations To Adsorption Isotherms Of Phenolic Compounds

- On Activated Carbons Prepared From Cork', Carbon, 44(12), Pp. 2422–2429. Available At: <https://Doi.Org/10.1016/J.Carbon.2006.05.015>.
- Nouri, S., Haghseresht, F. And Lu, G.Q.M. (2002) 'Comparison Of Adsorption Capacity Of P-Cresol & P-Nitrophenol By Activated Carbon In Single And Double Solute', Adsorption, 8(3), Pp. 215–223. Available At: <https://Doi.Org/10.1023/A:1021260501001>.
- Pandey, A. And Singh, R. (2014) 'Industrial Waste Water Treatment By Membrane Bioreactor System', Elixir Chemical Engineering, 70, Pp. 23772–23777.
- Rajeshwarisivaraj And Subburam, V. (2002) 'Activated Parthenium Carbon As An Adsorbent For The Removal Of Dyes And Heavy Metal Ions From Aqueous Solution', Bioresource Technology, 85(2), Pp. 205–206. Available At: [https://Doi.Org/10.1016/S0960-8524\(02\)00089-5](https://Doi.Org/10.1016/S0960-8524(02)00089-5).
- Rathi, A.K.A. And Puranik, S.A. (2002) 'Chemical Industry Wastewater Treatment Using Adsorption', Journal Of Scientific & Industrial Research, 61, Pp. 53–60.
- Singh, R.K., Gupta D.K., Kumar S., Kumar S. (2013) 'Utilization Of Fly Ash As Biologically Activated Carbon On Microbial Consortium For The Removal Of Pollutants From Waste Water', International Journal Of Advanced Biotechnology And Research, 4(1), Pp. 883-891.
- Singh, R.K., Kumar S., Kumar S., Kumar A. (2008) 'Development Of Parthenium Based Activated Carbon And Its Utilization For Adsorptive Removal Of P-Cresol From Aqueous Solution', Journal Of Hazardous Materials, 155(3), Pp. 523–535. Available At: <https://Doi.Org/10.1016/J.Jhazmat.2007.11.117>.
- Singh, R.K., Vats, S. And Tyagi, P. (2011) 'Industrial Wastewater Treatment By Biological Activated Carbon - A Review', Research Journal Of Pharmaceutical, Biological And Chemical Sciences, 2(4), Pp. 1053–1058.
- Tanada, M. Miyoshi T., Nakamura T., Tanda S. (1990) 'Adsorption Removal Of Cresol By Granular Activated Carbon For Medical Wastewater Treatment', Bulletin Of Environmental Contamination And Toxicology, 45(2), Pp. 170–176. Available At: <https://Doi.Org/10.1007/Bf01700179>.
- World Health Organization (Who) (1971) International Standards For Drinking-Water. Geneva.
- Wu, F.-C. And Tseng, R.-L. (2006) 'Preparation Of Highly Porous Carbon From Fir Wood By Koh Etching And Co<sub>2</sub> Gasification For Adsorption Of Dyes And Phenols From Water', Journal Of Colloid And Interface Science, 294(1), Pp. 21–30. Available At: <https://Doi.Org/10.1016/J.Jcis.2005.06.084>.



# Avifaunal Abundance of Lumding Forest Reserve Area, Assam, India

Prof. Jashodeb Arjun\* and Rajesh Paul

<sup>1</sup>Department of Zoology, School of Life Sciences, Royal Global University, Guwahati, Assam, India

## ABSTRACT

The solitary aim of our investigation was to explore the avian species richness in Lumding Forest Reserve area, Assam. The survey was cooperated by the team of Zoological Survey of India. After extended survey, 146 species of birds were spotted from Lumding Forest Reserve area, where birds from 17 different orders (Passeriformes, Columbiformes, Accipitriformes, Piciformes, Bucerotiformes, Gruiformes, Apodiformes, Coraciiformes, Trogoniformes, Cuculiformes, Galliformes, Ciconiiformes, Suliformes, Pelecaniformes, Strigiformes, Anseriformes, Charadriiformes) and 76 families were put down. The bird species thus obtained were compared with the IUCN Red list from where interesting cum astonishing facts were acquired. The IUCN Red list informed us that, 92.51% bird species spotted were under Least Concern category, where as 3.4% were Near Threatened, 2.72% were Vulnerable, 0.68% were Critically Endangered and Endangered. We considered ourselves successful for obtaining such astounded statistics about the avian species plentitude of Lumding Forest Reserve, which will provoke other researchers to come forward and explore more about this place.

**KEY WORDS:** LUMDING FOREST RESERVE AREA, BIRDS, BIODIVERSITY, IUCN.

## INTRODUCTION

Alike other animals, birds too are a remarkable part of ecosystem with amazing ecological values. They cover almost every corner of the planet. Globally there exists around 9000 species of birds of which india contributes 13% with 1300 species (Grimmett et al., 2000). Birds are watcher's delight, with their colorful appreance and melodious chirping, they not only increase the beauty of nature and provide intangible aesthetic enjoyment; they play many roles in ecosytem (Watanuki et al.,2022).

There exists a strong interdependence between birds and other organisms of this universe. These warm blooded vertebrates help in pollination; birds like bulbul, parakeets help in seed dispersal; act as scavengers; helps in pest control in crop fields; manages wetland grazing species like snails, periwinkles; acts as proficient gardeners(Beal et al., 2021) .

Prior to their sensitiveness to habitat change, birds are considered as one of the most important indicators of environmental change (Pitera et al., 2021). They play significant role in controlling insect outbreak. Birds like

Barn Swallow helps in mosquito control by eating around 850 mosquitoes each day (Vinod et al., 2023).

Lumding is a railway town and a municipal board of Nagaon district. It is the regional divisional headquarter of North –Eastern Frontier Railway, a big important junction and famous for Barak Valley Express, Cachar Express, Hill Queen Express, Agartala-Lumding Express, etc., which passes through scenic routes (Pawar and Salunkhe, 2014). It is also famous for cultural and educational excellence.

Initiated in 1964, The International Union for Conservation of Nature's (IUCN) Red List of Threatened Species has become the most reliable source regarding conservation of floral and faunal species of the earth (Miskelly et al., 2019). It accords information regarding the population, habitat, threats and conservation initiatives that assists us to take necessary steps towards protection. The Red List is an indicator of the exact position of worlds biodiversity (Sauve et al., 2021). It helps to protect natural resources on which we are dependent for survival. Hence, during our investigation, we checked our findings in the Red List to assure their future existence and safety.

## MATERIAL AND METHODS

Lumding is a hilly landscape which is covered with dense reserve forest from all sides. Summers are burning here as

**Article Information:**\*Corresponding Author: [jashodeb@gmail.com](mailto:jashodeb@gmail.com)

Received 15/04/2023 Accepted after revision 25/06/2023

Published: June 2023 Pp- 91-99

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.5>


well as winters can be chilling. Monsoon covers rainfall around 60-125mm. The only hill station of Assam i.e. Haflong is very closer to Lumding. Climate remaining favourable for agricultural practices (Arya and Rao, 2014).

Species density and diversity can be good indicators for measuring abundance of birds in any locality (Javed and Rahmani, 1993). Our survey was carried out from January,2016 to December,2018, where almost all the seasons were covered like Summer (March-June) and Winter (November-March).

For spotting, following methods were preferred, (1) point count method, (2) direct count method & (3) area search method.



**Figure 2: Graphical representation showing order wise abundance of avian fauna in Lumding Forest Reserve area**




Bird counting was performed from 5.00-8.00 am (morning) and 4.00-6.00 pm (evening). Various species of birds were observed during survey period. Observation was performed with the help of field binoculars (10x40). Pictures were captured with Digital Canon EOS 1300D, 18MP DSLR. Renowned Ornithologist Dr. Ghosh and his team from Zoological Survey of India, remained physically present and helped in species identification. Species identification was confirmed with the help of “Books of Birds of the Indian subcontinent” (Grimmett et al., 2000) and book of Salim Ali “The Book of Indian Birds”, Wikipedia and other resources.

## RESULT AND DISCUSSION

North-East India refers to the easternmost region of India consisting of contiguous seven sister states, Sikkim and parts of North Bengal (district of Darjeeling , Jalpaiguri and Kochbihar) (Javed and Rahmani, 1993). North East India is ethnically distinct from the other states of India, linguistically the region is distinguished by preponderance of Tibeto- Burman languages. Strong ethnic cultures that had escaped sanskritization effects permeate the region. The eight states from a special category is officially recognized (Peralta et al., 2020).

**Figure 3: Graphical representation showing IUCN status of avian fauna spotted in Lumding Forest Reserve area**



**Figure 4: Pictures of some birds spotted from Lumding Forest Reserve area**



**Figure 5: Pictures of some birds spotted from Lumding Forest Reserve area**



Lumding Reserve Forest is a compact large block of forest, measuring 252.9 sq. Kms and comparatively has less human disturbance. This forest gives shelter to a large variety of floral as well as faunal species (Arya and Rao, 2014). As the forest area of neighbouring Karbi Anglong district and N.C.Hills district are affected by shifting cultivation, the herbs of Asiatic Elephant takes shelter in Lumding Reserve Forest (Joshi and Shrivastava, 2012). The principal type of forest found in Lumding is degraded moist mixed deciduous forest type (Khan, Rao and Wani, 2012).

Lumding Reserve Forest is also habitat for a huge variety of medicinal plants like Sarpagandha, Basaka, Ahoi, Hilikha, etc. Among all the floral species Teak and Gamari are found to dominate in Lumding Reserve Forest (Javed and Rahmani, 1993). Among the climbers *Mikennia sp.* predominates where canopy is open and other climbers are *Bridelia sp.*, *Butea parviflora*, *discorea sp.*, *Clematis sp.* Himalayan black bear, Clouded Leopard, Crab eating mongoose, Slow loris, Assamese Macaque, Hoolock gibbon, Asiatic jackal, etc. comes under faunal diversity of Lumding. 146 species of birds were spotted from Lumding Forest Reserve area during our survey; Of which, 17 different avian orders including Passeriformes, Columbiformes, Accipitriformes, Piciformes, Bucerotiformes, Gruiformes, Apodiformes, Coraciiformes, Trogoniformes, Cuculiformes, Galliformes, Ciconiiformes, Suliformes, Pelecaniformes, Strigiformes, Anseriformes, Charadriiformes were detected.

**Figure 1. Checklist of avian fauna of Lumding Forest Reserve area**

SL NO.	ORDER	FAMILY	COMMON NAME	SCIENTIFIC NAME	Feeding habits*	IUCN Red List (2019-22) status**
1	Passeriformes	Pycnonotidae	Red vented bulbul	<i>Pycnonotus cafer</i>	Nectivorous, Frugivorous,	Least Concern
Insectivorous, Palynivorous						
2		Campephagidae	Scarlet minivet <i>speciosus</i>	<i>Pericrocotus</i> Carnivorous	Insectivorous, Concern Ophiophagous	Least
3		Oriolidae	Black headed oriole	<i>Oriolus larvatus</i>	Frugivorous, Insectivorous	Least Concern
4			Indian Golden oriole	<i>Oriolus kundoo</i>	Frugivorous, Insectivorous, Nectivorous	Least Concern
5		Pittidae	Black crown pitta	<i>Erythropitta ussheri</i>	Carnivorous	Least Concern
6			Indian pitta	<i>Pitta brachyura</i>	Carnivorous, Insectivorous,	
7		Orididae	Maroon oriole	<i>Oriolus trailii</i>	Insectivorous, Nectivorous	Least Concern
8		Alaudidae	Eastern skylark (Oriental skylark)	<i>Alauda gulgula</i>	Granivorous	Least Concern



Table 1a

9	Dicruridae	Racket tailed drongo	<i>Dicrorus paradiseus</i>	Insectivorous, Nectivorous	Least Concern
10		Black drongo	<i>Dicrorus macrocercus</i>	Insectivorous,	Least Concern
11		Ashy drongo	<i>Dicrorus leucophaeus</i>	Insectivorous	Least Concern
12		Bronzed drongo	<i>Dicrurus aeneus</i>	Insectivorous	Least Concern
13	Nectariniidae	Nepal sunbird	<i>Aethopyga nipalensis</i>	Nectivorous	Least Concern
14	Sturnidae	Hill myna	<i>Gracula religiosa</i>	Omnivorous, Insectivorous,	Least Concern
15	Laniidae	White crowned shrike	<i>Eurocephalus anguitimens</i>	Frugivorous Insectivorous, Carnivorous	Least Concern
16		Bull headed shrike	<i>Lanius bucephalus</i>	Insectivorous, Carnivorous	Least Concern
17		Long tailed	<i>Larius collaris</i>	Insectivorous ,	Least Concern
18			fiscal shrike		Carnivorous
18	Leiothrichidae	Jungle babbler	<i>Argya striata</i>	Insectivorous	Least Concern
19	Pycnonotidae	Black headed bulbul	<i>Pycnonotus atriceps</i>	Insectivorous, Frugivorous	Least Concern
20	Hirundinidae	Red rumped swallow	<i>Cecropis daurica</i>	Insectivorous	Least Concern
21	Passeri	Fairy blue bird	<i>Irena puella</i>	Insectivorous, Nectivorous	Least Concern
22	Chloropseidae	Gold fronted leafbird	<i>Chloropsis aurifrons</i>	Omnivorous	Least Concern
23		Orange bellied leafbird	<i>Chloropsis hard wickii</i>	Insectivorous, Nectivorous	Least Concern
24	Pycnonotidae	Brown eared bulbul	<i>Hypsipetes amaurotis</i>	Insectivorous, Frugivorous, Nectivorous	Least Concern
25	Mectarinidae	Large browed wagtail	<i>Motacilla moderaspatisensis</i>	Insectivorous	Least Concern
26	Estrildidae	Spotted munia	<i>Lonchura punctulata</i>	Granivorous	Least Concern
27		Black headed munia	<i>Lonchura malacca</i>	Granivorous	Least Concern
28	Ploceidae	Baya weaver bird	<i>Ploceus philippinus</i>	Granivorous, Insectivorous	Least Concern
29	Dicruridae	Lesser racket tailed drongo	<i>Dicrurus remifer</i>	Insectivorous	Least Concern
30	Muscicapidae	White rumped	<i>Copsychus magpie</i>	Insectivorous , <i>malabaricus</i>	Least Concern Carnivorous
31	Sturnidae	Common myna	<i>Acridotheres tristis</i>	Insectivorous, Granivorous, Carnivorous, Omnivorous	Least Concern
32		White headed starling	<i>Sturnia erythrogygia</i>	Insectivorous	Least Concern
33		Brahminy starling	<i>Sturnus pagodarum</i>	Omnivorous	Least Concern
34		Jungle myna	<i>Acridotheres fuscus</i>	Omnivorous	Least Concern
35		Indian Pied myna	<i>Gracupica contra</i>	Omnivorous	Least Concern
36	Motacilladae	Yellow wagtail	<i>Motacilla flava</i>	Insectivorous	Least Concern
37		Citrine wagtail	<i>Motacilla citriola</i>	Insectivorous	Least Concern
38	Turdidae	Common black bird	<i>Turdus merula</i>	Omnivorous	Least Concern
39	Vangidae	Indian wood shrike	<i>Tephrodornis pondicerianus</i>	Insectivorous	Least Concern

Table 1b

40		Motacillidae	Indian paddy field pipit	<i>Anthus rufulus</i>	Insectivorous	Least Concern	
41		Paridae	Great tit	<i>Parus major</i>	Insectivorous	Least Concern	
42		Alaudidae	Black lark	<i>Melanocorypha yeltoniensis</i>	Insectivorous	Least Concern	
43		Corvidae	Sand lark	<i>Alaudala raytal</i>	Insectivorous	Least Concern	
44			Jungle crow	<i>Corvus culminatus</i>	Frugivorous	Least Concern	
45			House crow	<i>Corvus splendens</i>	Omnivorous	Least Concern	
46		Monarchidae	Paradise flycatcher	<i>Terepsiphone paradisi</i>	Insectivorous	Least Concern	
47		Muscicapidae	Blue flycatcher	<i>Cyornis tickelliae</i>	Carnivorous	Least Concern	
48		Chloropseidae	Leaf bird	<i>Chloropsis flavipennis</i>	Insectivorous,	Vulnerable	
					Frugivorous, Nectivorous		
49		Laniidae	Brown shrike	<i>Lanius cristatus</i>	Insectivorous	Least Concern	
50		Muscicapidae	Shama bird	<i>Copsychus malabaricus</i>	Insectivorous,	Least Concern	
					Frugivorous		
51				Slaty backed forktail bird	<i>Enicurus schistaceus</i>	Carnivorous	Least Concern
52				Plumbeous redstart	<i>Rhyacornis fuliginosa</i>	Insectivorous,	Least Concern
						Carnivorous	
53		Leiotherichidae	Grey sibia	<i>Heterophasia gracilis</i>	Insectivorous,	Least Concern	
				Frugivorous			
54			Jungle babbler	<i>Turdoides striata</i>	Insectivorous,	Least Concern	
					Granivorous,		
					Nectivorous		
55			Common babbler	<i>Argya caudata</i>	Omnivorous	Least Concern	
56			White crowned shrike	<i>Eurocephalus ruppelli</i>	Carnivorous	Least Concern	
57	Psittaculidae	Alexandrine parakeet	<i>Psittacula eupatria</i>	Omnivorous	Near Threatened		
58	Laniidae	Red breasted parakeet	<i>Psittacula alexandri</i>	Frugivorous,	Near Threatened		
				Granivorous			
59	Pnoepygidae	Pygmy cupwing	<i>Pnoepyga pusilla</i>	Insectivorous	Least Concern		
60	Pellorneidae	Brown capped babbler	<i>Pellomeum fuscocapillus</i>	Insectivorous	Least Concern		
61			Marsh spotted babbler	<i>Pellomeum palustre</i>	Insectivorous	Vulnerable	
62	Paradoxornithidae	Grey headed	<i>Psittiparus gularis</i> parrot bill	Insectivorous	Least Concern		
63	Corvidae	Red whiskered bulbul	<i>Pycnonotus jocosus</i>	Insectivorous,	Least Concern		
				Frugivorous			
64			Black browed treepie	<i>Dendrocitta frontalis</i>	Carnivorous	Least Concern	
65	Passeridae	House sparrow	<i>Passer domesticus</i>	Insectivorous	Least Concern		
66			Tree sparrow	<i>Passer montanus</i>	Insectivorous	Least Concern	
67	Paridae	Oriental tit	<i>Parus minor</i>	Omnivorous,	Least Concern		
				Molluscivorous			
68	Anatidae	Pin tail duck	<i>Anas acuta</i>	Granivorous	Least Concern		
69	Piciformes	Megalaimidae	Large green barbet	<i>Psilopogon zeylanicus</i>	Frugivorous,	Least Concern	
					Insectivorous		
70			Coppersmith barbet	<i>Psilopogon haemacephalus</i>	Frugivorous	Least Concern	

Table 1c

71		Picidae	Golden backed woodpecker	<i>Dinopium benghalense</i>	Insectivorous	Least Concern
72			Large golden backed woodpecker	<i>Chrysocolaptes guttacristatus</i>	Frugivorous, Insectivorous, Granivorous	Least Concern
73			Rufous piculet	<i>Sasia abnormis</i>	Insectivorous, Carnivorous	Least Concern
74			Darjeeling pied woodpecker	<i>Dendrocopos darjellensis</i>	Insectivorous	Least Concern
75			Heart spotted woodpecker	<i>Hemicircus canente</i>	Insectivorous	Least Concern
76			Large yellow fronted woodpecker	<i>Melanerpes flavifrons</i>	Insectivorous, Frugivorous	Least Concern
77			Rufous woodpecker	<i>Micropternus brachyurus</i>	Insectivorous, Frugivorous, Nectivorous	Least Concern
78	Bucerotiformes	Bucerotidae	Grey hornbill	<i>Ocyrceros birostris</i>	Granivorous	Least Concern
79		Upupidae	Hoopoe	<i>Upupa sp</i>	Insectivorous, Carnivorous	Least Concern
80	Gruiformes	Rallidae	Coot	<i>Fulica atra</i>	Omnivorous	Least Concern
81		Rallidae	Common moorhen	<i>Gallinula chloropus</i>	Omnivorous	Least Concern
82			Western swampphen	<i>Porphyrio porphyrio</i>	Omnivorous	Least Concern
83		Ardeidae	Little egret	<i>Egretta garzetta</i>	Carnivorous	Least Concern
84	Apodiformes	Trochilidae	Ruby throated humming bird	<i>Archilochus colubris</i>	Insectivorous,	Least Concern
85	Accipitriiformes	Accipitridae	Indian white rumped vulture	<i>Gyps bengalensis</i>	Carnivorous	Critically Endangered
86			Shikra (Indian shikra)	<i>Accipiter badius</i>	Insectivorous	Least Concern
87			Pariah kite	<i>Milvus migrans</i>	Carnivorous, Avivorous	Least Concern
88			Pied harrier	<i>Circus melanoleucos</i>	Insectivorous, Avivorous	Least Concern
89			Crested serpent eagle	<i>Spilornis cheela</i>	Insectivorous, Ophiophagous	Least Concern
90			Black winged kite	<i>Elanus caeruleus</i>	Insectivorous	Least Concern
91			Black eared baza	<i>Aviceda leuphotes</i>	Insectivorous	Least Concern
92			Changeable hawk eagle	<i>Nisaetus cirrhatus</i>	Avivorous	Least Concern
93			Crested serpent eagle	<i>Spilornis cheela</i>	Carnivorous, Avivorous, Carnivorous	Least Concern
94			White tailed eagle	<i>Haliaeetus albicilla</i>	Carnivorous	Least Concern
95	Columbiformes	Columbidae	Spotted dove	<i>Stiloopelia chinensis</i>	Frugivorous, Granivorous,	Least Concern
96			Ring necked dove	<i>Streptopelia capicola</i>	Insectivorous Frugivorous, Granivorous, Insectivorous	Least Concern
97			Emerald dove	<i>Chalcophaps indica</i>	Frugivorous	Least Concern



Table 1d

98			Red turtle dove	<i>Streptopelia tranque baricha</i>	Granivorous,	Least Concern
99			Green imperial pigeon	<i>Ducula aenea</i>	Granivorous	Near Threatened
100			Rock pigeon	<i>Columba livia</i>	Granivorous, Frugivorous, Insectivorous	Least Concern
101			Imperial pigeon	<i>Dacula</i> sp	Frugivorous	Least Concern
102			Spotted dove	<i>Spoilopelia chinensis</i>	Insectivorous	Least Concern
103	Coraciiformes	Meropidae	Chestnut headed bee eater	<i>Merops leschenaulti</i>	Insectivorous	Least Concern
104			Green bee eater	<i>Merops orientalis</i>	Insectivorous , Carnivorous	Least Concern
105		Alcedinidae	White breasted kingfisher	<i>Halcyon smyrnensis</i>	Insectivorous, Carnivorous	Least Concern
106			Common kingfisher	<i>Alcedo anthis bengalensis</i>	Carnivorous, Insectivorous, Piscivorous	Least Concern
107	Trogoniformes	Trogonidae	Red headed trogon	<i>Harpactes erythrocephalus</i>	Insectivorous	Least Concern
108	Cuculiformes	Cuculidae	Crow pheasant	<i>Centropus sinensis</i>	Carnivorous	Least Concern
109			Asian koel	<i>Eudynamis scolopaceus</i>	Omnivorous	Least Concern
110	Galliformes	Phasianidae	Black francolin	<i>Francolinus francolinus</i>	Insectivorous	Least Concern
111			Grey partridge	<i>Perdix perdix</i>	Insectivorous, Granivorous	Least Concern
112			Jungle bush quail	<i>Perdica asiatica</i>	Frugivorous, Granivorous	Least Concern
113			Red jungle fowl	<i>Gallus gallus</i>	Frugivorous	Least Concern
114			Kalij pheasant	<i>Lophura leucomelanos</i>	Omnivorous	Least Concern
115		Dicacidae	Plain coloured flower pecker	<i>Diacacum minullum</i>	Frugivorous, Nectivorous	Least Concern
116		Pycnorotidae	Black bulbul	<i>Hypsipetes leucocephalus</i>	Insectivorous, Granivorous	Least Concern
117	Ciconiiformes	Ciconiidae	Greater adjutant	<i>Leptoptilos dubius</i>	Omnivorous, Carnivorous	Endangered
118			Lesser adjutant	<i>Leptoptilos javanicus</i>	Omnivorous, Carnivorous, Piscivorous	Vulnerable
119			Black necked stork	<i>Ephippiorhynchus asiaticus</i>	Carnivorous	Near Threatened
120			Scaly breasted stork	<i>Anastomus oscitans</i>	Carnivorous, Molluscivorous	Least Concern
121	Suliformes	Phalacrocoracidae	Little cormorant	<i>Microcarbo niger</i>	Carnivorous	Least Concern
122			Large cormorant	<i>Phalacrocorax carbo</i>	Carnivorous	Least Concern
123			Indian cormorant	<i>Phalacrocorax fuscicollis</i>	Carnivorous	Least Concern
124		Anhingidae	Darter	<i>Anhinga</i> sp	Piscivorous	Near threatened
125	Pelecaniformes	Ardeidae	Great egret	<i>Ardea alba</i>	Carnivorous	Least Concern
126			Medium egret	<i>Ardea intermedia</i>	Carnivorous	Least Concern
127			Cattle egret	<i>Bubulcus ibis</i>	Carnivorous	Least Concern

Table 1e

128			Yellow bittern	<i>Ixobrychus sinensis</i>	Carnivorous	Least Concern
129			Chinese pond heron	<i>Ardeola bacchus</i>	Carnivorous	Least Concern
130			Indian pond heron	<i>Ardeola grayii</i>	Carnivorous	Least Concern
131			Night heron	<i>Nycticorax nyctanassa</i>	Insectivorous, Avivorous,	Least Concern
132			Bittern	<i>Botaurus stephens</i>	Piscivorous Carnivorous	Least Concern
133			Tiger heron	<i>Tigrisoma lineatum</i>	Carnivorous	Least Concern
134	Strigiformes	Strigidae	Jungle owlet	<i>Glaucidium radiatum</i>	Insectivorous	Least Concern
135			Brown fish owl	<i>Ketupa zeylonensis</i>	Piscivorous	Least Concern
136		Tytonidae	Barn owl	<i>Tyto alba</i>	Carnivorous	Least Concern
137	Anseriformes	Anatidae	Common pochard	<i>Aythya ferina</i>	Molluscivorous, Insectivorous	Vulnerable
138			Lesser whistling duck	<i>Dendrocygna javanica</i>	Gregarious	Least Concern
139			Brahminy duck	<i>Tadorna ferruginea</i>	Omnivorous	Least Concern
140	Charadriiformes	jacanidae	Pheasant tailed jacana	<i>Hydrophasianus</i>	Insectivorous,	Least Concern
141					<i>chirurgus</i>	Molluscivorous
142		Charadriidae	Bronze winged jacana	<i>Metopidius indicus</i>	Insectivorous	Least Concern
142			Red wattled lapwing	<i>Vanellus indicus</i>	Insectivorous,	Least Concern
143					Granivorous, Molluscivorous	
143			Spur winged lapwing	<i>Vanellus spinosus</i>	Insectivorous	Least Concern
144		Laridae	River tern	<i>Sterna aurantia</i>	Insectivorous, Piscivorous	Vulnerable
145			Indian robin	<i>Saxicoloides fulicatus</i>	Insectivorous	Least Concern
146		Scolopacidae	Long toed stint	<i>Calidris subminuta</i>	Insectivorous, Molluscivorous, Granivorous	Least Concern

\***Piscivorous:** fish eater birds; Palynivorousbirds: pollen eater birds; Nectivorous: nectar eater birds; Ophiophagous: snake eater birds; Mucivorous: mucus eater birds; Molluscivorous: mollusce eater birds; Granivorous: grain eater birds; Frugivorous: fruit eater birds; Carnivorous: meat eater birds; Avivorous: bird eater birds.

\*\*Extinct in the wild (EW) - Known only to survive in captivity; Extinct (EX) - No known individuals remaining; Critically Endangered (CR) - Extremely high risk of extinction in the wild; Endangered (EN) - High risk of extinction in the wild; Vulnerable (VU) - High risk of endangerment in the wild; Near Threatened (NT) - Likely to become endangered soon; Least Concern (LC) - Lowest risk. Does not qualify for a more at risk category; Data Deficient (DD) - Not enough data to assess its risk of extinction; Not Evaluated (NE) - Has not yet ben evaluated against the criteria.

Noticably, Passeriformes showed highest species frequency with 46.25% abundance which was followed by Accipitriformes (6.8%), Piciformes and Pelecaniformes(6.12%) ;Columbiformes(5.44%); Galliformes and Charadriiformes(4.76%); Gruiformes, Coraciiformes, Suliformes, Ciconiiformes(2.72%); Strigiformes, Anseriformes(2.04%); Cuculiformes, Bucerotiformes (1.36%); Trogoniformes, Apodiformes (0.68%). Under 17 spotted orders, total 76 families were recorded. Passeriformes came out with maximum number of families (44).

The IUCN status of avian species richness of Lumding revealed that 92.51% birds of Lumding are Least Concern indicating their sufficient existence, where as 3.4% are Near Threatened, 2.72% are Vulnerable, 0.68% are Critically Endangered and Endangered. Birds like River tern, Common pochard, Lesser adjutant, Greater adjutant, Indian white rumped vulture, Marsh spotted babbler, Black necked stork, Alexandrine parakeet, Darter which were found under Critically Endangered, Near Threatened, Vulnerable and Endangered category of IUCN demands our attention to work for their conservation and protection from becoming extinct.

## CONCLUSION

Its extremely worrying that the number of Critically Endangered birds on the IUCN Red List continues to increase, despite successful conservation initiatives around the world”, says Simin Stuart, Chair of IUCN’s Species Survival Commission. Hence, it becomes immense important to work for the safety and security of birds at global level. Although our work was very small but we expect that it will motivate other researchers to work in this field so that rare species of birds can survive for long.

## ACKNOWLEDGEMENTS

Authors are grateful to Dr. Ghosh (Ornithologist) and his team from Zoological Survey of India, for their full hearted cooperation, involvement and remaining physically present during the investigation period and helping in identification of birds.

## REFERENCES

- Ainsworth, G.B., Fitzsimons, J.A., Weston, M.A. and Garnett, S.D. (2018). The culture of bird conservation: Australian stakeholder values regarding iconic, flagship and rare birds. *Biodiversity Conservation*, 27(2), 345-363.
- Arya, M., Rao, R.J. (2014). Avifaunal occurrence and distribution of wetland birds in Sakhya Sagar and Madhav lakes in Madhav National Park, Shivpuri, India. *Journal of Environmental Biology*, 35:703-708.
- Balasure, S., S. Dutta and Vyas, V. (2012). Avian diversity in Barna wetland in Narmada Basin in Central India. *Journal of Research in Biology*, 2(5):460-468.
- Beal, M., Dias, M.P., Phillips, R.A., Opper, S., Hazin, C., Pearmin, E.J., Cayrt, P. (2021). Global political responsibility for the conservation of albatrosses and large petrels. *Sci. Adv.*, 27, eabd7225.
- Borray- Escalante, N.A., Mazzoni, D., Ortega-Segalerva, A., Arroyo, L., Morera-Pujol, V., Gonzalez-Solis, J., Senar, J.C. (2020). Diet assessments as a tool to control invasive species: Comparison between Monk and Rose-ringed parakeets with stable isotopes. *J. Urban Ecol.*, 6, juaa005.
- Colin J, Bibby Nell, D. (1992). *Birds census techniques*. Academic press limited, London.
- Core Team. R. (2020). *A Language and Environment for Statistical Computing*, R. Foundation for Statistical Computing: Vienna, Austria.
- Gill, F. and Wright, M. (2006). *Birds of World: Recommended English Names*. Princeton NI: Princeton University Press, Version 2.2 generated on 2009-08-25.
- Grimmett, R., I., Inskipp, C. And Inskipp, T., (1998). *Birds of the Kotzebue south region, Alaska Jr. Columbia. Natural History of the Cook Inlet Region, Alaska*. 21:1-87.
- Grimmett, R., Inskipp, C. and Inskipp, T. (2000). *The pocket Guide to the Birds of Indian Subcontinent*. Oxford University Press, Delhi.
- Hernandez-Brito, D., Carrete, M., Blanco, G., Romero-Vidal, P., Senar, J.C., Mori, E., White, T.H., Luna, A. and Tella, J.L. (2021) *The Role of Monk Parakeets as Nest-Site Facilitators in Their Native and Invaded Areas*. *Biology*, 10, 683.
- ISSG. *Myiopsita monachus* (Monk Parakeet). *Global Invasive Species Database (GISD)*. Available online :<http://www.issg.org/database>.
- Javed, S. and Rahmani, A.R. (1993). Conservation of the avifauna of Dudhwa national Park, India. *Forktail*. 14:57-66.
- Joshi, P. and Shrivastava, V.K. (2012). Avifaunal diversity of Tawa Reservoir and its surrounding areas of Hoshangabad District (MP). *International Journal of Plant, Animal and Environmental Sciences*, 2(1): 46-51.
- Khan, A.K., Rao, R.J. and Wani, R.A. (2012). Studies on bird diversity of Overa-aru Wildlife Sanctuary of Jammu and Kashmir, India. *Journal of Threatened Taxa*. 4(13): 3228-3232.
- Miskelly, C.M., Gilad, D., Taylor, G.A., Tennyson, A.J., Waugh, S.M. (2019). A review of the distribution and size of gadfly petrel (*Pterodroma* spp.) colonies throughout New Zealand. *Tuhinga*, 30, 93-173.
- Pawar, S.K. and Salunkhe, P.S. (2014). The migratory birds in and around Pandharpur city with special reference to Takali (Padmavati) Lake, Pandharpur, Dist. Solapur (M.S.) *Avishkar-Solapur University journal*, Vol. 3. pp-38-44.
- Pitera, A.M., Branch, C.L., Sonnenberg, B.R., Benedict, L.M., Kozlovsky, D.Y. and Pravosudov, V.V. (2021). Reproduction is affected by individual breeding experience but not pair longevity in a socially monogamous bird. *Behav. Ecol. Sociobiol.*
- Peralta-Sanchez, J.M., Colmenero, J., Redondo-Sanchez, S., Ontanilla, J. And Soler, M. (2020). Females are more determinant than males in reproductive performance in the house sparrow *Passer domesticus*. *J. Avian Biol.*
- Sauve, D., Carmantier, A., Hatch, S.A., Friesen, V.L. (2021) Environmental conditions variably affect growth across the breeding season in a subarctic seabird. *Oecologia*, 1-12.
- Senar, J.C., Domenech, J., Arroyo, L., Torre, I.; Gordo, D. (2019). An evaluation of monk parakeets damage to crops in the metropolitan area of Barcelona. *Anim. Biodivers. Conserv.*, 39, 141-145.
- Vinod Shankar, R., Bhavanarayeni, U.R.S., Bayani, R, A. and Kunte. K. (2023). *Indian Birds in The IUCN Red List*. In Satose, Bayani, R, A., Ramachandran, V., Roy, P. and Kunte, K. (chief editors). *Birds of India*, v.2.17.
- Watanuki, Y., Yamamoto, M., Okado, J., Ito, M., Sydeman, W. (2022). Seabird reproductive responses to changing climate and prey communities are mediated by prey packaging. *Mar. Ecol. Prog. Ser.*, 683, 179-194.

# Standardization and Evaluation of Buffers: A One Step DNA Extraction Protocol from Microbial Cultures

Poornima Shyam<sup>1</sup>, Dr. Shyama Subramaniam<sup>2</sup> and

Dr. Shyam Subramaniam<sup>3</sup> and Edwina Jospiene M<sup>4</sup>

<sup>1,2 & 4</sup>Regenix Super Specialty Laboratories Private Limited

(Affiliated to University of Madras, Chennai India).

<sup>1,3</sup>-Apollo Lab Services, Chennai India

## ABSTRACT

Bacteria are one of the most diverse organisms that exist. Every day labs are flooded with cultures to detect the type of microorganisms and their antibiotic resistance patterns to treat patients. However, this reporting takes 3 days. Molecular biology is a field that is associated with quicker reporting. This work is an attempt to make molecular testing easier and quicker to help with one step DNA extraction method. 3 different buffers were tested for DNA extraction from cultures. The result of the best buffer is represented in agarose gel electrophoresis. Efficient extraction of DNA from bacteria was seen. The first step to standardization of buffers for the one-step extraction of DNA from bacteria. The buffers need to be tested and standardized with a wider variety of samples of both gram-positive and gram-negative organisms and an efficient method of extraction from the primary samples instead of the culture plates has to be standardized. This is a pilot study to screen the buffers for their effectiveness.

**KEY WORDS:** BACTERIAL DNA, GRAM-POSITIVE, GRAM NEGATIVE, BUFFERS, AGAROSE GEL ELECTROPHORESIS.

## INTRODUCTION

Infection by microorganisms is an overwhelming scenario that causes serious implications in the patients and sometimes even causes death. Bacteria are a major cause of these infections. Bacteria are defined as a group of small, single-celled organisms. The extent of damage is measured by the route, number, mode of transmission, and stability of the host. While the nose, throat, and cuts in the skin are foregrounds of microbial infection, the lungs, bowel, urinary tract, vagina, etc are also sites of serious infections, (Cars & Nordberg 2005, Weber & Stilianakis 2008 Reynolds & Kollef 2021).

The major mode of infection is entry through the skin and replication in the tissue. The incidence of infectious diseases was estimated at 71.8 per 10,000 population in 2022. These infectious diseases kill about nine million individuals per year among which children under the age of 5 are a major part. Bacterial diseases causing infection have reached various advanced levels of testing. Techniques of Lateral flow assay, ELISA, and molecular testing techniques have been employed (Lamballerie et al., 1992, Boehringer and O'Farrell, 2021).

However, for infections in wounds, the techniques of microbial identification and consequently identification of the antibiotics to treat the infection is a 3-day process. The samples for anti-microbial testing are in the order of sputum, urine, high vaginal swab, wound swab, and finally blood. Anti-microbial resistance due to prolonged use of antibiotics is one of the biggest challenges of the microbial industry today. Some of the major bacteria that cause infections are *Klebsiella spp.*, *Moraxella spp.*, *Escherichia spp.*, *Pseudomonas spp.*, *Staphylococcus spp.* etc., (Bharathi et al., 2010, Reynolds & Kollef 2021).

The constant evolution of bacteria poses a major challenge in identification. Testing like miRNA and SNP may be an apt solution (Ali et al., 2017). The bacterial cell wall is made up of rigid structures of uniform thickness around the cell. This rigidity ensures the various types of shapes like spiral, rod, and coccus and patterns of colonization. Based on cell wall components bacteria are divided into Gram-positive and Gram-negative bacteria (Schleifer et al., 1972). The presence of a peptidoglycan layer in the outer layer of the Gram-negative bacteria makes the need for the buffer to lyse the cell wall different from that of the gram-positive bacteria.

This study deals with the standardization of a buffer that extracts bacterial DNA from overnight culture plates in one step. The purity of the DNA obtained was screened with A

**Article Information:**\*Corresponding Author: [rksingh1@amity.edu](mailto:rksingh1@amity.edu), huda.

Received 15/04/2023 Accepted after revision 28/06/2023

Published: June 2023 Pp- 100-103

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.6>

260/ 280 and confirmed with Agarose gel electrophoresis. The results displayed below are of the best methods that yielded a good quantity of DNA.

## MATERIALS AND METHODS

The bacteria for testing was taken from ATCC strains. The confirmation was done by staining and biochemical studies. One gram- Positive and one gram-negative bacteria were taken for study. Gram Positive – *Staphylococcus aureus*- ATCC @ 25923 Gram Negative - *Escherichia coli* ATCC 25922. The same was inoculated for overnight growth in nutrient broth. One ml of broth was taken centrifuged and used for the rest of the experiment.

**Buffers:** The below table contains the constituents of the best 3 buffers which were taken for further studies.

Absorbance at 260 nm, 280nm, and 230nm.: NanoDrop® ND-1000 was used to take the readings (5). Quantification of DNA : dsDNA concentration =  $50 \mu\text{g/mL} \times \text{OD}_{260} \times$  dilution factor.

**Gel Electrophoresis:** A 1% gel was used for the analysis of the result. Only pure DNA was used for this analysis. The gel pictures of the best results are represented in the results section. The running buffer of TAE and TBE was used. To visualize DNA EtBr (Ethidium Bromide) was added and visualized through a UV transilluminator and documented through a gel documentation system (Lee et al., 2012).

## RESULTS AND DISCUSSION

All experiments have been carried out in triplicate. The below results are the representatives of the same. The purity of the DNA obtained is seen from the A260/280 ratio. A value between 1.7 -2.0 is considered pure for ds DNA. A lower value indicated the presence of phenol, proteins, and other contaminants as they are easily absorbed at 280nm. A higher value of the ratio shows the presence of RNase. In this experiment as proteinase K is not used to avoid the barrier of storage the need to rule out contamination becomes important. DNA Purity is calculated using the below formula and represented in the table (Ning et al., 2009).

Buffer 1:	Buffer 2:	Buffer3:
Tris HCL	MgCl <sub>2</sub>	Tris HCL
KCL (Pottasium chloride)	EDTA	TE buffer
Mgcl <sub>2</sub> (Magnesium chloride)	SDS	MgCl <sub>2</sub>
SDS 2% (Sodium Dodecyl Sulfate)	Triton X100	EDTA
Triton X	PVP (Polyvinylpyrrolidone)	SDS
Ethanol	CH3COOK	PVP
	TE buffer (Tris –EDTA)	CH <sub>3</sub> COOK
	Ethanol	Ethanol
		NaCl(Sodium Chloride)

DNA Purity ( $A_{260}/A_{280}$ ) =  $(A_{260} \text{ reading} - A_{320} \text{ reading}) \div (A_{280} \text{ reading} - A_{320} \text{ reading})$  (Tiwari et al., 2017).. A260/230 is needed to rule out contaminants such as guanidine thiocyanate, guanidine HCL, triazole, phenol, etc. A ratio between 2.0 to 2.2 is considered pure for this ratio. The

below table contains the results of the nanodrop for the evaluation of the buffers: The results of the effective buffers are only represented in the table. buffers and methods which showed values of ratios below 1.5 or above 2.2 for ratios A260/A280 are not represented.


S.no.	Buffer name	Method	A260/280	A2260/230	DNA Quantification ng/μl
1.	Buffer 1	A	1.68	2.2	4.1
		B	1.75	2.19	8.3
2.	Buffer 2	A	1.85	2.18	27.2
		B	1.71	2.1	10.7
3.	Buffer 3	A	1.72	2.19	11.2
		B	1.91	2.02	33.4




This result shows that buffer 2 is more efficient for *S. aureus* – gram-positive, while buffer 3 is more efficient for *E. coli* – gram-negative. This shows PVP and  $\text{CH}_3\text{COOK}$  increase the efficiency of extraction.  $\text{CH}_3\text{COOK}$  helps with the breaking down of proteins (Angela 2017). PVP increases the DNA yield (Khan et al., 2007). Buffer 2 with the second detergent, Triton X is more efficient in the case of Gram –Positive bacteria. While Tris HCL and NaCl proved greater efficiency in the breakdown of the peptidoglycan layer presenting Gram Negative bacteria. Tris HCL helps in the maintenance of the buffer layer whereas NaCl proves efficient with the lysis of the peptidoglycans by creating pores on the surface of the cell wall.

**Figure 1: Lane 1 –ladder Lane 2 and 3, Buffer 3 –*Escherichia coli*. Lane 4 and 5 , Buffer2 – *Staphylococcus aureus***

### Gel electrophoresis



**Figure 2: DNA after ethanol wash. Lane 1: DNA Ladder Lane 2 : gram positive bacteria from kit method. Lane 3 : gram negative bacteria from kit method . Lane 4 and 5- Gram negative bacteria from buffer 3. Lane 6 and 7 Gram positive bacteria from buffer 2**



Separation of DNA is seen. Addition of Ethanol to collect the DNA yields pure DNA that can be used for further study.

Efficient extraction comparable to a CE-approved extraction kit was seen.

## CONCLUSION

Efficient extraction of DNA from bacteria was seen. The first step to standardization of buffers for the one-step extraction of DNA from bacteria. The buffers need to be tested and standardized with a wider variety of samples of both gram-positive and gram-negative organisms and an efficient method of extraction from the primary samples instead of the culture plates has to be standardized. This is a pilot study to screen the buffers for their effectiveness.

**Conflict of Interest:** Authors have no conflict of interest

## REFERENCES

- Ali, N., Rampazzo, R. C. P., Costa, A. D. T., & Krieger, M. A. (2017). Current Nucleic Acid Extraction Methods and Their Implications to Point-of-Care Diagnostics. *BioMed research international*, 2017, 9306564.
- Bharathi, M. J., Ramakrishnan, R., Shivakumar, C., Meenakshi, R., & Lionalraj, D. (2010). Etiology and antibacterial susceptibility pattern of community-acquired bacterial ocular infections in a tertiary eye care hospital in south India. *Indian journal of ophthalmology*, 58(6), 497-507.
- Boehringer HR and Brendan J O'Farrell (2021) Lateral Flow Assays in Infectious Disease Diagnosis *Clinical Chem*. 2021 Dec 30;68(1):52-58. doi: 10.1093/clinchem/hvab194.
- Cars, O., & Nordberg, P. (2005). Antibiotic resistance–The faceless threat. *International Journal of Risk & Safety in Medicine*, 17(3-4), 103-110.
- De Lamballerie, X., Zandotti, C., Vignoli, C., Bollet, C., & De Micco, P. (1992). A one-step microbial DNA extraction method using “Chelex 100” suitable for gene amplification. *Research in microbiology*, 143(8), 785-790.
- Desjardins, P., & Conklin, D. (2010). NanoDrop microvolume quantitation of nucleic acids. *Journal of visualized experiments : JoVE*, (45), 2565.
- Angel DS (2017). Measuring Elasticity of Force sensing proteins using Single Molecule Force Spectroscopy (Doctoral dissertation, IISER-M).
- Khan, S., Qureshi, M. I., Alam, T., & Abdin, M. Z. (2007). Protocol for isolation of genomic DNA from dry and fresh roots of medicinal plants suitable for RAPD and restriction digestion. *African Journal of Biotechnology*, 6(3), 175.
- Lee, P. Y., Costumbrado, J., Hsu, C. Y., & Kim, Y. H. (2012). Agarose gel electrophoresis for the separation of DNA fragments. *Journal of visualized experiments : JoVE*, (62), 3923.
- Ning, J., Liebich, J., Kästner, M., Zhou, J., Schäffer, A., & Burauel, P. (2009). Different influences of DNA purity indices and quantity on PCR-based DGGE and functional



gene microarray in soil microbial community study. *Applied Microbiology and Biotechnology*, 82, 983-993.

Reynolds D & M Kollef (2021) *The Epidemiology and Pathogenesis and Treatment of Pseudomonas aeruginosa Infections: An Update Drugs* volume 81, pages2117–2131

Schleifer, K. H., & Kandler, O. (1972). Peptidoglycan types of bacterial cell walls and their taxonomic implications.

*Bacteriological reviews*, 36(4), 407-477.

Tiwari, S., Tomar, R. S., Tripathi, M. K., & Ahuja, A. (2017). Modified protocol for plant genomic DNA isolation. *Indian Research Journal of Genetics and Biotechnology*, 9(4), 478-485.

Weber, T. P., & Stilianakis, N. I. (2008). Inactivation of influenza A viruses in the environment and modes of transmission: a critical review. *Journal of infection*, 57(5), 361-373.

# Didactic Strategy for the Use of Virtual Itineraries in the Training of Graduates in Education Geography

Oliday Aguilar Espinosa<sup>1</sup>, Erich Rodríguez Vallejo<sup>1</sup>,

Rafael Armiñana García<sup>1</sup> and Rigoberto Fimia Duarte<sup>1,2\*</sup>

<sup>1</sup>Central University “Marta Abreu” of Las Villas, Villa Clara, 50100 Cuba.

<sup>2</sup>University of Medical Sciences of Villa Clara (UCM-VC), 50200 Cuba.

## ABSTRACT

The training of teachers has always been a necessity of the planet aligned to the space and time where this process takes place. Future training of graduates in education, Geography who study geographic space need novel ways to appropriate these contexts and knowledge. The objective of this research was to propose a didactic strategy with the use of virtual didactic itineraries, in the teaching-learning process for the training of education, Geography in the first year of the career: Bachelor of Education. Geography, through the subject Introduction to Labor Practice, which contributed to deepen the knowledge about different landscapes and at the same time developed professional skills related to the study of the landscape. For the development of the research, information collection methods were used (participatory observation, document review, interview, expert criteria) and information processing methods (analysis - synthesis, inductive-demonstrative, logical history, descriptive statistics, modeling and analysis). The methods allowed to determine the needs of the students from the cognitive, educational and procedural point of view, regarding the teaching-learning process. The didactic strategy was implemented in practice, achieving very satisfactory results from the 2019-2022 academic years. It allows contributing to the development of professional pedagogical skills while deepening the knowledge of different landscapes in the students of the Degree in Education. Geography of the Faculty of Secondary Education of the Central University “Marta Abreu” of Las Villas, Villa Clara, Cuba.

**KEY WORDS:** DIDACTIC STRATEGY; GEOGRAPHY; LABOR PRACTICE; TEACHING-LEARNING PROCESS; VIRTUAL ITINERARIES.

## INTRODUCTION

The International Geographical Union and the Commission on Geographical Education (1992, 2000 and 2007) propose International Declarations for the teaching of Geography where proposals are made to promote the idea of fostering knowledge of the environment close to the society where we live. Threats and risks often arise from the lack of knowledge of the inhabitants themselves, from forgetting the history accumulated in each of them (Sánchez, 2020).

The 1992 Declaration outlines that, when selecting geographic content, one can opt for or combine regional studies (e.g., studies of the region, country) or thematic studies (e.g., through the study of physical or human geography, or through centers of interest, such as environmental quality or population growth). In any case, it is pointed out that there should be a balance in the scale of analysis: local, regional,

national or global (International Geographical Union and the Commission on Geographical Education, 1992). The authors of the research consider that it is oriented to study not only the near or local space but also the distant or international space, for which it is necessary to use information and communication technologies for the purpose of teaching Geography, so it is essential for the training of teachers to develop skills related to these media.

This statement also addresses the methods for its teaching, the statement points out that they should stimulate in students the desire to engage in questioning and inquiry. It is essential that students exercise and develop geographical skills with a tendency to search for solutions to the problems posed in the organization of space, present and future (International Geographical Union and the Commission on Geographical Education, 1992). Another Declaration of great interest is the Seoul Declaration of 2000, which proclaims how geographic education is fundamental for people to develop different capacities in relation to cultural diversity such as:

**Article Information:**\*Corresponding Author: [rigoberto.fimia66@gmail.com](mailto:rigoberto.fimia66@gmail.com)

Received 23/02/2023 Accepted after revision 18/06/2023

Published: June 2023 Pp- 104-114

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <http://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.7>

The ability to defend and be sensitive to human rights;

The ability to understand, accept and appreciate cultural diversity;

The ability to understand, empathize and criticize alternative points of view about people and their social conditions;

To be aware of the impact of their own lifestyles on their social contexts;

Appreciate the need to protect the environment and provide environmental justice to regions and local communities that have suffered environmental devastation;

Ability to act as an informed and active member of both their own society and the global society (International Geographical Union and the Commission on Geographical Education, 2000).

It is a Declaration that insists on intercultural studies from the perspective that geographic education also helps to understand the need to protect the environment at different scales, both local and international, and contributes to educate people active in the global society. (International Geographical Union and the Commission on Geographical Education, 2000).

Subsequently, the 2007 Lucerne Declaration on Geographical Education for Sustainable Development prioritizes issues affecting ecological-holistic thinking, and focuses on three a

A. The Contribution of Geography to Education for Sustainable Development.

B. Criteria for Developing a Geographic Curriculum for Education for Sustainable Development.

C. The Importance of Information and Communication Technologies (ICT) in Education for Sustainable Development in Geography (International Geographical Union and the Commission on Geographical Education, 2007).

This same Declaration emphasizes that the teaching of Geography and Education for Sustainable Development are closely linked, because the acquisition of geographic competencies can improve the understanding of sustainable development. For this reason, it insists on the three competence areas: knowledge, skills and attitudes.

Through geographic knowledge and understanding of natural systems, interaction between ecosystems, and socio-economic systems helps students understand the world through key geographic concepts such as: location, distribution, distance, movement, region, scale, spatial association, spatial interaction, and change over time; Work on communication skills, reasoning, and practical and social skills help explore geographic issues at a range of levels from local to international. With the development of attitudes and values to seek solutions to local, regional,

national and international questions and problems on the basis of the Universal Declaration of Human Rights (International Geographical Union and the Commission on Geographical Education, 2007).

Education, in its dynamics, does not escape the significant changes in its teaching-learning process (TLP), by taking into account the growing use of information and communication technologies, as essential elements for the fulfillment of the programs offered by educational institutions (Lima and Fernandez, 2016). Current education does not take advantage of the technological resources at its disposal, much less, all the possibilities offered by connectivity (Garrison, 2016). Another problem is related to the consensus about the common denomination to express e-learning (Lin et al., 2017).

The authors of this research consider that for the success of this purpose it is necessary to train teachers capable of fulfilling this social task, therefore their training must meet the demands of contemporary Geography teaching. With this purpose of studying the geographic space in an integral way at all scales (global, regional and local), from the ASP, multiple researchers at the international and national level have addressed, in a precise and contextualized manner, the subject matter of the research, the following stand out in recent years (Vilarrasa, 2003; Berger, 2006; Ruiz and González, 2007; Maderuelo, 2007; Sánchez, 2010; Garrido, 2014; Licerias, 2018; Alcántara and Medina, 2019).

Many strategies, projects, research, study programs and other elements such as work practice and field work have been proposed, but there are insufficient proposals from science for the use of connectivity as a teaching tool for the study of landscapes at different scales (Licerias, 2018; Ruiz and González, 2007; Berger, 2006).

From the use of the observation method, it has been possible to observe: weak preparation of the graduates in Education. Geography for the use of virtual didactic itineraries, insufficiencies related to this strategy such as: scarce use of didactic itineraries to promote the study of geographical space and scarce use of local values and technological means available in educational institutions, in addition, geographical knowledge is prioritized while professional skills are little promoted from the (TLP), scarce theoretical and methodological foundations for the use of didactic itineraries to contribute to the training of graduates in education, Geography and the contributions related to the study of geographic space at different scales in a virtual way are insufficient.

In order to carry out this didactic strategy, it is important how this knowledge is transmitted, therefore, active methodologies based on collaborative and/or individual learning are needed. The combination of other teaching strategies and the variety of didactic resources is also a necessity to favor motivation, such as: case studies, field work, didactic itineraries, games and simulations, involving the use of new technologies, handling of Geographic Information Systems, digital cartography, statistical data, use of the web, creation of a notebook, portfolio or dossier,

virtual trips, project work, worksheets, analysis of images, stories and poems, creation and solution of problems, among others, that help them to understand and deepen their knowledge, to acquire geographic skills and abilities, and to become aware of the world in which they live, in order to be sensitive to the problems that affect them, both near and far. Elements that should characterize the Geography TLP for the training of graduates in Education, Geography and thus influence the future of the new generations of this society to ensure the Education for Sustainable Development to which we aspire (De la Calle, 2012).

Although different agencies have emphasized the implementation of a sustainable model with an increasing amount of didactic resources, there is still progress to be made in linking science in society. It also states that this is possible through the use of geographic information technologies that allow the creation of virtual databases using virtual globes such as Google Earth and other geometric applications (Oyarce, 2018).

A didactic itinerary is a route or path to follow with different stopping points or points of interest in certain elements of heritage or cultural value. A field trip, previously organized in the classroom by the teacher, and which aims to make a series of stops in those places that you want to make known to the students, but that do not have to be heritage or cultural, because, if we want to know, for example, the commercial activities of a city, such as the market and the different types of existing trade, this didactic tool can also be used (Liceras, 2018; Campo et al., 2016).

According to the same author, it is a tool that provides the keys to the understanding and teaching of geographic space, constituting rich and deep learning moments that awaken interest, while teaching the student to observe, value and reflect on the surrounding environment (Liceras, 2018).

Virtual didactic itineraries allow the study of various localities through the unobstructed observation of the surrounding landscape and the understanding of the history of the land recorded in rocks, structures, and geographical features (Migon, 2017). In addition, the place is identified from points or sites of interest that allow to recognize and understand a geographical space as a whole according to Palacio et al. (2016).

Taking into consideration what has been expressed in previous paragraphs, the objective of this research is to propose a didactic strategy to use virtual didactic itineraries in the TLP for the training of graduates in Education. Geography.

## MATERIAL AND METHODS

The research was conducted at the Faculty of Pedagogical Sciences "Félix Varela Morales" (Figure 1) of the Central University "Marta Abreu" of Las Villas, in the city of Santa Clara, province of Villa Clara, Cuba (Figure 2)

For the development of the research, the authors assume the classification of methods given by Bermúdez & Rodríguez

(2016) and Expósito et al. (2021). Information collection methods and methods for processing the information collected were used. The information collection methods used were the following:

**Figure 1: Satellite photo of the location of the Faculty of Pedagogical Sciences "Félix Varela Morales" of the Central University "Marta Abreu" of Las Villas, Santa Clara, Villa**



**Figure 2: Map of the province of Villa Clara, with its 13 municipalities. Taken from Google Scholar.**



**Participatory observation:** to detect the inadequacies of the Teaching-Learning Process (TLP) in the sample and evaluate the transformations experienced by the students during the application of the didactic strategy.

**Document review:** to verify the shortcomings and potentialities of the sample and the TLP being developed.

**Interview:** to verify the geographic knowledge, professional skills and professional motives and interests of the students before and after applying the didactic strategy.

**Expert criteria:** to evaluate the didactic strategy developed and improve it according to the suggestions made by the experts in order to achieve the proposed objective.

An expert is assumed to be a person or persons with experience in the subject, with the capacity to make judgments, criteria and evaluations that allow for the improvement of the proposal developed. The competence of the experts was determined by the K coefficient, calculated



according to the opinion of each of the candidates, regarding their level of knowledge on the subject to be addressed and the relations with the sources to argue their criteria (Armiñana, 2020).

**Table 1. Level of competence of the experts consulted for the assessment of the strategy.**

EXPERT	K	PROFICIENCY LEVEL
1	1	High
2	1	High
3	1	High
4	1	High
5	0,97	High
6	0,89	High
7	1	High

The evaluation criteria used to determine the level of competence of the experts were: high competence (0.8 to 1); medium competence (0.5 to 0.7) and low competence less than 0.5. The application of the competence coefficient made it possible to select 7 highly competent experts, on the basis of their knowledge in the training of graduates in Education, Geography and Higher Education.

The aspects on which they had to make their judgment were: objective of the strategy, logical structure, contextualization, contribution to the training of B.Ed., Geography, uniformity for the treatment of geographic knowledge and professional skills in the career of B.Ed. in Geography. Geography.

For the evaluation of these aspects, a scale from 1 to 5 was proposed, where (1) is inadequate, (2), not very adequate, (3) moderately adequate (4) quite adequate and (5) very adequate. The methods used to process the information collected (intellectual) were as follows:

**Analysis-synthesis:** to analyze the subject of study, approached by various researchers at national and international level; in the choice of methods; the analysis and synthesis of the results of the initial and final survey, and in the design of the didactic strategy.

**Inductive-demonstrative:** to verify from the bibliographic exploration and applied methods, the knowledge, skills and professional pedagogical attitudes of the sample of selected students of the Bachelor's degree in Education. Geography and to demonstrate the certainty of the research proposal.

**Historical-logical:** to implement the theoretical and methodological foundations related to the evolution of the use of virtual didactic itineraries in Cuba and the world.

**Modeling:** to model the didactic strategy in the Bachelor's Degree in Education. Geography; starting from the general objective, the stages, actions and the interrelations among them to fulfill the proposed objective.

The methods of processing the information collected (Statistics) were descriptive statistics for the analysis of frequency distribution tables and elaboration of graphs; which allowed quantifying the results for their subsequent description in the results.

The population coincides with the sample according to the interests of the researchers, and consisted of 14 students, which is the totality of the enrollment of the first year of the Bachelor's Degree in Education. Geography of the Regular Day Course 2021.

## RESULTS AND DISCUSSION

The application of the scientific methods allowed obtaining the following results:

The review of guiding documents such as the Study Plan and Model of the Professional of the Bachelor's degree programs in Education. Geography, evidenced that the professional skills of landscape interpretation and work with technological media are oriented from the objectives and curricular contents of the career. The use of virtual didactic itineraries as a didactic strategy is not required.

An initial interview was carried out individually with the 14 students of the first year of the Regular Day Course of the Geography career. The objective was to ascertain the geographic knowledge, professional skills and motives and interests of the students about the TLP they would like to receive, as well as the elements that give them pleasure during the Geography TLP.

In the first question related to the geographical phenomena found in a given landscape, only 14.0%, recognized them, but did not establish the relationships between the phenomena that make it up. In the second question, 10.0% of the students identified the problems of a given landscape, but failed to explain them in context. In the third question, the same number of students recognize the potential of this discipline for the study of geographic space, through virtual itineraries, but none of them managed to exemplify it.

In the fourth question, 3.0% of the sample, in the fourth question, show the professional skills achieved and no student is capable of proposing other examples to develop it in the school. Question 5 addresses the level of satisfaction of the students with the TLP they receive, which is 50.0%. With this instrument it was evidenced the characteristics that students demand in the TLP for the formation of the future B.A. in Education., Geography, being these:

- Expanding their knowledge.
- Develop cartographic skills.
- Knowledge exchange between students and teachers.
- Collaborative learning.
- Variety of learning environments, both real and virtual.
- Scientific research.
- Search for information from different sources.
- Students as protagonists of this process.
- Use of different forms of organization.



- Varied means of teaching.
- Use of the most effective methods for learning.
- Development of positive human attitudes and values.

With the participant observation of teaching activities, a total of seven were observed, with the purpose of verifying the development of the Geography TLP in the main integrating discipline and the subjects of the academic year, distributed as follows: two of Physical Geography, one of Seminar and another of Lecture, two of Cartography, one Lecture and another Practical Class, three Workshops, two of Work Practice and another of Research Methodology. The following difficulties were identified:

- Scarce treatment of the knowledge of cause-effect and geographical regularities.
- Poor adjustment of the process to the students' motivations and interests.
- Insufficient linkage of the subjects of the academic year with professional pedagogical skills.
- Scarce technological resources are used.
- Insufficient representations are used to illustrate knowledge.
- A large amount of information is transmitted, with little explanation of how to transmit it.
- Little use of varied learning environments (virtual or real) and at different scales (global, regional and local).
- Insufficient use of knowledge to strengthen emotions, feelings, values and positive attitudes towards the profession.

The analysis of the documents consulted, the results of the interview with students and teachers, as well as the participant observation, made it possible to determine the following Potentialities: The use of technological means, the study of various spaces and at different scales is included in the normative guiding documents such as Study Plan and Model of the professional of the Bachelor's Degree in Education. Geography, the use of technological means, the study of various spaces and at different scales.

- The motivation and interest of the students for the profession and the career.
- The experience of the teaching staff.

The main inadequacies in the teaching-learning process of Geography for the formation of the Bachelors in Education, Geography were:

- Little use of the Geography ADP for educational work and the development of professional skills.
- Little use of other forms of organization such as: seminar, practical class, workshop and geographic excursion.
- Insufficient use of the technological resources available to them.
- Insufficient motivation, meaning and involvement of students in learning.
- Insufficient use of the potential of the study plan for the benefit of the Geography teacher's training.
- Lack of interest by the teacher of how his student learns

and how much he contributed to his formation as a teacher and as a man for life in society.

- Insufficient treatment of cause-effect knowledge and geographical regularities.

Based on the results achieved with the applied methods, the didactic strategy is proposed with the use of virtual didactic itineraries in the TLP for the training of graduates in Education, Geography, which was submitted to the experts' criteria. From the analysis carried out by the experts, the following results are specified:

- 93, 3%, 6 of the experts consulted, estimated that the objective of the strategy was adequate.
- For 5 of the experts, 86.6%, the structure and contextualization of the strategy was very adequate, while the rest considered it to be quite adequate.
- In relation to the actions to achieve unity in the treatment of professional knowledge and skills, only 20.0% considered it to be quite adequate, while 80.0% considered it to be very adequate.
- 93.3% considered the way to establish integration between professional knowledge and skills to be very adequate; only one expert considered it to be not very adequate.
- 100% of the experts consulted agree that the proposal contributes to the training of graduates in Education, Geography of the first year of these careers, they evaluate this aspect as very adequate.

The following are some of the assessments made by the experts. What they expressed has been taken literally.

Expert 1 "I consider that the didactic strategy is pertinent, applicable, shows a logical structure; the professional knowledge and skills have a specific character for the Geography career".

Expert 2 "The theoretical and methodological conception of the proposal responds to the proposed objective".


Expert 4 "The actions conceived in stages respond to a logical, coherent procedure, which integrates professional knowledge and skills in correspondence with the professional's Model".

Expert 5 "I consider that the strategy is novel and opportune to incorporate students who are starting out in a more dynamic and motivating process, so that they acquire knowledge that they can apply in the profession and in daily life and that a change of attitude is evidenced in them, the transformation into a professional consistent with the world where he/she lives. It is necessary to involve the faculty so that, together with the students, the objective proposed in the strategy can be achieved".

Expert 7 "The proposal evidences the principles of education and instruction and preparation for life, which are not circumscribed to a career, discipline or year, but should be approached in an integral manner in all university processes, with the participation of students and professors in achieving a better teacher training".

The criteria, judgments and suggestions issued by the experts were given by: to continue perfecting the actions to achieve an adequate use of the Internet, to establish adjustments for its use in the strategy, taking into account the diagnosis of the group. After reviewing the proposal by the experts, the researchers made adjustments, reworked the didactic strategy and sent it back to the experts, who evaluated it and expressed that it was pertinent to be applied. The following is the model of the Didactic Strategy designed (Fig. 3).

Figure 3: Didactic Strategy Modeling.



The execution of this didactic strategy was carried out through the use of virtual didactic itineraries in four steps: preparation, research, realization and conclusions, each with specific characteristics, where the student is the protagonist of the activity and the teacher serves as a guide, controller and animator.

### Preparation stage

- It began with the elaboration of a program of Introduction to Labor Practice for the use of virtual didactic itineraries.
- The proposal was presented to the discipline and year group for its approval, planning and integration of different contents and subjects of the year.
- Administrative procedures were carried out.
- The necessary material resources for the activity were organized.
- Virtual didactic itineraries, points of interest or stops were analyzed and selected with the students, using cartographic means, taking into account the combination of different scales: global, regional and local.
- The motivation of the activity was carried out.

- Each of these virtual didactic itineraries was visited and studied in order to know the potentialities and possibilities offered by the networks for their study, as well as to define which one is the most suitable for the proposed objectives.
- A preliminary study of the itinerary to be carried out was carried out.
- The students were organized into study teams and those in charge were selected.
- The information search and scientific research activities carried out by the students in the following stage were planned and oriented, as well as the bibliography they used or the Internet sites they consulted.

### In the research stage:

- The teacher examined various means to interest and motivate his students for the topic of the activity.
- The students elaborated and applied research techniques such as: interview, survey or any other that the teacher determined, to specialists or experts on the topic.
- Maps of different scales and themes were consulted to contrast and compare the regularities of the phenomenon in different contexts (digital maps).
- The didactic potentialities of each studied itinerary were determined.
- Students located on maps the areas of study and traced the itinerary to be followed together with their teacher, according to their cognitive interests.
- The students searched for information in different internet sites about the proposed itineraries and elaborated the observation guide for each studied landscape, then compared it with other teams and reached a consensus about the worksheet to be solved during the realization.
- The students, with the help of their teacher, analyzed the information obtained and determined the didactic resources to be addressed during the activity and determined the points of interest during the tour.
- They expressed their opinions about the results obtained in the research.
- A portfolio or dossier was prepared with the information collected, to be consulted in the next stage.

### In the realization stage:

- The teacher guided, oriented, directed, proposed, facilitated, asked questions where to direct the observation and what elements to look for, at the end settles the knowledge.
- The students, organized in work teams, investigated what was guided, taking into account the didactic points of interest.
- Procedures such as observing, identifying, describing, inventorying, analyzing, understanding, relating, explaining, evaluating, concluding, comparing, extrapolating, generalizing and enjoying while learning were developed.
- The perception of the landscape was enhanced through the following sequence: perceive-observe-read-analyze-conclude, from different angles of vision or

perspectives.

- The observations, descriptions and activities carried out made up the reading of the study landscape.
- Virtual spatial boundaries were established.
- The most relevant elements of the landscape were distinguished and described, going through the entire study area.
- The observation and perception of the landscape was compared with the research carried out.
- A common objective perspective was established for all.
- The reading of the landscape began taking into account the following levels: perception, analysis, structure and application.
- The points of interest were located and located with the use of the superimposition of images and digital maps.
- A dense description of the landscape was made, taking into account the relationship between components, causes and reasons for processes and phenomena.
- The didactic principles of Geography were addressed, such as: location, distribution, permanence, change and continuity.
- The aesthetic, patrimonial and identity values of the study landscape, the history and traces of history in it were specified.
- Nature-society relationships were established, environmental damage was identified and actions to minimize it were proposed.
- Interdisciplinary relationships were established.
- The causes of the processes and phenomena studied were understood, explained and valued, taking into account the contents studied, observed and investigated as a result of thinking.
- The feelings experienced during the activity, the elements that provoked those emotions, as well as the most significant lived experiences for the students were described.
- The study of global spaces and local landscapes, as well as the virtual local, were compared with the real local.
- Human values such as solidarity, responsibility, love of nature, humility, honesty, patriotism, sense of belonging and identity were strengthened.
- Professional skills and didactic procedures were developed and the didactic potentialities of the activity were addressed, as well as its contribution to their training.
- An integral evaluation of the studied space and its potentialities for the development of the Geography teaching process was carried out.

#### **In the conclusion stage.**

- Conclusions were reached and the fulfillment of the objectives of each didactic itinerary was verified.
- The experience was reflected and recapitulated, deepening in the contributions of the activity to the integral formation of the students.
- The results were presented in the form of seminars or round tables.
- The presentation of the work was organized in

exhibitions or publications where there were exhibitions of samples of natural elements, photographs, creative works made by the students, posters, murals, banners, collages, models, maps or computer products, such as web pages or interactive blogs or publications in social networks and creation of groups.

- The teacher evaluated his students, comprehensively and individually, taking into account their evolution during the whole process of the activity.
- The activities carried out were specified, the best and most complete works were selected, as well as the most outstanding students and teams in the activity. In addition, the main difficulties and achievements were specified and the students proposed goals for the next virtual didactic itinerary.

At the end of these steps and actions, a new one is started, which allowed its improvement throughout the research. A total of 6 virtual didactic itineraries were carried out, which make up this elaborated didactic strategy. At the end of its application, a final interview was applied to the students of the first year of the Bachelor's Degree in Education. Geography, with the objective of verifying the geographic knowledge and professional skills acquired and the degree of satisfaction they have in relation to the actions developed. The results obtained are presented below.

The first question related to the total determined, 99.0% of the students answered satisfactorily. Regarding the recognition of environmental problems and explanation of the same in the context (second question), 97.0% already do it correctly.

100.0% of the students recognize the potential of this discipline for the study of geographic space and exemplified it (question 3).

Question 4, which refers to the professional skills achieved, 99.0% manage to do it correctly and 90.0% are able to propose other examples to develop it at school.

Question 5 addresses the students' level of satisfaction with the process achieved, reaching 100%.

The graph (Fig. 4) shows the satisfactory results achieved by the students after applying the didactic strategy.

The results showed that the students in the sample acquired geographic knowledge while developing professional skills, according to the requirements of the TLP of Higher Education. From the actions carried out, it was possible to contribute to the formation of graduates in Education, Geography with greater deepening and integration of knowledge to a geographic space and how to carry out similar activities in the students of the Geography career in the Faculty of Secondary Education of the Central University "Marta Abreu" of Las Villas, Villa Clara, Cuba.

In the research process, it was possible to confirm that the use of virtual didactic itineraries is an effective, efficient and effective didactic strategy for the training of graduates in Education. Geography, as part of the initial training of this professional, where the teacher is the most important



element in the intellectual, affective and moral formation, focused on achieving changes in attitudes in the subjects involved, which does not differ from what is expressed by (Marrero and Méndez, 2018).

As results of the use of virtual didactic itineraries in the teaching-learning process for the training of Bachelors in Education. Geography the authors consider that:

It contributed to enhance the training of future Bachelors in Education. Geography in the following elements: greater participation of the students, as active agents within the process; they became researchers at the same time they acquired new knowledge; they were motivated and interested not only in Geography but also in their profession; this form of professional performance became a reference for them; it promoted the search for these particularities in their localities and the study of them; it promoted the systematic application of this form of professional performance;

strengthened values and positive attitudes for a culture of TLPce; promoted Education for Sustainable Development; developed professional skills and the ability to solve problems related to their profession; promoted the use of technologies as a means of teaching; the use of social networks; favored the development of skills such as observing, describing, explaining, characterizing, arguing and valuing; developed the perception of the landscape.

The influence of these technological tools in the TLP is evident in self-learning, the acquisition of knowledge, the development of personal and interpersonal skills, the promotion of creativity, and the stimulation of collaborative and committed attitudes, in addition to the use of technologies focused on the landscape as an educational resource (Botella and Hurtado, 2016). The combination of strategies and skills contributes to the achievement of different objectives, competencies and contents that characterize the training process of future teachers.

In order to teach school geography, didactic itineraries are an important resource that should be known and practiced as active teaching. In addition, it promotes motivation in students to disconnect the rhythm and routines of daily classes or to discover new emotions by performing different experiences (Campo et al., 2016).

There is correlation between students' effort, active participation in the virtual environment and reading the document prepared by the student with the evaluation of their academic performance (Dominguez et al., 2016).

The teaching-learning process contributed to the study of different geographical spaces at different scales, as Marrón (2007) points out, when the student is able to explain the diversity of existing spaces, the territorial and social inequalities that occur on the planet and within the countries or nations themselves, the effects that human action is having on Nature and the environment and the causes that motivate them, and try to provide solutions, since it is

necessary to put limits to an unlimited development in a space - the planetary - that is limited.

Batllore (2011) suggests the need to combine the large scale with the small scale because it provides a more complex view of reality. The use of virtual tools has advantages such as: lower costs. They are more motivating for learning. It has no defined limits, neither of time nor of space. It can be accessed from anywhere in the world. It can be documented with a great variety of materials and in very diverse formats.

It allows unlimited interactivity. It is an experiential experience in which concepts, skills and procedures are developed and emotions are produced. It promotes a deep knowledge of space. It demands the active participation of students. It integrates the stages of traditional fieldwork. It reinforces real observation with digital literacy while obtaining deep knowledge of the place (González and Marrón, 2012).

The authors agree with the criteria expressed by Évora (2011) when he asserts that the strategy, in the educational field, constitutes the pedagogical direction to transform the real state to the desired state of the object to be modified, from the conditioning of the system of actions between the directing subsystem and the directed subsystem to achieve the proposed objectives at its highest level.

We also agree with Charbonet (2009), when he asserts that strategies arise from dissatisfactions or contradictions between the actual and the desired state, pursue defined objectives, are organized by stages, require transforming actions that cause changes in the actual state and in their conception are present control and feedback mechanisms. In this sense, the didactic strategy assumes the postulates issued by these authors.

For Chávez et al. (2019) and coinciding with Orfa et al. (2019), the strategy becomes the teacher's actions with the purpose of facilitating training and learning, using didactic techniques which allow building knowledge in a creative and dynamic way.


The actions implemented in the didactic strategy influence the transformation of the personality of the future teacher of Geography; manifested in the acquisition of knowledge, its implementation in new situations, teamwork for the solution of problems related to the pedagogical task, willingness for active and conscious participation in environmental activities, and the planning, preparation and implementation of actions that can be put into practice from the Cuban school.

The criteria issued by the teachers and experts consulted ratify the importance of the studies conducted by Villamandos et al. (2019), related to the search for alternatives as the didactic strategy for the development of the TLP in Geography students.

The strategy perfected how to express the scientific results achieved by the students in order to socialize them, which

can be applied in performance contexts similar to the one in which the research is developed; in the opinion of the researchers, actions should be conceived in the strategy that involve Educational Research Methodology, which is the subject that has within its task, to prepare the student for this purpose.

**Figure 4: Results of the interviews before and after applying the strategy.**



The authors agree with García & Reques (2012) about the use of these tools in the teaching of Geography where it should be kept in mind that the Internet is a means not an end, that data should not be confused with information, nor information with knowledge.

In addition, García and Pascual (2012) point out the dual role of the teacher as a model of discipline, authority and for teaching to think, and as a social animator who creates learning situations and promotes their development in an adequate manner. It is also necessary to count on the disposition of students towards learning with the use of Information and Communication Technologies.

In this regard, Esteban et al. (2018) expressed about adapting to the changes that today's society demands, where the master class has already been left behind to give way to the class with technological, audiovisual and interactive media, which has greater prominence.

On the other hand, the training of students must be in accordance with today's society, in an integral way where it enables them to apply professional knowledge, in a responsible and regulated way under the respect, not only to the procedures and adequate decision-making processes, but also within the respect for values and attitudes proper to the profession such as professional ethics, as expressed by González and Méndez (2017), elements with which the researchers agree.

In conclusion, the actions applied as a result of the didactic strategy contributed to improve the TLP in the students of the first year of the Regular Day Course of the Geography career of the Faculty of Secondary Education of the Central University "Marta Abreu" of Las Villas.

## REFERENCES

Alcántara, M.J., Medina, Q.S. (2019). El uso de los itinerarios didácticos (SIG) en la educación ambiental. Enseñanza de las ciencias (Innovaciones didácticas).

Córdoba, España. 37 (2): 173-188. <https://doi.org/10.5565/rev/ensciencias>

Armiñana, G.R., Fimia, D.R., Castillo, F.Y., López, P.R.T., Fernández, P.J.A., Iannacone, J. (2021). *Lissacathina fulica* (Bowdich, 1822) (Mollusca: gastropoda: Stylommatophora: Achatinidae), amenaza los ecosistemas cubanos y la salud humana. *Neotropical Helminthology* 15(1): 41-55.

Armiñana, G.R., Castillo, F.Y., Mesa, C.N., Fimia, D.R., Leyva, H.J., Iannacone, J., Durán, F.Y., Fábrega, O.G. (2020). Nueva concepción didáctica para el proceso de enseñanza - aprendizaje de la Zoología de los cordados. *PAIDEIA XXI* 10: 33-57.

Batllore, R. (2011). Enseñar y aprender el espacio geográfico” en Santisteban, A. & Pagès, J. (Coords.) *Didáctica del Conocimiento del Medio Social y Cultural en la Educación Primaria. Síntesis*, Madrid, p. 249-268.

Botella, N.A.M., Hurtado, A. (2016). Formación del profesorado en innovación educativa. La experiencia de L’Hort 2.0. *Aulas virtuales: fórmulas y prácticas*. Universidad de Valencia, España. 13 pp.

Campo, P.B., Rodríguez, P.L.A., Colomer, R.J.C. (2016). Enseñanza de itinerarios geográficos para el estudiante de magisterio; propuestas de aprendizajes autónomo en el aula universitaria y en el Geo foro iberoamericano. En *Nativos digitales y Geografía en el siglo XXI; educación geográfica y estilos de aprendizajes* Universidad de Valencia y Colombia. 142 pp.

Charbonet, M.E. (2009). Mejoramiento del desempeño profesional pedagógico para la educación ambiental de los profesores de ciencias naturales de los preuniversitarios. Una estrategia pedagógica [Tesis de doctorado, Universidad de Ciencias Pedagógicas “Enrique José Varona”, La Habana, Cuba]. 228 pp.

Chávez, L.E.D., Chávez, M.N., Villafructe, M.J.A. (2019). Estrategia Pedagógica de educación ambiental en función de una vida saludable en la Universidad de Guayaquil. *Revista Conrado* 15(70): 63-66 <http://conrado.ucf.edu.cu/index.php/conrado>

De la Calle, C. (2012). La enseñanza de la Geografía ante los nuevos desafíos ambientales, sociales y territoriales. *La educación geográfica digital*. Universidad de Valladolid, España. Grupo de Didáctica de la Geografía de la Asociación de Geógrafos Españoles. 123-137 pp. Disponible en: <http://www.age-didacticageografia.es>

Domínguez, D., Álvarez, J., Gil-Jaurena, J. (2016). Analítica del aprendizaje y Big. Data: heurísticos y marcos interpretativos. *International Journal of Applied Ethics* 22: 87-103. <http://www.dilemata.net/revista/index.php/dilemata/aeticle/view/41200042/450>

Esteban, S.N., Rivas, R.B., Sánchez, M.R., Cáceres, T.C. (2018). Itinerarios formativos para docentes URJC online. Centro de Innovación en Educación Digital: URJC online. Universidad Rey Juan Carlos Móstoles, Madrid,



- España. 157-159 pp. Disponible en: <http://riull.ull.es/xmlui/handle/915/8045>
- Évora, O.E. (2011). La educación ambiental para el desarrollo sostenible en la formación inicial de los profesionales de la educación en el proceso de enseñanza-aprendizaje de la física. [Tesis en opción al grado científico de Doctor en Ciencias Pedagógicas, Universidad de Ciencias Pedagógicas "Félix Varela", Villa Clara, Cuba]. 212 pp.
- Expósito, M., Armiñana, R., Fimia, R., Gómez, M. A., Artiles, L. A., Olivera, D., Iannacone, J., Carballo, B.M. (2021). Vinculando a los escolares de la enseñanza media al conocimiento de las especies exóticas invasoras de origen vegetal en Cuba. *The Biologist (Lima)* 19(2): 175-185.
- Fernández, N.F.A., Lima, M.S. (2017). La educación a distancia en entornos virtuales de enseñanza aprendizaje. *Reflexiones didácticas. Atenas* 3(39): 31-47. Disponible en: <https://www.redalyc.org/articulo.oa?id=478055149003>
- García, P. (2012). Posibilidades de la plataforma IDERioja para el análisis de la ciudad de Logroño. La educación geográfica digital. Universidad de La Rioja, España. Grupo de Didáctica de la Geografía de la Asociación de Geógrafos Españoles. 65-72 pp. Disponible en: <http://www.age-didacticageografia.es/>
- García, R. (2012). Los desafíos futuros de la población mundial: materiales para su análisis y discusión en el aula. La educación geográfica digital. Universidad de Barcelona y Cantabria, España. Grupo de Didáctica de la Geografía de la Asociación de Geógrafos Españoles. 131-157 pp. Disponible en: <http://www.age-didacticageografia.es/>
- Garrido, P.F. (2014). Topofilia, paisaje y sostenibilidad del territorio. *Enrahonar. Quaderns de Filosofia* 53. Andalucía. España. 63-75 pp.
- Garrison, D. (2016). ELearning en el siglo XXI. Un marco comunitario de investigación y la práctica. New York, Estados Unidos. *Revista Iberoamericana de Educación a Distancia* 23(1): 09-28. Disponible en: <http://dx.doi.org/10.5944/ried.23.1.25495>
- González, M. (2017). Desarrollo de un Prácticum virtual: fundamentos y clases. *Revista Prácticum* 2(2): 50-68.
- Hernández, F.M., Cárdenas, G.M., Armiñana, G.R., Fimia, D.R., Iannacone, J. (2021). El trabajo de campo: una herramienta para la enseñanza del patrimonio y la historia local. *Biotempo* 18(1): 21-35.
- International Geographical Commission, Commission on Geographical Education. (1992). International Statement on Geographical Education. 160-162 pp. Available at: <http://www.raco.cat/index.php/RevistaGeografica/article/viewFile/46086/56892>.
- International Geographical Union. Commission on Geographical Education. (2000). International Declaration on Geographic Education for Cultural Diversity. 195-200 pp. Available at: [http://www.agegeografia.es/v2/diversidad\\_cultural.php?TB\\_iframe=true&height=430&width=70](http://www.agegeografia.es/v2/diversidad_cultural.php?TB_iframe=true&height=430&width=70)
- International Geographical Union. Commission on Geographical Education. (2007). International Declaration on Geographical Education for Sustainable Development. 217-240 pp. Available at: <http://www.igu-cge.org/Charterspdf/spanish.pdf>
- Liceras, A. (2018). Los itinerarios didácticos en la enseñanza de la Geografía. Reflexiones y propuestas acerca de su eficacia en la Educación. *Revista de la Universidad de Granada, España. Universidad, Escuela y Sociedad* 5: 66-81.
- Lin, C., Liu, K. (2017). A study of the effects of digital learning on learning motivation and learning outcome. *Eurasia Journal of Mathematics, Science and Technology Education* 13(7): 3553-3564.
- Maderuelo, J. (2007). El paisaje. Génesis de un concepto. Editorial Abada. Madrid, España. ISBN 9788-4962-5856-3. 344 pp.
- Marrero, R D., Santos, I.E. (2018). Aproximación a los valores del educador ambiental. *Revista Humanidades Médicas* 19(1): 160-179.
- Marrón, G.M.J. (2007). Desarrollo sostenible, globalización y educación en valores ambientales desde la geografía. Una propuesta metodológica en el marco europeo de Educación Superior. En Marrón Gaite, Mª J. Salom Carrasco, J. & Souto González, X.M. (Eds.) *Las competencias Geográficas para la educación ciudadana*, Valencia: Grupo de Didáctica de la AGE- Universidad de Valencia, p. 133-146.
- Marulanda, N., Rojas, M.D. (2019). Ética en Instituciones de Educación Superior para la Construcción de Relaciones de Confianza con Grupos de Interés (Stakeholders). *Información Tecnológica* 30(3): 269-276. Disponible en: <http://dx.doi.org/10.4067/S0718-07642019000300269>
- Migon, P. (2017). View point geosites-values, conservation and management issues. *Proceedings of the Geologists' Association*, volume Elsevier Ltd 128, Issue 4. 518-522 pp. Disponible en: <https://doi.org/10.1016/I.pgeola.2017.05.007>
- Orfa, R.M., Garzón, R.L., Pinzón, M.L., Salamanca, A.L., Pabón, L.C. (2019). Estrategias pedagógicas como herramienta educativa: la tutoría y el proceso formativo de los estudiantes. *Revista Iberoamericana de Educación* 50 (3): 1-16. Disponible en: <https://doi.org/10.35362/rie5031869>
- Oyarce, A. (2018). Caracterización, valoración y promoción de la geodiversidad del sendero Sierra Velluda en el parque nacional Laguna del Laja mediante la creación de un itinerario geológico virtual utilizando Google Earth y código QR. Facultad de ingeniería de la Universidad "Andrés Bello", Concepción, Chile.
- Palacio, J., Sánchez, J., Schilling, M. (2016). Patrimonio

geológico y su conservación en América Latina: situación y perspectivas nacionales. Instituto de Geografía de la Universidad Nacional Autónoma de México. Colección: Geografía para el siglo XXI. Serie: Libro de investigación, No 18. 659-668 pp

Sánchez, R. (2020). Proyecto de Innovación docente para la educación patrimonial en Madrid. Innovación educativa en la sociedad digital. Madrid, España. s/p. Disponible en: <https://www.ucm.es/proyectos-de-innovación-2019-20>

Seoul Declaration of 2000. In: International Geographical Union. Commission on Geographical Education. 2000.

International Declaration on Geographic Education for Cultural Diversity. 195-200 pp. Available at: [http://www.agegeografia.es/v2/diversidad\\_cultural.php?TB\\_iframe=true&height=430&width=70](http://www.agegeografia.es/v2/diversidad_cultural.php?TB_iframe=true&height=430&width=70)

Vilarrasa, A. (2003). Salir del aula. Reapropiarse del contexto. Revista Iberoamericana 36: 13-25.

Villamandos, F., Gomera, A., Antúnez, M. (2019). Conciencia ambiental y sostenibilización curricular, dos herramientas en el camino hacia la sostenibilidad de la Universidad de Córdoba. Revista de Educación Ambiental y Sostenibilidad 1(1): 13-25. Doi: 10.25267/Revista\_educación\_ambiental\_sostenibilidad. 2019. v1.i1.1301

# Localization of neuropeptide Y(NPY) Immunoreactivity in the Proximal and Distal Intestinal Region of Teleost Fish, *Notopterus Notopterus*

Tulika A. Khadse<sup>1</sup>, Yashashree. A. Gadhikar<sup>2</sup>, Trupti S. Khedkar<sup>3\*</sup> and Sudhir G. Chird<sup>4</sup>

<sup>1</sup>Department of Zoology, Jagadamba Mahavidyalaya, Achalpur city(MS), India.

<sup>2</sup>Department of Zoology, Govt. Vidarbha Institute of Science and Humanities, Amravati (MS), India.

<sup>3</sup>Department of Zoology, Nabira Mahavidyalaya, Katol 441302, Dist. Nagpur (MS), India.

<sup>4</sup>Department of Zoology, Arts Commerce and Science College Maregaon District Yavatmal (MS), India

## ABSTRACT

The purpose of the present study was to investigate the immunocytochemical aspects of the intestine of the teleost fish *N.notopterus*. The distribution and relative frequency of the endocrine cells in the intestine were studied immunocytochemically using Streptavidin biotin peroxidase complex method. The aim was analyse the distribution and localization of immunoreactivity in the intestine of *Notopterus notopterus* against the antisera Neuropeptide Y (NPY). The samples were taken from the proximal and distal intestine. The NPY immunoreactive endocrine cells were found in the distal intestine at low frequencies than proximal intestine. These immunoreactive cells were distributed among more in mucosa epithelium and very low in the lamina propria and submucosa of intestine of *N.notopterus*.

**KEY WORDS:** IMMUNOCYTOCHEMISTRY , INTESTINE, TELEOST, *NOTOPTERUS NOTOPTERUS*

## INTRODUCTION

The intestine in the carnivorous fish is shorter than that of omnivores and herbivores. The intestine of the fish plays an important role in digestion absorption of dietary nutrients and also involved in immunological functions (Cao et al.2011; Salias et al .2011).Morphometric features are important in identifying the fish species and their interaction in various habitats like freshwater, rivers and seas (Akombo et al.2011). Gastrointestinal tract (GI) of fishes can be divided into four topographical regions headgut, foregut, midgut and hindgut. Although in some cases there is no clear morphological distinction between the midgut and hindgut, morphometric characteristic of intestine of *N. notopterus* was performed .

The endocrine cells secreted the gastrointestinal hormones which are distributed throughout the mucosa of gastrointestinal tract(GI). A great variety of endocrine type secretions are generated in the mucosa of the gastrointestinal tract of fish. Peptides are the name given to these secretions. For the last 15 years more than 45 gastrointestinal

peptides have been determined in studies on fish GI tract. Neuropeptide Y(NPY) was isolated and sequenced for the first time from swine brain (Tatemoto, 1982, Bjorgen and Koppang 2021).

It has been characterized as one of the most highly conserved neuroendocrine peptide through out the evolution. NPY has been involved in the regulation of a wide range of physiological central effects like the control of appetite, body weight homeostasis, the modulation of reproductive processes. This paper examines the distribution and localization of endocrine cells in the proximal and distal intestine of teleost fish *N. notopterus* using immunocytochemical study.

The Asian Knifefish (*N. notopterus*) is a carnivorous and predatory fish. Intestine of the carnivorous fishes is short or more or less straight, because meaty foods can be digested more readily than vegetable ones. This fish accepts most kind of live frozen food. Some specimen accepts pellets and dry food. It is available in India, Pakistan, Bangladesh, Nepal, Thailand, Malaysia, and Indonesia.. Body is highly compressed, dorsal and ventral profile almost equally convex. Hence in the present investigation, immunocytochemical were carried out.

**Article Information:**\*Corresponding Author: [truptikhedkar9@gmail.com](mailto:truptikhedkar9@gmail.com).

Received 25/04/2023 Accepted after revision 15/06/2023

Published: June 2023 Pp- 115-119

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.8>

## MATERIAL AND METHODS

**Sample collection:** Twelve live adult *N. notopterus* of either sex with body length ranging from 19-22 cm, weighing 80-100 gm were collected from fresh water pond called as Chatri Talab, Amravati, Maharashtra, India ( 200 56' N, 770 47' E). Fishes were anaesthetized with 2-phenoxy ethanol (1:2000) for 10-15 mins in the anaesthetic tank. After being anaesthetized, the fish was scarified by transcardial perfusion by ice cold Phosphate Buffer Saline (PBS, pH 7.45) solution and processed for immunocytochemical studies.

**Immunocytochemical Study:** For immunocytochemical study, the intestine was excised, separated into proximal

and distal region and fixed in Bouins fluid for 24 hours. After fixation, fixative was removed by using cold sucrose solution prepared in PBS. The tissue were cryoprotected in 10% sucrose for 2 hours, 20% sucrose for 2 hours and 30% sucrose for overnight at 4<sup>o</sup> C. Cryoprotected tissues were embedded in 15% Polyvinyl Pyrrolidone Phosphate (PVP) on cryoprotected peg. 12-20 micron thick sections were obtained by cryostat -29<sup>o</sup> C and spread on the slides subbed with poly L-lysine (sigma). The slides were preserved in deep freezer and proceed for immunocytochemical staining. Immunocytochemical staining was carried out by using Streptavidin-biotin peroxidase complex method-The slides were transferred to humidity chamber and sections were washed in PBS (3×5 min) and then washing with Triton X 0.5 % (20 min).

**Table 1. Morphometric characteristic of *Notopterus notopterus***

Notopterus notopterus	Fish total body length(cm)	Fish standard length(cm)	Digestive tract length(cm)	Intestine length (cm)	Weight (gm)
Mean ± SD	20 ± 1	19.5 ± 1	6.333 ± 0.3	5 ± 0.5	90 ± 10

**Table 2. Morphological different cells in proximal and distal intestine of *Notopterus notopterus***

Types of cells	Proximal per villi/per sec	Distal per villi/per sec
Round	++	++
Tadpole	+++	++
Ovale	++	+
Triangular	++	+
Sac like	+	+
Rod like	+	+
Elongated	+	+
Cucurbit like	+	+

+ -- average, ++ -- moderate, +++ -- maximum

**Table 3. Meannumber of endocrine cells per intestinal folds (Mean ± SD) in *N. notopterus* (120 intestinal fold were examine for NPY antisera)**

NPY like immunoreactivitiy from 120 folds divided into 3 groups		
Intestinal region ( <i>N. notopterus</i> )	Proximal	Distal
Mean ± SD	68.66 ± 1.52752	35.6 ± 6.4291005

The sections were treated with blocking solution 1% BSA (30 min).3)The sections were incubated with primary antibody (NPY Sigma, Cat # N-9528) at dilution of 1:2000 with PBS containing 0.3 % Triton X-100 1% BSA and then

washed with PBS(3×5min).4) The sections were incubated with secondary antibody (Biotin IgG complex, Sigma, Cat #BA2) for 2 hours and then washed in PBS (5 min).4) The sections were incubated with Streptavidin peroxidase conjugate at dilution (1:100) for 2 hours and then washed in PBS(3×5 min) and then double distilled water (2×5 min).5) For visualization of reaction 3-Amino -9-ethylcarbazole (AEC) solution were applied on sections for 5-10 min until reddish brown colour appear .6)Then slides were washed in double distilled water (2×5 min) and mounted with glycerol gelatin.7) The sections were examined with light microscope and photograph were taken.

## RESULTS AND DISCUSSION

The sections obtained from the proximal and distal intestinal tissue of *N. notopterus* were formed by mucosa, submucosa, muscularis and serosa layer which is similar to the basic organization of intestinal wall of other teleost fishes according to the histological studies (Fig.1).

In the present investigation, essentially NPY immunoreactive (IR) endocrine cells were observed in the epithelium of intestinal mucosa and few existed in the lamina propria and submucosa of the *Notopterus notopterus*. Density of NPY IR endocrine cells were more pronounced in the proximal intestine than the distal intestine. All NPY IR cells were stained dark brown colour. Two types of NPY IR endocrine cells were observed in both proximal and distal intestine –open type and closed type. Numerous open type tadpole like NPY IR endocrine cells having long cytoplasmic processes extending to the lumen and basal process extending to the lamina propria were observed in the mucosal epithelium (Fig.2,4).

Open type rod like,(Fig2),Spindle shaped(Fig.6),Pyramid like with short cytoplasmic process extending to the lumen and basal broad process extending to the



lamina propria,(Fig.,6),elongated (Fig.5), Piriform like(Fig.6),Triangular like (Fig.6).NPY IR endocrine cells were observed in the intestinal mucosal epithelium. Open type Sac like NPY IR endocrine cells with apical cytoplasmic processes extending to the lumen, the basal part narrow extending to the lamina propria & middle part of the cell body to be broader are observed in the apical, middle and basal part of the intestinal mucosal epithelium. (Fig.3)Closed type oval like and round like(Fig.7) NPY IR endocrine cells were observed in the intestinal mucosal epithelium.

**Figure 1: Transverse section through proximal intestine showing different layers Mucosa (MU), Sub mucosa (SM), Muscularis (MU), Longitudinal Muscle Layer (LML),Circular Muscle Layer (CML), Lamina propria (Lp) and Lumen(Lu) (X 100).**


**Figure 2: Magnified view of open type tadpole like and rod like NPY IR endocrine cells.**

**Figure 3: Magnified view of open type sac like and pyramid like NPY IR endocrine cells.**

**Figure 4: Open type triangular and tadpole like NPY IR endocrine cells.**

**Figure 5: Magnified view of open type triangular and elongated NPY IR endocrine cells (X400).**

**Figure 6: Magnified view of open type pyramid like and spindle shaped open type NPY IR endocrine cell (X1000).**



Tadpole like open type NPY IR endocrine cells with one long cytoplasmic processes (fig.8), two long cytoplasmic processes(fig.11) extended to the lumen and basal process extending to the lamina propria are observed in the intestinal mucosal epithelium.Open type elongated, Spindle shaped and Piriform like (fig.8), Cucurbit like(fig.9) NPY IR endocrine cells were observed in the intestinal mucosal epithelium Open type triangle like NPY IR endocrine cells with long cytoplasmic process extending to the lumen and basal process to the lamina propria in the intestinal mucosal epithelium.(fig.8).

Sac like open type IR endocrine cells were observed in the intestinal mucosal epithelium (fig.10) and in sub-mucosa. Closed type oval like NPY IR endocrine cells were observed

in the lamina propria. Closed type round like NPY IR endocrine cells were observed in the intestinal mucosal epithelium (fig.12).

**Figure 7: Magnified view of closed type round NPY IR endocrine cells.**

**Figure 8-12: Transverse section through distal intestine of *Notopterus notopterus* showing NPY immunoreactive (IR) endocrine cells.**


**Figure 8: Open type tadpole like, piriform, triangular, spindle shaped NPY IR endocrine cell (X400).**

**Figure 9: Open type cucurbit like NPY IR endocrine cell (X400).**

**Figure 10: Magnified view of sac like NPY IR endocrine cell (X1000).**

**Figure 11: Magnified view of elongated NPY IR endocrine cell with two long cytoplasmic processes (X1000).**

**Figure 12: Closed type round NPY IR endocrine cell (X400).**



Studies concerning regulatory peptides in fish intestine have been carried out since last several years. But intestinal NPY like endocrine cells have not been previously documented in species of *N.notopterus*. In the present, we have studied the distribution and localization of NPY regulatory peptide from intestine of the fish *N.notopterus*. In the present study immunoreactive endocrine cells were detected more in the mucosal epithelium of proximal intestine than distal intestine. Similar results of NPY were reported in *Anguilla anguilla* (Domeneghini et al.2000 ; Youson et al.2001) and with variation to above results less amount of NPY immunoreactive cells were observed in the intestine of *Salmotrutta* (Dezfuli et al. 2000) and *Stizostedianlucio perca* (Ozen et al.2010).

Open and closed types of endocrine cells were observed in the intestine of *N.notopterus* in the present study. In open type, hormones were carried to the gut lumen (lumen endocrine)by long apical cytoplasmic processes (Liu et al. 2003; Pan and Fang 1995) sand in closed type cells it is covered by lumen epithelia (Liu et al.2003 ;Talatar and Simsek 1993).The endocrine cells of the gastrointestinal tract were different in the relative frequency of regional distribution and cell types between species (Leet al.2004;



Pan et al. 2000;Reverter et al.2001; Salianas et al.2011 Bjorgen and Koppang 2021).

## CONCLUSION

The distribution and relative frequency of immunoreactive endocrine cells in *N. notopterus* are similar to that of other teleost fishes. However few difference in morphology of immunoreactive endocrine cells were observed. From morphometric analysis of the intestine of *N. notopterus* it was confirmed that intestine is relatively shorter in length as compared to other teleost fishes. NPY may involved in many physiological functions like regulation of food intake and modulation of reproductive processes. These studies could be extended to study for the immunology of this fish.

**Authors' Contribution:** YG, TK and SC, were responsible for conceptualization and design of the research. TUK, YG were responsible for specimen collection. SC and TK were responsible for result analysis and interpretation. TUK and YG were responsible for laboratory techniques. All authors contributed equally in literature research, manuscript preparation, editing and review.

**Statement of Ethics:** The study was approved by the Institutional Animal Ethics Committee ((IAEC) Of Government Vidharbha Institute of Science and Humanities, Amravati.

**Declaration of Conflicting Interests:** The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**Funding:** The authors did not receive any financial support for the research, authorship and publication of this article.

## REFERENCES

- Akombo, P.M., Atile, J.I., Adikwu, I.A., &Araoye, P.A. (2011). Morphometric measurements and growth patterns of four species of the genus *Synodontis* (Cuvier, 1816) from lower Benue river, Markurdi, Nigeria. *International Journal of Fisheries and Aquaculture*, 3(15), 263-270.
- Akombo PM, Atile JI, Adikwu IA, Araoye PA. Morphometric measurements and growth patterns of four species of the genus *Synodontis* (Cuvier, 1816) from Lower Benue River, Makurdi, Nigeria. *International Journal of fisheries and Aquaculture*. 2011 Dec 23;3(15):263-70.
- Bjorgen H and Koppang EO (2021). Anatomy of teleost fish immune structures and organs *Immunogenetics*. 2021; 73(1): 53–63. Published online 2021 Jan 11. doi: 10.1007/s00251-020-01196-0.
- Cerdá-Reverter JM, Martínez-Rodríguez G, Zanuy S, Carrillo M, Larhammar D. Molecular evolution of the neuropeptide Y (NPY) family of peptides: cloning of three NPY-related peptides from the sea bass (*Dicentrarchus labrax*). *Regulatory peptides*. 2000 Nov 24;95(1-3):25-

34

- Cao XJ, Wang WM, Song F. Anatomical and histological characteristics of the intestine of the topmouth culter (*Culter alburnus*). *Anatomia, Histologia, Embryologia*. 2011 Aug;40(4):292-8.
- Cerdá-Reverter JM, Martínez-Rodríguez G, Zanuy S, Carrillo M, Larhammar D. Molecular evolution of the neuropeptide Y (NPY) family of peptides: cloning of three NPY-related peptides from the sea bass (*Dicentrarchus labrax*). *Regulatory peptides*. 2000 Nov 24;95(1-3):25-34.
- Dezfuli, B.S., Arrighi, S., Domeneghini, C., &Bosi, G. (2000). Immunohistochemical detection of neuromodulators in the intestine of *Salmotrutta L.* naturally infected with *Cyathocephalustruncatus Pallas* (Cestoda). *Journal of Fish Diseases*, 23, 265-273.
- Dezfuli, B.S., Arrighi, S., Domeneghini, C. and Bosi, G., 2000. Immunohistochemical detection of neuromodulators in the intestine of *Salmo trutta L.* naturally infected with *Cyathocephalus truncatus Pallas* (Cestoda). *Journal of Fish Diseases*, 23(4), pp.265-273.
- Domeneghini, C., Radaelli, G., Arrighi, S., Macarello, F., &Vegetti, A. (2000). Neurotransmitters and putative neuromodulators in the gut of *Anguilla anguilla* (L.). Localizations in the enteric nervous and endocrine system. *European Journal of Histochemistry*, 44(3), 295-306.
- Lee, J.H., Ku, S.K., Park, K.D., & Lee, H.S. (2004). Immunohistochemical study of the gastrointestinal endocrine cells in the Korean aurchaperch. *Journal of Fish Biology*, 65, 170-181.
- Liu, Y., Tytgat, G.N.G., Xiao, S.D., & Ten-kate, F.J.W. (2003). Gastric endocrine cells. *Chinese Journal of Digestive Diseases*, 4, 160-167.
- Ozen, R.M., Kenan, Cinar., & Nurgul, Senol. (2010). Endocrine cells in the gastrointestinal tract mucosa of Zander (*Stizostedionlucioiperca*). *Kafkas University Veterinary Faculty Journal*, 16(Suppl-B), S339-S345.
- Pan, Q.S., & Fang, Z.P. (1995). Present progress in the study of the APUD cells in gastro-entero-pancreatic endocrine system of the fishes. *Acta Hydrobiologica Sinica*, 19, 275-282.
- Pan, Q.S., Fang, Z.P., & Zhao, Y.X. (2000). Immunocytochemical identification and localization of APUD cells in the gut of seven stomachless teleost fishes. *World Journal of Gastroenterology*, 6(1), 96-101.
- Salianas, I., Yang, A. Z., &Sunyer, O. (2011). Mucosal immunoglobulins and B cells of the teleost fish, Dve. *Comparative Immunology*, 35, 1346-1365.
- Talatar, H., &Simsek, H. (1993). Hacettepeuniversitesi tip fakultesiichastalıkları ABD gastroenteroloji: unitesiHekimlerYayınBirligi, Ankara, pp. Gastroenteroloji, 551.
- Tatemoto, K. (1982a). Neuropeptide Y: Complete amino-acid sequence of the brain peptide. *Proceedings of the*

Khadse et al.,

National Academy of Sciences of the United States of America, 79, 5485-5489.

Youson JH, Al-Mahrouki AA, Naumovski D, Conlon JM.

The endocrine cells in the gastroenteropancreatic system of the bowfin, *Amia calva* L.: an immunohistochemical, ultrastructural, and immunocytochemical analysis. *Journal of Morphology*. 2001 Dec;250(3):208-24.

# Ecology of Biofouling Phytoplankton in Chinnamuttom Harbour Waters Southeast Coast of India

P. Nithya\*<sup>1</sup>, B. Dhanalakshmi<sup>2</sup> and P. Santhanam<sup>3</sup>

<sup>1</sup>PG and Research Department of Zoology, Vivekanandha College of Arts and Sciences for Women, Tiruchengode, Namakkal -637205, Tamil Nadu, India.

<sup>2</sup>PG and Research Department of Zoology, Nirmala College for women, Coimbatore – 641018, Tamil Nadu, India.

<sup>3</sup>Marine Planktonology and Aquaculture Laboratory, Department of Marine Science, School of Marine Sciences, Bharathidasan University, Tiruchirappalli – 620 024, Tamil Nadu, India.

## ABSTRACT

Phytoplankton are the primary source of food, attributed most zooplankton communities and larval stages, and many meroplankton such as bivalves, crustaceans, and fishes. In the present study, phytoplankton samples were collected from surface water and substratum of the ship hull during the period of June 2015 to May 2016. Population density recorded from water and substratum of the ship hull ranged from  $1.5 \times 10^4$  to  $3.4 \times 10^4$  cells/L. Shannon-Wiener diversity index ( $H'$ ) of station I and station II ranged from 5.448 to 6.134 bits/ind. Simpson's richness range was 0.973 to 0.984 at station I and station II. Pielou's Evenness index ( $J'$ ) was found to be from 0.950 to 0.979 and Chlorophyll 'a' range was 0.277 to 1.326 mg/m<sup>3</sup> in the station I and station II. During this entire study period, 74 species of phytoplankton and 54 species of phytoplankton in station II have been recorded. Data of the present study show that significant differences occurred in over all percentage composition of phytoplankton species, both in water and substratum. In water samples, recorded data showed that 85% were of *Bacillariophyceae*, 8 % of *Chlorophyceae* and, 7% of *Dinophyceae*. On the other hand, 75 % *Bacillariophyceae*, 7 % *Chlorophyceae*, 16% *Dinophyceae* and 2% *Cyanophyceae* were recorded in the substratum. The marine biofouling phytoplankton density, diversity, richness, evenness and chlorophyll 'a' were recorded in their maximum levels during summer season, and minimum were obtained in the monsoon. The diatoms were the predominant groups among the phytoplankton. Among the diatoms, species like *Nitzschia longissima*, *Coscinodiscus centralis* and *Fragilaria* sp. were record almost in all the seasons in the present study.

**KEY WORDS:** CHLOROPHYLL, PHYTOPLANKTON, POPULATION DENSITY, SUBSTRATUM, WATER

## INTRODUCTION

Phytoplankton by virtue of drifting habits in illuminated waters of our planet and short turnover periods link the oceanic food web and its trophic marine structures. These are extremely diverse energy sources, distributed across major eukaryotic lineage, in food web of the aquatic ecosystems, playing a key role in planktonic communities, ecological structure, function, and the dynamic fate of the biota, (Mattei and Scardi 2021). In marine ecosystems, the phytoplankton (<1µm to ≈1mm length in diameter) represent at least five eukaryotic phyla, which are one of the world's

simplest, basic and dominant key producers of the primary marine food chain, (Sun et al. 2000).

Anthropogenic discharge of nutrient load to sea waters, increase of global carbon emission and their mix with wide oceanic surface water, multitude seasonal hydrogeochemical changes impacts on the primary producers. Hence about 45% of global net primary productivity plays a critical role in biogeochemical functions and production of climatically active gases. The earlier estimate of Geider et al. (2014) indicates that, the extensive range in size and desperate genetic diversity challenges the phytoplankton quantification and characterization, (Kim and Kim 2021). Species composition, density, abundance, richness and primary productivity of phytoplankton vary in coastal geography, which is liable to physiological and hydro-

**Article Information:**\*Corresponding Author: [neethujeni@gmail.com](mailto:neethujeni@gmail.com)

Received 15/05/2023 Accepted after revision 25/06/2023

Published: June 2023 Pp- 120-125

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.

Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.9>

biological fluctuation (Al-amri et al. 2020). Physicochemical values, phytoplankton and zooplankton species, and their composition, abundance, seasonal variations have been studied in various regions of Indian coastal waters (Vajravelu et al. 2018).

Significantly, the phytoplankton are the major contributors in absorbing atmospheric carbon than the terrestrial plants, also, these plankton takes a major role in global carbon dioxide sequestration, oxygen production, nutrient cycle, and carbon fixation and contributes to global warming and the existence of aquatic lifeforms (Paffenhofer, 1993). Consequently, assessing the species composition and their density diversity is necessary to assess a healthy ecosystem.

Changes in phytoplankton ecology and diversity are driven by several major unresolved conceptual challenges, perhaps the foremost of which is the marine pollution of fouling. Analysis of diversity and ecology of phytoplankton has largely benefited many research fields in biology as it is entering a new era with the advent of fostering the improvement of the predictive tool of phytoplankton distribution at global scales. Hence, the present study was focused on diversity, density, species composition, richness, and evenness of phytoplankton community in the fishing harbor environment concerning the prevailing hydrographical conditions of Chinnamuttam area which harbours waters of southeast coast of India.

## MATERIAL AND METHODS

**Sampling program:** A total of four sampling sessions were conducted during the study period i.e. June 2015 to May 2016, between the hours of 06.00 hrs and 12.00 hrs.

**Collection of samples:** The phytoplankton sampling was done using standard Indian Ocean hand net (48 $\mu$ m) by towing horizontally in water surface for 30 minutes, as well the sampling from substratum of ship hull was collected by scraping using sterile spatula (Manickam et al. 2017).

**Preparation of the sample:** 10ml/L Lugol's iodine was added into samples and kept standing for 24hrs to ensure complete sedimentation followed by centrifugation if necessary. The supernatant liquid was removed using pipette and the sample was further concentrated up to 10-100ml depending on the number of plankton.

**Fixation and Preservation of samples:** The formalin was added in the ratio of 5-10 to 90 parts(v/v) and the bottle was inverted for fine dispense. The pH of the solutions was maintained at 7.6 – 8.3.

**Observation of phytoplankton:** The observation of phytoplankton was done by exposing the concentrated sample to the Light microscope (COSLAB). The phytoplankton observed under microscope were taken with the support of a digital camera (MDCE-5C).

**Identification of phytoplankton:** The identification of phytoplankton was carried out with the help of standard

books, identification manuals and classical works of Subramanyan (1946); Subramanyan (1968); Venkataraman (1939); Tomas (1997) and Mitra et al. (2004).

**Plankton enumeration using Sedgwick-Rafter Cell method:** One ml of well mixed plankton samples was poured into the Sedgwick-Rafter cell using graduated pipette and uniformly spread as thin layers. The plankton numbers were counted by selecting squares randomly. The plankton density was estimated using the following formula. Calculation  $N = n \times v/V$  Where, N = Total number of plankton cells / L, n = Average number of plankton cells in 1 ml of sample v = Volume of plankton concentrate, V = Total volume of water filtered (L).

**Diversity Analysis of plankton samples:** The diversity indices (DI) of phytoplankton in Chinnamuttom harbour was calculated using Shannon Weiner's Index (H'), Simpson richness index (D') and Evenness index (E'), which was calculated by using PAST– Palaeontological Statistics Ver. 2.00 software packages.

**Chlorophyll- 'a':** 250 ml of collected water sample was filtered through (0.45 diameter GF/ C filter paper) Millipore filtering unit. Then filtered paper was soaked in acetone and kept in dark for 20–24 hours under refrigeration to extract the chlorophyll. The centrifuged concentration of chlorophyll extracts measured spectrometric at 665nm (APHA, 2010).

## RESULTS AND DISCUSSION


**Species composition:** The phytoplankton samples collected from Chinnamuttom fishing harbour was analysed for the one-year period from 2015 to 2016. In the one year study period, totally 74 phytoplankton species were recorded, viz.. 57 species of Bacillariophyceae, 9 species of Dinophyceae and 8 species of Chlorophyceae in water sample, whereas in the substratum of the ship hull 59 phytoplankton species were recorded. Among the total of 59 species, 45 Bacillariophyceae species, 8 Dinophyceae species, 4 Chlorophyceae species, and 2 species of Cynophyceae were noted. Similar, study was conducted recently in southeast coast of India and recorded five different classes (Vajravelu et al. 2017)

Among the recorded species of sampling stations (Bacillariophyceae, Dinophyceae, Cyanophyceae and Chlorophyceae) diatoms were found to be dominant contributor (75–85%) of the total phytoplankton. Especially, species such as *Nitzschia* spp., constituted the maximum numbers followed by *Navicula* spp., and *Coscinodiscus* spp. The domination in phytoplankton group was in the order: Diatom > Dinoflagellates > Green algae > Blue-green algae.


**Percentage composition:** Phytoplankton comprised of Bacillariophyceae (diatoms), Dinophyceae (Dinoflagellates), Chlorophyceae (green algae) and Cyanophyceae (blue green algae). The recorded phytoplankton number from the collected samples were composed of 85% by Bacillariophyceae, 8% Dinophyceae and 7% by Chlorophyceae, were recorded in water sample (Figure 1

and 2 ). All these groups were present in all four seasons. In substratum of the ship hull, the recorded compendial count of 75% Bacillariophyceae, 16% Dinophyceae, 7% Chlorophyceae and 2% Cyanophyceae (Figure 1). Therefore, totally 74 and 59 phytoplankton species were recorded in water and substratum of the ship hull in Chinnamuttom harbour respectively during the study period.

**Figure 1: Percentage composition of phytoplankton in Chinnamuttom Harbour water during June 2015-May 2016**



**Figure 2: Percentage composition of phytoplankton in Chinnamuttom Harbour substratum during June 2015 - May 2016**




**Population density:** Population density of phytoplankton was recorded from water and substratum of the ship hull, in the range between 15000 - 34000 cells/L. The minimum density was recorded during monsoon (15000 cells/L) in the substratum of the ship hull, while the maximum value (34000 cells/L) was noticed during summer in the water sample (Figure 3). In the present study, the phytoplankton population density was found to be high (15000 cells/L) during summer could be attributed to stability in hydrographical conditions, and minimum density (34000 cells/L) was obtained during monsoon due to low salinity and high turbidity due to rainfall (Senthilkumar et al. 2002). A similar increased number of phytoplankton species significant with high salinity increase in Bay of Bengal were reported (Rajasekar et al. 2005).


**Species richness:** An analysis of the phytoplankton species richness at Chinnamuttom harbour was noticed to be the range between 0.973 and 0.984. The minimum (0.973) richness during monsoon in substratum, and maximum

(0.984) was during summer in the water sample (Figure 5).


**Figure 3: Seasonal variations in phytoplankton density during June 2015 - May 2016**



**Figure 4: Seasonal variations in phytoplankton species diversity during June 2015-May 2016**



**Figure 5: Seasonal variations in phytoplankton species richness during June 2015-May 2016**




The richness index of species (0.973-0.984 cells/L) are between 0.973 and 0.984. In substratum, minimum species richness was 0.973 during monsoon, and the maximum was 0.984 during summer in the water. The recorded species richness variation was correlated with the recorded salinity values (Mani, 1992).

The species evenness: The evenness index was varied 0.950 to 0.979 cells/L in phytoplankton species. The minimum species evenness was recorded 0.950 cells/L during




monsoon in the substratum and the maximum 0.979 cells/L was observed during summer in

**Figure 6: Seasonal variations in phytoplankton species evenness during June 2015-May 2016**



**Chlorophyll ‘a’:** chlorophyll ‘a’ ranged between (0.277 to 1.326 mg/m<sup>3</sup>), the higher concentration of 1.067mg/m<sup>3</sup> was found in summer season. Significant high population of phytoplankton density during summer season was reported in Palk Bay coral reef region (Sridhar et al.,

**Figure 7: Seasonal variations in chlorophyll ‘a’ concentration during June 2015-May 2016**



2010). The 0.277 mg/m<sup>3</sup> concentration of low chlorophyll ‘a’ was obtained on monsoon season may be due to low phytoplankton production owing to high turbidity and low light availability due to high land drainage and cloudy sky (Kawabata et al. 1993, Godantaraman 2002, Rajasekar et al. 2005). The phytoplankton density, richness, composition, diversity, and evenness indices are significant to the variations in seasonal physicochemical condition variation (Figure 7).

**Table 1. List of phytoplankton species recorded in Chinnamuttom Harbour during June 2015 – May 2016**

S. No.	Species Name	Water	Substratum
<b>BACILLARIOPHYCEAE (Diatoms)</b>			
1	<i>Asterionellopsis glacialis</i>	+	+
2	<i>Bacteriastrium comosum</i>	+	+
3	<i>B. paradaxa</i>	+	+
4	<i>B. delicatula</i>	+	+
5	<i>Biddulphia aurita</i>	+	-
6	<i>B. sinensis</i>	+	+
7	<i>B. mobiliensis</i>	+	+
8	<i>Chaetoceros affinis</i>	+	+
9	<i>C. compressus</i>	+	+
10	<i>C. dydymus</i>	+	-
11	<i>C. longissimi</i>	+	+
12	<i>C. lorenzianus</i>	+	+
13	<i>C. messanensis</i>	+	+
14	<i>C. paruvianus</i>	+	+
15	<i>Coscinodiscus centralis</i>	+	+
16	<i>C. concinnes</i>	+	+
17	<i>C. gigas</i>	+	+
18	<i>C. jonesianus</i>	+	+
19	<i>C. marginatus</i>	+	+
20	<i>C. radiates</i>	+	+
21	<i>Diploneis robusta</i>	+	+
22	<i>Ditylum brightwelli</i>	+	+
23	<i>Fragilaria sp.</i>	+	+
24	<i>F. intermedia</i>	+	-
25	<i>F. oceanica</i>	+	+

26	<i>Gyrosigma balticum</i>	+	+
27	<i>Hemidiscus cuneiformis</i>	+	+
28	<i>H. hardmannianus</i>	+	-
29	<i>Hyalodiscus stelliger</i>	+	-
30	<i>Lauderia annulate</i>	+	-
31	<i>Leptocylindrus danicus</i>	+	-
32	<i>Navicula forcipate</i>	+	+
33	<i>N. granulate</i>	+	-
34	<i>N. longa</i>	+	+
35	<i>Nitzschia Closterium</i>	+	+
36	<i>N. longissimi</i>	+	+
37	<i>N. paradoxa</i>	+	+
38	<i>N. sigma</i>	+	+
39	<i>N. sigmoidea</i>	+	+
40	<i>Odontella mobiliensis</i>	+	+
41	<i>O. sinensis</i>	+	+
42	<i>Planktoniella sol</i>	+	+
43	<i>Pleurosigma angulatum</i>	+	+
44	<i>P. elongatum</i>	+	+
45	<i>Rhizosolenia setigera</i>	+	-
46	<i>R. robusta</i>	+	-
47	<i>R. setigera</i>	+	+
48	<i>R. styliformis</i>	+	+
49	<i>Skeletonema costatum</i>	+	+
50	<i>Thalassionema nitzschiodes</i>	+	+
51	<i>T. mobiliensis</i>	+	+
52	<i>T. subtilis</i>	+	+
53	<i>Thalassiothrix frauenfeldii</i>	+	+
54	<i>T. longissimi</i>	+	+

## CONCLUSION

Data of the present study show that significant differences occurred in over all percentage composition of phytoplankton species, both in in water and substratum. In water samples, recorded data showed that 85% were of *Bacillariophyceae*, 8 % of *Chlorophyceae* and, 7% of *Dinophyceae*. On the other hand, 75 % *Bacillariophyceae*, 7 % *Chlorophyceae*, 16% *Dinophyceae* and 2% *Cyanophyceae* were recorded in the substratum. The marine biofouling phytoplankton density, diversity, richness, evenness and chlorophyll 'a' were recorded in their maximum levels during summer season, and minimum were obtained in the monsoon. The diatoms were the predominant groups among the phytoplankton. Among the diatoms, species like *Nitzschia longissima*, *Coscinodiscus centralis* and *Fragilaria* sp. were record almost in all the seasons in the present study.

## ACKNOWLEDGEMENTS

Authors are thankful to the Head, Department of Zoology, Nirmala College for Women (Autonomous), Coimbatore-18 and authorities of Bharathidasan University, Tiruchirappalli-24 for the facilities provided.

## REFERENCES

- Coast of India. In Basic and Applied Zooplankton Biology (pp. 45-86). Springer, Singapore.
- Geider, R.J., Moore, C.M. and Suggett, D.J., 2014. Ecology of Marine Phytoplankton 17. *Ecology and the Environment*, p.483.
- Godhantaraman, N., 2002. Seasonal variations in species composition, abundance, biomass and estimated production rates of tintinnids at tropical estuarine and mangrove waters, Parangipettai, southeast coast of India. *Journal of Marine Systems*, 36(3-4), pp.161-171.
- Kawabata, Z., Magendran, A., Palanichamy, S., Venugopalan, V.K. and Tatsukawa, R., 1993. Phytoplankton biomass and productivity of different size fractions in the Vellar estuarine system, southeast coast of India.
- Mani, P., 1992. Natural phytoplankton communities in Pichavaram mangroves.
- Mitra, A., Banerjee, K. and Gangopadhyay, A., 2004. *Introduction to marine plankton*. Daya Publishing House.
- Paffenhofer, G.A., 1998. On the relation of structure, perception and activity in marine planktonic copepods. *Journal of Marine Systems*, 15(1-4), pp.457-473.
- Rajasekar, K.T., Peramal, P. and Santhanam, P., 2005. Phytoplankton diversity in the Coleroon estuary, Southeast coast of India. *Journal of Marine biological association of India*, 47, pp.127-132.
- Reynolds, C.S., 1993. Scales of disturbance and their role in plankton ecology. In *Intermediate Disturbance Hypothesis in Phytoplankton Ecology* (pp. 157-171). Springer, Dordrecht.

55	<i>Triceratium fавus</i>	+	+
56	<i>Triceratium robertsonianum</i>	+	-
57	<i>T. reticulatum</i>	+	-
DINOPHYCEAE (Dinoflagellates)			
58	<i>Ceratium extensum</i>	+	+
59	<i>C. declinatum</i>	+	+
60	<i>C. furca</i>	+	+
61	<i>C. harridum</i>	+	+
62	<i>C. macroceros</i>	+	+
63	<i>C. minutam</i>	+	+
64	<i>C. trichoceros</i>	+	+
65	<i>Dinophysis caudate</i>	+	+
66	<i>Dinophysis</i> sp.	+	-
CYANOPHYCEAE(Blue-Greens)			
67	<i>Trichodesmium erythraeum</i>	-	+
68	<i>Spirulina meneghiniana</i>	-	+
CHOLOPHYCEAE(Greens)			
69	<i>Chlorella vulgaris</i>	+	-
70	<i>C. marina</i>	+	+
71	<i>Clostriopsis longissimi</i>	+	+
72	<i>Tetraselmis gracilis</i>	+	+
73	<i>Oedogoniumde sikaeharyii</i>	+	+
74	<i>Pediastrum simplex</i>	+	-
75	<i>Spirogyra indica</i>	+	-
76	<i>Spirogyra</i> sp	+	-
+ Presence; - Absence			

**Figure 8: Dominant phytoplankton species recorded at Chinnamuttom harbour during May -June months**



- Saravanakumar, A., Serebiah, J.S., Thivakaran, G.A. and Rajkumar, M., 2007. Benthic macrofaunal assemblage in the arid zone mangroves of gulf of Kachchh-Gujarat. *Journal of Ocean University of China*, 6(3), pp.303-309.
- Senthilkumar, S., Santhanam, P. and Perumal, P., 2002, January. Diversity of phytoplankton in Vellar estuary, southeast coast of India. In *Proc. 5th Indian Fisheries Forum (Eds.: S. Ayyappan, JK Jena and M. Mohan Joseph). Published by AFSIB, Mangalore and AeA, Bhubanewar, India* (pp. 245-248).
- Subrahmanyam, R., 1946, October. A systematic account of the marine plankton diatoms of the Madras coast. In *Proceedings of the Indian Academy of Sciences-Section B* (Vol. 24, No. 4, pp. 85-197). Springer India.
- Subrahmanyam, R., 1968. The dinophyceae of the Indian Sea. *Marine Biological Association of India. Mar Biol Assoc India*, 3, pp.118-133.
- Sun, J., Liu, D.Y. and Qian, S.B., 2000. Study on phytoplankton biomass, III. Estimated bulk measurement biomass of phytoplankton in the Jiaozhou Bay. *Acta Oceanol. Sin*, 22, pp.293-299.
- Tomas, C.R. ed., 1997. *Identifying marine phytoplankton*. Elsevier.
- Venkataraman, G., 1939, December. A systematic account of some south Indian diatoms. In *Proceedings of the Indian Academy of Sciences-Section B* (Vol. 10, No. 6, pp. 293-368). Springer India.

# Knowledge and Management Assessment of Temporomandibular Joint (TMJ) Disorders Among Dental interns at King Saud University

Hamad Albagieh<sup>1</sup>, Abdulrahman Alghamdi<sup>2</sup>, Musaad Almefyaz<sup>2</sup>,  
Abdulrahman Alshehri<sup>2</sup> and Osama Alsaiady<sup>2</sup>

<sup>1</sup>Oral Medicine and Diagnostic Sciences Department, College of Dentistry,  
King Saud University, Riyadh, Saudi Arabia.

<sup>2</sup>College of Dentistry, King Saud University, Riyadh, Saudi Arabia

## ABSTRACT

The temporomandibular joint (TMJ) complex is mainly composed of bone, cartilage, muscles, ligaments, and neurovascular channels that supply the tissues. The temporomandibular disorders (TMD) are collective term embracing a number of clinical problems that involve the masticatory muscles, the temporomandibular joints (TMJs) and associated structures, or both. The most common symptoms of temporomandibular joint disorders (TMDs) are pain, noises in the joint, limitation of mouth opening, or a combination of these. The objective of this study was to assess the knowledge and management of temporomandibular joint (TMJ) disorders among dental interns at King Saud University, Saudi Arabia. The sample size of this study consisted of 100 dental interns at King Saud University. Informed consent from the participants was obtained. The participants were informed that their responses and personal information would be kept confidential. A cross-sectional survey was prepared in English language using Google Forms. The survey contains 12 questions was distributed among the selected sample of dental interns at King Saud University. The average knowledge score of the dental interns on the knowledge regarding management of (TMJ) disorders was 4.04 (SD = 1.42). The maximum score obtained was 8 out of 8, the minimum was zero. that the majority of the dental interns (67%) had fair knowledge. The finding of the present study showed, there was a fair level of knowledge about (TMJ) disorders among dental interns. Many interns are unable to diagnose and treat patients suffering from temporomandibular joint disorders.

**KEY WORDS:** TEMPOROMANDIBULAR JOINT, KNOWLEDGE AND MANAGEMENT, DISORDERS AMONG.

## INTRODUCTION

The temporomandibular joint (TMJ) complex is mainly composed of bone, cartilage, muscles, ligaments, and neurovascular channels that supply the tissues. The temporomandibular disorders (TMD) are collective term embracing a number of clinical problems that involve the masticatory muscles, the temporomandibular joints (TMJs) and associated structures, or both. The most common symptoms of temporomandibular joint disorders (TMDs) are pain, noises in the joint, limitation of mouth opening, or a combination of these (Glick et al., 2015). The most common musculoskeletal diseases affecting the temporomandibular joint and associated structures in the orofacial region

are temporomandibular joint disorders. Many dentists are unable to diagnose and treat patients suffering from temporomandibular joint disorders. the misdiagnosis of the orofacial pain of the temporomandibular joint associated with the dentoalveolar region, which leads to incorrect management (Al-Huraishi et al., 2020).

TMDs are diagnosed using a variety of methods including clinical examination, radiographic image such as: panoramic X-rays, conventional tomography (CT), Digital Volume Tomography, Arthrography, Magnetic Resonance Imaging (MRI), and Cone Beam-Computed Tomography (CBCT) (Pettersson 2010). Pain relievers, nonsteroidal anti-inflammatory medications (NSAIDs), muscle relaxants, anticonvulsants (gabapentin), and tricyclic antidepressants (TCAs) are among the medications used to treat TMDs (López-Frías 2019).

**Article Information:**\*Corresponding Author: [Halbageah@ksu.edu.sa](mailto:Halbageah@ksu.edu.sa)

Received 15/04/2023 Accepted after revision 25/06/2023

Published: June 2023 Pp- 126-131

This is an open access article under Creative Commons License,

<https://creativecommons.org/licenses/by/4.0/>.


Available at: <https://bbrc.in/> DOI: <http://dx.doi.org/10.21786/bbrc/16.2.10>

In another study, Rabaa Aboubakr and Elkwatehy (2021) found that the knowledge level about TMDs was insufficient among undergraduates and dental intern students. The aim of this study was to assess the knowledge and management of temporomandibular joint (TMJ) disorders among dental interns at King Saud University, Saudi Arabia.

### MATERIAL AND METHODS

The sample size of this study consisted of 100 dental interns at King Saud University. Informed consent from the participants was obtained. The participants were informed that their responses and personal information would be kept confidential. A cross-sectional survey was prepared in English language using Google Forms. The survey contains 12 questions was distributed among the selected sample of dental interns at King Saud University. The first three questions were designed based on an established Likert scale, the questions were related to the ability of undertake a clinical examination regarding TMJ, assess the treatment needs and management of TMJ and the third question was related to satisfaction of lectures and clinical training giving at undergraduate curriculum. The followed eight questions were to measure and assess the knowledge and management regarding temporomandibular joint (TMJ) disorders.

**Data Analysis:** All questionnaire data were included for analysis purposes in the SPSS 22 software (SPSS, Chicago, USA). Each element has been assigned to the value 1 in case of a correct answer and 0 if it is incorrect. Descriptive statistics were used to show the results in terms of frequencies, percentages, means and the standard deviation.



**Table 1. The total number of participants in this study (N=100):**

		Frequency	Percent
GENDER	Male	60	60.0%
	Female	40	40.0%
	Total	100	100.0%

### RESULTS AND DISCUSSION


The total number of participants in this study was 100. Participants were recruited in this survey, 60% males and 40% females.

Competencies to undertake a clinical examination, assess the treatment needs and management regarding temporomandibular joint (TMJ) disorders: The participants were asked question about their competencies to undertake a clinical examination, assess the treatment needs and management regarding temporomandibular joint (TMJ) disorders. The results were presented in the following tables:

**Table 2. Competency to undertake a clinical examination (N=100):**

Responses	No	Percent	Mean	SD
Strongly agree	5	5.0	3.03	0.96
Agree	28	28.0		
Neutral	36	36.0		
Disagree	27	27.0		
Strongly disagree	4	4.0		
Total	100	100.0		

**Figure 2**



**Table 3. Competency to assess the treatment needs and management (N=100):**


Responses	No	Percent	Mean	SD
Strongly agree	2	2.0	2.84	0.84
Agree	19	19.0		
Neutral	43	43.0		
Disagree	33	33.0		
Strongly disagree	3	3.0		
Total	100	100.0		

As shown in Table 2, that only 33% of dental interns were able to undertake a clinical examination regarding (TMJ) disorders and reach to the correct diagnosis, while 31% of




them reported they do not have the ability to undertake a clinical examination regarding (TMJ) disorders and reach to the correct diagnosis. However, the high percentage of participants were not sure about that i.e. (did not give an exact response). Moreover, the average response toward this question was (3.03± 0.96) which tend to be neutral.

As shown in Table 3, the high percentage of participants (43%) (mean ±SD 2.84, 0.84) were not sure about their ability to assess the treatment needs and management regarding (TMJ) disorders, while 36% were not able to do that. However, only 21% of dental interns were able to assess the treatment needs and management regarding (TMJ).



**Table 4. Satisfaction of lectures and clinical training (N=100):**

Responses	No	Percent	Mean	SD
Strongly agree	3	3.0	2.40	1.07
Agree	12	12.0		
Neutral	31	31.0		
Disagree	30	30.0		
Strongly disagree	24	24.0		
Total	100	100.0		



**Satisfaction of lectures and clinical training during undergraduate curriculum to diagnosis and manage (TMJ) disorders:** The participants were asked question

about if they are satisfied about the coverage of lectures and clinical training during undergraduate curriculum to diagnose and manage (TMJ) disorders. The result was presented in the following table:

As shown in Table 4, more than half of dental interns (54%) reported that lectures and clinical training covered during undergraduate curriculum were not enough to diagnose and manage (TMJ) disorders. However, 31% were neutral about that i.e. (did not give an exact responses). On the other hand, only 15% of participants reported that lectures and clinical training covered during undergraduate curriculum were enough to diagnose and manage (TMJ) disorders.

Knowledge and management of (TMJ) disorders among dental interns at King Saud University: In this section of the study the knowledge and management of (TMJ) disorders were assessed among dental interns at King Saud University in Saudi Arabia. As shown in Tables 5, regarding dental interns' knowledge about (TMJ) disorders at King Saud University, that the overall percentage of correct answers to this subscale was 46.5%.

However, the results revealed that 68% of dental interns were knowledgeable that Ankylosis of the temporomandibular joint (TMJ) most often results from Trauma or infection. Moreover, 56% of dental interns had the knowledge that the most probable diagnosis for the case (A well-nourished 41-year-old female patient present with a chief complaint of painful limited mouth opening, her mouth deflects to the right side while opening, she reported receiving a trauma from her teenage son one week ago) was acute disc displacement without reduction in the right TMJ.

Moreover, the majority of dental interns involved in this study failed to answer the item (A 23-year-old former ballet dancer came to your clinic complaining of a loud clicking sound in both sides of her TMJs without any pain, her mouth opening is 55mm, with slight deviation to the right then to the left, started suddenly 16 months ago, during examination; clicking sound existed even after mandibular advancement, the most probably diagnosis for this case is) correctly by 76%. Also, 62% of dental interns failed to answer item (Which of the following statements is true regarding the temporomandibular joint?) correctly. As shown in Tables 6, regarding dental interns' knowledge about (TMJ) disorders at King Saud University, that the overall percentage of correct answers to this subscale was 36%.

However, the results revealed that the most item pertained to management of (TMJ) disorders that dental interns aware about the treatment of dislocation, that 70% of dental interns had the knowledge that dislocation is treated by forcing the mandible downward and backward. As well, 51% of dental interns had the knowledge that best way to control painful anterior disc displacement cases was by administering non-steroidal anti-inflammatory drugs.

Moreover, that the most item that dental interns failed to answer correctly was "A 35-year-old female teacher came to your clinic complaining of severe pain in the left side

of her jaw started three months ago, her medical history was significant for systemic osteoarthritis, that she's been diagnosed last year. Her pain is severe with dull aching

quality. She is using paracetamol, but it doesn't help anymore, her mouth opening is limited because of the pain. The best way to control the pain in this case is achieved by:", 90% of them failed to answer this item correctly.

Items	Answers	
	Correct	Incorrect
Which of the following statements is true regarding the temporomandibular joint?	38%	62%
Ankylosis of the temporomandibular joint (TMJ) is defined as immobility or fusion of the joint. This condition most often results from which of the following?	68%	32%
A 23-year-old former ballet dancer came to your clinic complaining of a loud clicking sound in both sides of her TMJs without any pain, her mouth opening is 55mm, with slight deviation to the right then to the left, started suddenly 16 months ago, during examination; clicking sound existed even after mandibular advancement, the most probably diagnosis for this case is	24%	76%
A well-nourished 41-year-old female patient present with a chief complaint of painful limited mouth opening, her mouth deflects to the right side while opening, she reported receiving a trauma from her teenage son one week ago. What is the most probable diagnosis for this case?	56%	44%

Items	Answers	
	Correct	Incorrect
A 35-year-old female teacher came to your clinic complaining of severe pain in the left side of her jaw started three months ago, her medical history was significant for systemic osteoarthritis, that she's been diagnosed last year. Her pain is severe with dull aching quality. She is using paracetamol but it doesn't help anymore, her mouth opening is limited because of the pain.	10%	90%
The best way to control the pain in this case is achieved by: In painful anterior disc displacement cases, which one of the following modalities is the best way to control pain?	51%	49%
A 23-year-old former ballet dancer came to your clinic complaining of a loud clicking sound in both sides of her TMJs without any pain, her mouth opening is 55mm, with slight deviation to the right then to the left, started suddenly 16 months ago, during examination; clicking sound existed even after mandibular advancement, the most probably diagnosis for this case is:	13%	87%
Dislocation is treated by forcing the mandible	70%	30%

Also, 87% of dental interns involved in this study failed to answer the item "A 23-year-old former ballet dancer came to your clinic complaining of a loud clicking sound in both sides of her TMJs without any pain, her mouth opening is 55mm, with slight deviation to the right then to the left, started suddenly 16 months ago, during examination; clicking sound existed even after mandibular advancement, the most probably diagnosis for this case is".

**Knowledge scores regarding Knowledge and management of (TMJ) disorders among dental interns at King Saud University:** The knowledge score was classified as follow: Poor knowledge (<4) , Fair knowledge (4-6) and good knowledge (7-8) (Table 7).

Knowledge Level	No.	Percentage			
Poor (<4)	31	31%			
Fair (4-6)	67	67%			
Good (7-8)	2	2%			
Total	100	100.00%			
Correct answers		Min.	Max.	Mean	SD
		0	8	4.04	1.42

As shown in the above Table, that the average knowledge score of the dental interns on the knowledge regarding management of (TMJ) disorders was 4.04 (SD = 1.42). The maximum score obtained was 8 out of 8, the minimum was zero. As presented in Table 6, that the majority of the dental interns (67%) had fair knowledge. Many studies have been done to assess the knowledge and management regarding TMDs among general practitioners and dental students but none of them has been done on dental interns. This the study has been conducted on dental Interns only.

According to the current survey, the average score of the dental interns on the knowledge and management regarding (TMJ) disorders was fair. The participants were asked question about if they are satisfied about the coverage of lectures and clinical training during undergraduate curriculum to diagnose and manage (TMJ) disorders, the result was more than half of dental interns (54%) reported that lectures and clinical training covered during undergraduate curriculum were not enough to diagnose and manage (TMJ) disorders.

Many studies have been conducted to assess the level of knowledge and management regarding TMDs among general dental practitioners (GDPs) and dental students. Aboubakr and Elkwehly (2021) study which was conducted among undergraduate dental students and interns revealed that the knowledge level among the participants regarding TMDs was not sufficient as the percentage of participants who answered correctly to the questions was less than 50% for all questions except in two questions they were more than 50% which were asking about the common medications prescribed for TMDs, another survey by Elyas

et al. (2021) revealed that general dental practitioners had fair level of knowledge regarding TMDs.

This finding is also in accordance with a survey by Patil and Iyengar (2016) showed evident that GDPs had fair level of knowledge regarding TMDs, whereas the majority of the experts had a good level of knowledge. Thus, a significant difference in the level of knowledge was observed between the two groups. Also, Rahmeier et al. (2021) revealed that the students had little knowledge regarding TMD and bruxism due to the insufficient credit hours offered by the curriculum. Le Resche et al. (1993) reported that general dental practitioners (GDPs) had low knowledge level regarding pathophysiology, diagnosis and treatment in comparison to TMDs specialists. However, Ashwin and Siri (2018) revealed good knowledge level among their postgraduate participants. In addition, López Frías et al. (2019) showed sufficient knowledge among postgraduate students regarding etiology and diagnosis of TMDs in comparison to lower knowledge in management of the same condition among their GDPs participants.

In dentistry and educational programs regarding TMDs, importance is given to applying and converting learned knowledge to diagnose and manage TMDs in clinical practices. Thus, the current study emphasizes the importance of developing and improving undergraduate curricula to provide dental practitioners with both theoretical and practical knowledge regarding the diagnosis and management of TMDs. Continuing dental education programs, fellowships, and TMJ workshops may help dental practitioners to improve their skills and knowledge.

## CONCLUSION

The anatomy and pathophysiology of the TMJ are important topics in the curriculum of dental students that help with the proper diagnosis and management of TMJ disorders. The finding of the present study showed, there was a fair level of knowledge about (TMJ) disorders among dental interns. Many interns are unable to diagnose and treat patients suffering from temporomandibular joint disorders. Further studies on a larger scale with a larger sample size are required to get a proper overview regarding this topic among dental interns.

**Ethical approval:** This article does not include any studies involving human participants or animals performed by the author. An exemption letter was provided by the Institutional Review Board (IRB) (No. E-22-7415).

**Funding:** This research did not receive any specific grant from funding agencies in the public commercial or not for profit sectors.

## REFERENCES

Aboubakr, R., & Elkwehly, W. (2021). Knowledge about temporomandibular joint disorders among dental students in Saudi Arabia. *Egyptian Dental Journal*, 67(4), 2885-2892.

Al-Huraisi, H.A., Meisha, D.E., Algheriri, W.A., Alasmari,

- W.F., Alsuhaimeh, A.S. and Al-Khotani, A.A., 2020. Newly graduated dentists' knowledge of temporomandibular disorders compared to specialists in Saudi Arabia. *BMC Oral Health*, 20, pp.1-9.
- Ashwin, P.S. and Siri, P.B., 2018. Assessment of knowledge, attitude and practice on temporomandibular joint disorders among post-graduate dental students. *Int J Dent Res*, 3(4), pp.59-62.
- Elyas, H., Alomran, S., Almosharraf, E., Alomar, G. and Alkharaan, M., 2021. Assessment of knowledge, attitude, and practices of dental practitioners regarding temporomandibular joint disorders in Saudi Arabia. *International Journal of Medicine in Developing Countries*, 6(1), pp.6-13.
- Glick, M., 2015. *William M. Burket's Oral Medicine*. 12th ed. People's Medical Publishing House: USA, pp.104-8.
- Le Resche, L., Truelove, E.L. and Dworkin, S.F., 1993. Temporomandibular disorders: a survey of dentists' knowledge and beliefs. *Journal of the American Dental Association* (1939), 124(5), pp.90-4.
- López-Frías, F.J., Gil-Flores, J., Bonilla-Represa, V., Ábalos-Labruzzi, C. and Herrera-Martinez, M., 2019. Knowledge and management of temporomandibular joint disorders by general dentists in Spain. *Journal of clinical and experimental dentistry*, 11(8), p.e680.
- Patil, S. and Iyengar, A.R., 2016. Assessment of knowledge, attitude and practices of dental practitioners regarding temporomandibular joint disorders in India. *Journal of Advanced Clinical and Research Insights*, 3(2), pp.64-71.
- Petersson, A., 2010. What you can and cannot see in TMJ imaging—an overview related to the RDC/TMD diagnostic system. *Journal of oral rehabilitation*, 37(10), pp.771-778.
- Rahmeier, G., Irineu, G.L., Maracci, L.M., Salbego, R.S., Nascimento, L.P., Marquezan, B., Silva, B. 2021. Dental students' knowledge assessment of TMD and Bruxism at UFMS. *Revista da ABENO*, 21(1), pp.1-10. DOI: 10.30979/revabeno.v21i1.1253.

1. Life Members and Fellows of Society of Science & Nature (MSSN/FSSN), Bhopal, India will be entitled to receive free early on line issues of Biosc.Biotech.Res.Comm for life. They will get substantial waivers for publication of their research papers.
2. Selected life members on the basis of their academic and research contributions will be conferred with Honorary Fellowship of SSN (FSSN), who will be instrumental in scientific awareness programs, particularly encouragement and popularization of science. These members will be appointed reviewers / editors of the Journal in different subject areas. Life Fellow members of SSN will be invited to attend society sponsored conferences and seminars in India.

**Form**

For Member, Society for Science & Nature and  
Bioscience Biotechnology Research Communications  
(MSSN & BBRC)

AND

Fellow, Society For Science & Nature (FSSN) &  
Member, Bioscience Biotechnology Research Communications  
(BBRC)

Website: Society: [www.ssnb.org.in](http://www.ssnb.org.in) Publisher Email-  
[Publisher@ssnb.org.in](mailto:Publisher@ssnb.org.in) Website Journal: [www.bbrc.in](http://www.bbrc.in)  
E-mail: [bbrc.in.info@gmail.com](mailto:bbrc.in.info@gmail.com)

Kindly download the form from the Societys website [www.ssnb.org.in](http://www.ssnb.org.in) (Photocopies will be accepted). Forms can also be downloaded from our journals, Bioscience Biotechnology Research Communications website [www.bbrc.in](http://www.bbrc.in)  
Send completed forms by email to [editor@bbrc.in](mailto:editor@bbrc.in) or [Publisher@ssnb.org.in](mailto:Publisher@ssnb.org.in)

Life Membership Fellow SSN (One Time Subscription (FSSN) including Life Member-ship of BBRC is Rs. 6000/- Direct NEFT to be made on contacting the Managing Editorat [bbrc.in.info@gmail.com](mailto:bbrc.in.info@gmail.com) Foreign Members will have to obtain separate invoices from [editor@bbrc.in](mailto:editor@bbrc.in) for making payment to the Society/ BBRC Fellowship.

Name: Dr./Prof. (IN CAPITAL LETTERS): \_\_\_\_\_

Designation & Organization: \_\_\_\_\_

Qualification & Specialization of Research: \_\_\_\_\_

Present and Past Academic Positions: \_\_\_\_\_

Research Publications Experience Enclose Biodata\* with full publications list: \_\_\_\_\_

Academic Achievements Memberships of Societies etc.: \_\_\_\_\_

Mailing Address (With tel./Mob./Email id): \_\_\_\_\_

I wish to become life Member / Fellow of Society for Science And Nature Bhopal, India. I have read the details and agree to abide by them.

Signature

Name and Address / Email

Details of Accompanying Payment NEFT No..... DD No.....Date.....

**JOIN AS LIFE MEMBER BIOSCIENCE BIOTECHNOLOGY  
RESEARCH COMMUNICATIONS  
SUBSCRIPTION FORM FOR BBRC (ONLY JOURNAL)**

Kindly complete this form if you want to become life member/ of BBRC only (Individual Life Member BBRC Rs. 5000/-) I wish to become Life Member of Bioscience Biotechnology Research Communications.

Name: \_\_\_\_\_

Address: \_\_\_\_\_

E-mail: \_\_\_\_\_ Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Type of Membership /Life Member, ( MSSN with BBRC) / Fellow, (FSSN with BBRC) Amount in Rs-----/(In words) Paid By NEFT No.....

NEFT favoring Treasurer, Bioscience Biotechnology Research Communications, For NEFT contact the Managing Editor.

Website Society: [www.ssnb.org.in](http://www.ssnb.org.in) Website Journal: [www.bbrc.in](http://www.bbrc.in)  
E-mail: [bbrc.in.info@gmail.com](mailto:bbrc.in.info@gmail.com) Publisher Email:- [publisher@ssnb.org.in](mailto:publisher@ssnb.org.in)

**BBRC SUBSCRIPTION RATES**

- |   |              |
|---|--------------|
| 1. Life Member (MSSN & BBRC) Only   | INR 5000/-   |
| 2. Institutional Annual Member India Libraries for (Hard Copies of 4 Issues per year Postage Extra) | INR 16,000/- |
| 3. Single Hard Copy of Journal per Issue (to be ordered in advance, Postage Extra)                  | INR 4000/-   |



**Declaration about the ownership of  
Bioscience Biotechnology Research Communications  
Form (IV) [See Rule 3]**

1. Place of Publication : Bhopal, India
2. Periodicity of its Publication : Six Monthly
3. Printer's Name : Ayesha S. Ali On behalf of Society  
For Science & Nature
- (Whether Citizen of India) : Yes
- Address : H. No. C-52, H.B. Colony, Kohefiza  
Bhopal-462001, India
4. Publisher's Name : Ayesha S. Ali on Behalf of Society  
For Science & Nature
- (Whether Citizen of India) : Yes
- Address : H. No. C-52, H.B. Colony, Kohefiza  
Bhopal-462001, India
5. Editor's Name : Dr. Sharique Ali
- (Whether Citizen of India) : Yes
- Address : H. No. C-52, H.B. Colony, Kohefiza  
Bhopal-462001, India
6. Name & Address of the individual/ : Ayesha S. Ali  
who own the newspaper & : H. No. C-52, H.B.Colony, Kohefiza,  
partners or share holders Bhopal-462001, India  
holding more than one percent  
of the total capital
- (Whether Citizen of India) : Yes

I, Ayesha S. Ali hereby declare that the particulars given above are true to the best of my knowledge and belief.

Date : 31<sup>st</sup> June 2023  
Place : Bhopal

# Bioscience Biotechnology Research Communications

Open Access International Journal Indexed by Clarivate Analytics USA, Web of Science ISI, ESCI

Bioscience Biotechnology Research Communications

P-ISSN: 0974-6455 E-ISSN: 2321-4007

CODEN (USA): BBRCBA

Indexed in Thomson Reuters ISI Now Clarivate Analytics Web of Science (ESCI)

Publishers: Society for Science and Nature, Bhopal India

Journal Unique Identifier: Cross Ref DOI: <http://dx.doi.org/10.21786>

Periodicity: Jan-Feb-March, April-May-June, July-Aug-Sep and Oct-Nov-Dec

Journal Website: <https://bbrc.in/>

## (Important Links of Journal)

1. **Manuscript Processing Flow Chart:** <https://bbrc.in/bbrc/wp-content/uploads/2019/05/Flowchart1.pdf>
2. **Manuscript Template:** <https://bbrc.in/manuscript-template/>
3. **Author Ethical Statement & Copyright form / Plagiarism Check Report:** <https://bbrc.in/plagiarism-and-ethical-statement/>
4. **Cover letter with Reviewers and their addresses (see template):** <https://bbrc.in/wp-content/uploads/2021/10/Cover-letter-Bioscience-Biotechnology-Research-Communications.pdf>
5. **Manuscript On Line Submission:** <https://bbrc.in/homepage/submit-article-2/>

## Instructions for Authors / Detailed MS Submission Guidelines For Bioscience Biotechnology Research Communications

All manuscripts must be submitted to Bioscience Biotechnology Research Communications

Only through the journals online submission system at <https://www.bbrc.in> (<https://bbrc.in/homepage/submit-article-2/>)

Author submitting the manuscript for the first time is required to register online and create a profile as an author. This enables the authors to receive login credentials for manuscript submission. Manuscripts must consist of duly completed Author Ethical Statement / Copyright Form along with plagiarism / similarity level Certificate of the submitted MS, (which should be less than 20%. Attach Certificate checked by Ithenticate / Turnitin Software). **This is a mandatory part of manuscript submission.**

Before final submission, please make sure that the manuscript conforms to the journal guidelines and instructions to authors for the preparation of the manuscript.

**MS not prepared as per instructions to authors will not be entertained and will be returned as incomplete submission.**

Please note that the journal does not charge any fees for submission of articles, and we do not give any fixed frame of time to publish an article, since the review of articles depends upon the reviewers processing time, the editorial assessment, and production. Roughly a MS takes about 60 to 90 days from the date of submission to publication, depending upon the review process and number of revisions envisaged.

### **1. Ethical & Plagiarism Policies of Bioscience Biotechnology Research Communications:**

([Author Ethical Statement / Copyright form / Plagiarism Check Report](#))

Plagiarism is the unauthorized use or close imitation of the language and thoughts of another author and representing them as one's own original work and Biosc. Biotech. Res. Comm. strictly condemns all forms of plagiarism, following a very vigilant policy of removing this malady. Within the academia, it is considered dishonesty or fraud and offenders are subject to academic censure. Plagiarism can be unintentional or intentional, reproducing academic material without appropriate credit to the original authors (Citations).

Similarly self-plagiarism is the re-use of significant, identical or near identical portions of one's own work without citing the original work. This is also known as recycling fraud. Worst form of plagiarism is to steal the whole article or in parts from some source and publish it under one's own name in another journal. Plagiarism, fabrication, unethical or redundant publication grossly violates the editorial policies of Biosc Biotech Res Comm. which follows best practice guidelines given by the International Committee of Medical Journal Editors (ICMJE) and Committee on Publication Ethics (COPE), as mentioned in the Journals Instructions for Authors. Biosc. Biotech. Res. Comm. strongly condemns any form of plagiarism and unethical practices.

**All authors submitting their MS to Biosc Biotech Res Comm must complete and sign the ethical statement form**

**(downloaded from above link) and append the Plagiarism Check Certificate of their MS along with ethical statement form, failing which their MS will be not processed further.**

Authors submitting their work to Biosc.Biotech.Res.Com must also mention the names, addresses and email ids of three subject experts to serve as independent reviewers for their submitted MS, in their cover letter. The reviewers must not be of their Institution, it is not necessary the same reviewers will be appointed for their submitted manuscript, selection of independent unbiased reviewers is under the purview of editorial board / editors.

**The following files need to be submitted with every article:**

- 1. Cover Letter** stating the originality of research and why you think it should be published in Biosc Biotech Res Comm. along with names / addresses and emails of 3 external reviewers must be attached, (See [Cover Letter template](#)).
- 2. Manuscript Text:** For preparation and style of MS (See [Manuscript Template](#)):

The full manuscript should contain first page with full author names, affiliation, ORCID No and the corresponding author email / ORCID details, followed by full text of the MS file in word format, not exceeding 4000 words or 20 pages. All data/tables/figures/Images (images must be submitted with the MS in high print-reproducible resolution).

**2. Article Types:** Submission of the following article types is considered for publication in Biosc. Biotech.Res. Comm.

- 1. Original Research Articles**
- 2. Critical Meta Reviews**
- 3. Case Reports with Discussion**
- 4. Short Communications**
- 5. Letters to the Editor / Editorials / Perspectives / Correspondence**

#### **(I) Original Research Articles**

Manuscript must be written in good English, typewritten using Times New Roman font size 12 only, double-spaced with one inch margin on all sides. All manuscripts must be accompanied by author declaration with ethical certificate signed by the corresponding author and all co-authors that they have seen and approved the final version of the manuscript and that the article has NOT been published or submitted to any other journal for publication. The corresponding author is responsible for obtaining permission from the copyright owner for the use of any copyrighted material in the submitted article.

Each original article must contain the following in the order as:

#### **Title page: Title page should contain the following information:**

Main Title of the article followed by short running title, Name (s) of author(s), Department (s)/Institution(s) City / Code & Country, where the work was performed, with all author ORCID links, (<https://orcid.org/login>). E-mail address of the corresponding author marked with an asterisk \* is necessary.

#### **2. Abstract:**

Abstract should be factual summarization of the entire work and should NOT TO EXCEED 250 words, with 5 keywords written below it. Abstract must have following subheadings:

#### **Introduction (Objectives / Rationale), Brief Methods, Results and Conclusion**

#### **3. Main Text of the Manuscript: Text must be arranged under the following headings:**

- 1. Introduction**
- 2. Material and Methods**
- 3. Results (Including Tables/Fig/Images)**
- 4. Discussion**
- 5. Conclusion followed by Funding Statements /Acknowledgements (if any).**
- 6. References (Strictly in Harvard Style)**

**Introduction:** This section must provide a brief review of literature, purpose of the study, objectives and the rationale of the research undertaken should be given with proper clarity.

**Material and Methods:** This section of material and methods /procedures should be concise but detailed enough to enable the reader to reproduce the experiments / methodology. Commonly used procedures and methods in detail need not be described, but require a reference to the original source.

**Results (Including Tables/Fig/Images):** Give only brief findings, presented in the form of tables or figures, should be included without duplication of presentation and no discussion of the significance of the data, either tables or figures be given, avoid duplication of data.

**Discussion** should present the significance of the present data under the prevalent understanding and interpretation of the phenomenon. Speculative discussion is allowed but it should be concise and corroborated by the presented data.

**Conclusion** summarizes the study and is drawn from the results and discussion, should not be more than 100 words.

**Acknowledgements/ Financial Acknowledgements** if any, should be placed at the end of Conclusion before References.

## 6. References: (Strictly as per Harvard Style)

References in text of the manuscript should be written using last author name (s) without their initials with year in PARENTHESES ( ).

The final bibliography in the **References Section** should be **arranged alphabetically using last name of the author** and written in **Harvard Style** as shown below in examples of references: **All references must be written in 11 point font Roman letters.**

*Use Italic styles only for scientific names of organisms, genera, species in the entire MS as well as in the Reference section.*

*In this section et al should be used only after three names of authors.*

In reference section, DOIs / Links of the references from PubMed, WoS–Clarivate Analytics, Scopus, Google Scholar and others must also be provided.

All references should be checked minutely, for their appearance in text as well as in References, incomplete or missing references in the text or in Reference List & Vice versa will not be accepted, and the MS will be returned as **Incomplete Submission**.

### a. Example of Reference from a Standard Journal Article:

Ali Sharique A, S Salim, Sahani T, Peter J and Ali AS (2012c) Serotonergic receptors as novel target for optimizing skin pigmentary responses in Indian bull frog, *Hoplobatrachus tigerinus* British Journal of Pharmacology Vol 165 No 5 Pages 1515-1525.

### b. Example of Reference from a book:

Falconer DC (1960) Introduction to Quantitative Genetics. Oliver & Boyd Edinburgh 165-185.

### c. Reference from article in a book:

Ali, Sharique A, N Parveen and Ayesha S Ali (2021) In Herbal Medicine: Back to The Future, Promoting Melanocyte Regeneration Using Different Plants and Their Constituents – Vol 3 (Ed. Ferid Murad, Nobel Laureate) Bentham Science, USA Pages 247-276.

**Tables and Figures (or Images):** Short, Precise Tables and sharp image figures must be included, complete with legends /footnotes / explanation / units should be right below them. The tables and figures pages should be consecutively numbered, and arranged between results and discussion. Position of the tables or figures in the text of the MS must be indicated using same numbers.

**Instructions for Preparation of Images:** An image refers to the following: Graphs, photographs, maps, charts, paintings, drawings, diagrams, etc. Images must be embedded within the manuscript text between Results and Discussion of the article, not separately or at the end of the article. Once the article is accepted for publication, the author may be asked for submission of image in high resolution file formats. It is strongly recommended before embedding images in the manuscript, images must be prepared as mentioned below in the image specifications section.

**Image specifications:** Images must be prepared in accordance with the instructions mentioned on the PubMed Central website: <https://www.ncbi.nlm.nih.gov/pmc/pub/filespec-images/> The key factor for preparation of MS images for sufficient quality is images must have a minimum resolution of 300 dots per inch (dpi) for the grayscale (or black and white) and at least 600 dpi for color scale. The acceptable image formats are tiff, jpeg, gif, psd or png.

**Image Copyright:** For any image that the authors have not made themselves, the authors will need to have written permission to reproduce that image, even if the image is posted on the internet. It is the author's responsibility to obtain permission to use the images, not the publishers. Permission must be obtained in writing before the article can be submitted. For complete information, please visit the Copyright Agency Limited website: <http://www.copyright.com.au/get-information/about-copyright>.

## (II) Critical Review Articles / Systematic Reviews / Meta-Analysis

**(Simple Reviews Are not considered for publication in Biosc.Biotech.Res.Comm.)**

Systematic Reviews or Meta-Analysis should be systematic, critical assessments of most recently updated literature and data sources pertaining to basic biological or bio-medical science topics that include a statistical technique for quantitatively combining the results of multiple studies that measure the same outcome into a single pooled investigation. Data must be searched for and selected systematically for inclusion and critically evaluated, and the search and selection process for compiling the review must be mentioned. The text should NOT exceed 5000 words excluding abstract, references, tables and figures.

Each of the sections of the **Systematic Review or Meta Analysis** articles should include specific sub-sections as follows:

### 1. Structured Abstract: (Not exceed 250 words):

Objectives, Methodology, Results and Conclusion

2. **Introduction:** Rationale, Objectives, Research questions

3. **Methodology:** Study design, Participants, interventions, comparators

4. **Systematic Review Protocol:** Search strategy, Data sources, Studies Sections and Data Extraction, Data analysis/ Statistical tools used

5. **Results and Discussion:** In results provide flow diagrams / attractive tables / figures of the studies retrieved for the review, study selection characteristics synthesized findings, risk of bias etc.

**6. Summary:** Abstract of main findings, Limitations, Conclusions etc.

**For all other information including title page, typing and reference style etc, please follow the instructions to authors for Research Articles.**

### **(III) Case Reports with Discussion**

The case reports, of two or more patients must contain genuinely new interpretational information, discussed with up to date literature. The reports should have clinical significance, new adverse effect(s) of a drug or other unique first time observations, etc. Patient consent for publication must be obtained from the patient in written or, if this is not possible, the next of kin before submission. The author(s) must have been involved in the care of the patient.

Case Report /case description should start with a single paragraph abstract followed by text, which should not exceed 2000 words (excluding references, tables and figures) with maximum 10 bibliographic references and either three figures or three tables. Case report / case presentation must contain:

- 1. Brief Abstract (should not exceed 150 words)**
- 2. Introduction**
- 3. Case Presentation**
- 4. Reviews & Discussion**
- 5. Conclusion**
- 6. References**

Patient Consent, Competing interests, Funding Statement, Acknowledgements (if any). For all other information including title page, typing and reference style, please follow the instructions for original articles.

### **(IV) Short Communications**

Short communication should be original work, such as complete results of a short pilot study, not merely a preliminary report and should not exceed 2000 words with one or two figures and/or one table. An editorial decision will be provided rapidly without reviews. For writing and references style, follow the same instructions listed above.

### **(V) Letters to the Editor/Editorials / Perspectives / Correspondence**

Opinions on topics and articles recently published in the journal will be considered for publication if they are objective and constructive in nature and provide academic interest to the readers. These letters may also be forwarded to the author of the cited article for possible response. The editor reserves the right to shorten these letters, delete objectionable comments, make other changes, or take any other suitable decision to comply with the style and policies of the journal. For writing and references style, follow the same instructions listed above.

### **(VI) Editorials**

Editorial will be written by one member of the editorial board as solicited by the Editor-in-Chief. The editorial is generally a scientific review on one or two of the current topics pertaining to biomedical sciences.

## **4. Article Processing Charges (APC) and Waivers**

Bioscience Biotechnology Research Communications does not have any article submission charges, however authors will be required to pay only article processing charges (APC) that too after acceptance of their peer reviewed manuscripts.

**We do not have any other charges for publication of MS in Biosc. Biotech. Res. Comm. like color print charges or reprint charges, author subscription charges or any other fees.**

The moderate APC taken from authors contributes to the handling/ editorial/ production / open access/ HTML/ DOI / costs and hence is non-refundable. APC is to be deposited via Net Banking/ Electronic Transfer after acceptance of the manuscript only.

Article Processing Charges (APC) for Authors from India- Rs.7000/\*

Article Processing Charges (APC) for SAARC Countries – US Dollars 175

Article Processing Charges (APC) for Low Income Countries- US Dollars 250

For All other Countries the APC is US dollars 425

\*Waivers available for Non Funded Research.

**Publication Fee Waivers:** In order to meet the rigorous academic standards on a fast track, the open access journal has some expenses as stated above, and for these reasons we charge a very modest article processing fee. **Nevertheless, as we believe that lack of funds should not be a barrier to quality open access publication, Biosc Biotech Res Comm has a policy to provide significant waivers to deserving authors from middle and low income countries without any financial support. Authors can request for a waiver in such cases.**

## **5. Conditions of Acceptance of Manuscripts**

Acceptance of Manuscript: On acceptance, the editors retain the right to make stylistic changes in the MS, shorten the material as necessary and decide on the date and periodicity of publication.

**6. Galley Proofs:** Authors will be sent an online copy of the galley proofs to the email id of only the corresponding author. Corrections should be confined to typographical errors or matters of accuracy. Authors should return their galley proofs within two days of receipt. If there is a delay in



returning the proofs beyond the given deadlines the MS will be published in next issue, no changes in the MS will be possible once the author sends the corrected galley.

## 7. Early On Line Ahead of Print Publication / Final Publication

Early on Line E- Prints, ahead of final publication, is provided by Bios Biotech Res Comm to enable authors and readers to have early and free access to their published work.

## 8. Checklist for Authors While Submitting Their Manuscripts

- **As part of the on-line submission process, authors should carefully check their submission, using the below Check List for careful compliance with the following items as the manuscript will be returned to the authors as Incomplete Submission if any of the following points is missing.**
- The main manuscript has been prepared by all the concerned authors, after carefully reading all the Instructions to Authors.
- All authors through the corresponding or principal author have filled and enclosed the Author Ethical Statement, Copy Right and Plagiarism-Check Certificate along with their manuscript.
- **The above forms of Author Ethical Statement, Copy Right and Plagiarism-Check Certificate should be downloaded from journals website [www.bbrc.in](http://www.bbrc.in) and must be filled, signed by all authors and attached with the MS.**
- The submission file format is in ~Microsoft Word document file and not a PDF.
- The text is double-spaced and should be within the word limit of 4000 words or 20 pages with a 12-point Roman font
- Italics must be used only for all scientific / Latin / Greek names.
- **A single manuscript word file has been submitted that contains title page, short running title, author details, abstract followed by main manuscript.**
- **Check all correct authors names, their addresses, email ID of corresponding author and ORCID link of all authors.**
- **A brief cover letter stating why the submission is suitable for Bios Biotech Res Comm must be attached mandatorily giving names, Addresses and e-mail ids of 3 subject experts to serve as unbiased reviewers, who should be from different universities and institutions.**
- Main MS file must be in word format, single and must contain all text matter headings such as Title, Short Running Title, Abstract, MS Main Text Matter Tables / Figures / and References in it.
- Abstract of 250 words must be written under headings: **Background / Introduction, Objectives and Rationale, Brief Methods, Results and Conclusion.**
- Mention 5 key words below the abstract in alphabetical order
- Only 5 subheadings are required in the main MS: **Introduction, Material and Methods, Results & Discussion, Conclusion and References.**
- Ethical approvals / consent to participate must come in Methodology. Acknowledgements / funding details (if any) must come after Conclusion before References.
- All illustrations, figures, and tables are properly numbered and should be arranged between Results & Discussion.
- Size of tables / figures must not be more than half a page. All legends of tables / figures must be written right below them.
- **References should be written in text with AUTHOR LAST NAME WITH YEAR IN PARENTHESES ( )**
- **Strictly as per Harvard Style of References. Do not use any italics for names of Journals or their Volumes Numbers or years**
- **All references in the References Section must be alphabetically arranged using only the first author's last name as per Harvard style.**
- **Use of et al in Reference Section must only be used after writing three author names.**
- Name, designation, institution and email address of three independent reviewers related to the **Subject area of research must be provided in the cover letter along with the manuscript.**

### **Journal's Address:**

*Head Office: Editor in Chief*

*Bioscience Biotechnology*

*Research Communications,*

Post Box No 01 GPO Bhopal 462001 India

### ***Delhi Office: Bioscience Biotechnology Research Communications***

Care of AIHMS 31, Gautam Nagar, Behind

AIIMS New Delhi -110049, India

[editor@bbrc.in](mailto:editor@bbrc.in) website: [www.bbrc.in](http://www.bbrc.in)

### **Publisher's Name & Address:**

Society For Science & Nature,

C-52 HB Colony, Kohe-Fiza, Bhopal 462001, India

**Country: India Website: [ssnb.org](http://ssnb.org)**

Name of the Scholar	Fellow Society for Science & Nature (FSSN) and Member BBRC	Designation and Address of the Scholar
Dr. Sharique A. Ali, FLS FRSB (UK)	FSSN/BBRC	Professor and Head, Department of Biotechnology Saifia Science College, 462001 Bhopal, India
Dr. Ayesha S. Ali	FSSN/BBRC	Professor Department of Biotechnology Saifia Science College, Bhopal 462001, India
Dr. J. Peter	FSSN/BBRC	Associate Professor, RKDF University Gandhi Nagar, Bhopal 462023 India
Dr. M. Miraj	FSSN/BBRC	Director, Institute of Health & Management Studies, Gautam Nagar New Delhi India
Prof. D K Belsare PhD DSc FNASc	FSSN/BBRC	Biosciences Baylor College of Medicine Houston USA. & Barkatullah University Bhopal, India
Dr. GN Wankhede	FSSN/BBRC	Professor & Head, Department of Zoology, SGB University Amravati (MS) India
Dr. Sarita Shrivastava	FSSN/BBRC	Professor of Chemistry, Govt Motilal Vigyan Mahavidyalya, Bhopal India
Dr. P.M. Makode	MSSN/BBRC	Associate Professor of Zoology, Venue, Park, Shegaon Naka VMV Road, Amravati (MS) India
Dr. Sunita Yadav	MSSN/BBRC	Assistant Professor Department of Zoology, Satya Sai College for Women BHEL, Bhopal India Bhopal
Dr. Romsha Singh	MSSN/BBRC	Associate Professor, Department of Zoology, MLB Girls College, Bhopal India
Dr Shaima Miraj PhD	FSSN/BBRC	College of Health Sciences, Saudi Electronic University Riyadh Saudi Arabia
Dr Sushma Prasad PhD	MSSN/BBRC	Zarifa Farm, Kachhwa Road, Karnal, Haryana 132001, India
Dr Kamal Zaidi PhD	MSSN/BBRC	Department of Microbiology Peoples University Peoples Campus, Bhanpur, Bhopal, 462037 India
Dr. A. D. Lakha	FSSN/BBRC	Associate Professor of Zoology, Nagazari Area, MIT Road, Ambajogai, Beed (MS) 431517 India
Dr. R. S. Virani	MSSN/BBRC	Associate Professor Karimabad Society, Pandhar Kawada, District, Yeobtmal (MS) India
Dr. Madhulika. Pal	MSSN/BBRC	Assistant Professor, Department of Biotechnology Sadhu Vaswani College Bhopal, India
Dr. V.R. Wankhede	MSSN/BBRC	Assistant Professor, Department of Zoology, Deccan College, Pune, India

Dr. Mrs. V. Ingole	MSSN/BBRC	Department of Zoology Vidya Bharti Mahavidyalaya, Amravati, India
Dr. U.N. Bhale	MSSN/BBRC	Associate Professor RTM University, Nagpur, India
Dr. A.P. Sawane	MSSN/BBRC	Associate Professor RTM University, Nagpur, India
Dr. A.D. Bobde	MSSN/BBRC	Associate Professor RTM University, Nagpur, India
Dr. R.G. Jadhaw	MSSN/BBRC	SGB University, Amravati, Amravati 444604 India
Dr SK Pal PhD	FSSN/BBRC	Professor of Genetics Skyline University, Kano, Nigeria
Dr. O.N. Tiwari	FSSN/BBRC	Senior Scientist, Department of Biotechnology NE Region Imphal, Manipur, India
Dr. R. K. Singh	MSSN/BBRC	Research Scientist US Department of Agriculture Washington DC , USA
Dr Anjali Choudhary PhD	MSSN/BBRC	Department of Biochemistry Opposite to Dussehra Maidan, BHEL Square, Sector A, Govindpura, Bhopal, 462023 India
Dr. V. Meitei	FSSN/BBRC	Department of Biotechnology NE Region Imphal, Manipur, India
Dr. N. Qayyumi	MSSN/BBRC	Assistant Professor of Zoology, Mittal College Bhopal, India
Dr LK Jakkala PhD.	FSSN/BBRC	Director Macrocare, Macrocare Towers, Hyderabad Telangana 500081, India
Dr. M. Sajid	MSSN/BBRC	Head Department of Biotechnology Bonnifie College, Bhopal, India
Dr. V. Jaiswal	MSSN/BBRC	Research Scholar, SGB University, Amravati, India
Dr. A. Kumar	MSSN/BBRC	Associate Professor, Department of Biotechnology, SMD Teerth University, Haryana, India
Dr. A.S. Dighde	MSSN/BBRC	Professor of Pomology, Horticulture University of Horticulture Science, Bagalkot
Dr. P Babu	MSSN/BBRC	Plot-18, Street-1, Ashish Nagar (East) Risalt, Bhilai, Durg, CG
Dr. R. Khaliq	MSSN/BBRC	Department of Zoology Holkar Science Colege, Indore India
Dr. A. Siddiqui	FSSN/BBRC	Department of Biotechnology Holkar Science College, Indore India
Dr Raj Sharma PhD	MSSN/BBRC	Pharmaceutical Sciences Chhattisgarh Institute of Medical Sciences (CIMS), Bilaspur, CG, India
Dr. A. Eberle	FSSN/BBRC	Professor Emeritus Biology, University of Basel, Switzerland
Dr. S. Newton	FSSN/BBRC	Professor of Aquaculture University of Virginia, Virginia VA USA

Dr. J Galgut	FSSN/BBRC	Biochemist, Allied Healthcare Systems, Department of Biomedicine, Qatar
Dr. S. Salim	FSSN/BBRC	Research Analyst NIH Bethesda Maryland, USA
Dr. Ruchi Shivle	MSSN/BBRC	Department of Zoology Devi Ahilya Vishvidyalaya, Indore India
Dr. Kirti Dubey	MSSN/BBRC	Department of Fisheries Holkar Science College Indore MP India
Dr. AT Kalse	FSSN/BBRC	Department of Life Sciences North Eastern Maharashtra University, Jalgaon (MS)
Dr. F Kabinwar	FSSN/BBRC	Professor of Oncology, University of California, Los Angeles, USA
Dr Neelu Qayyumi PhD	MSSN/BBRC	Professor and Head Life Sciences Mittal College Opposite Bhopal Memorial Hospital Bhopal India Hospital Research Centre (BMHRC), Navi Bagh, Karond, Bhopal, 462008 India
Dr. Arjun Deb	FSSN/BBRC	Professor of Zoology Lumding College Lumding Assam, India
Dr. Z.Pir	FSSN/BBRC	Department of Biosciences University of Kashmir, Srinagar India
Dr. Razia Sultan	FSSN/BBRC	Department of Zoology Devi Ahilya Vish Vidyalaya, Indore India
Dr. Thingujam I. Devi	FSSN/BBRC	Institute of Bioresources and Sustainable Development, Imphal, Manipur
Dr. I Onyesom	FSSN/BBRC	Professor of Microbiology Abraska, Delta State Nigeria
Dr. K. Sudhakar	FSSN/BBRC	Assistant Professor Energy Centre, Maulana Azad National Institute of Technology Bhopal, India.
Dr. Ravi Jain	FSSN/BBRC	Associate Professor of Physics Samrat Ashok Technical Institute Vidisha India
Dr Shiv Kumar Jayant	MSSN/BBRC	Department of Biochemistry All India Institute of Medical Sciences Bhopal India
Prof. C Rama Mohan	MSSN/BBRC	Narayana Engineering College, Jawaharlal Technological University Anantpur AP India
Dr. Sushil Kumar Upadhyay D. Phil	FSSN/BBRC	Assistant Professor, Department of Biotechnology Maharishi Markandeshwar (Deemed to be University) Mullana Ambala 133207 (Haryana) India



# Bioscience Biotechnology Research Communications

An Open Access International Journal [www.bbrc.in](http://www.bbrc.in)

Post Box 01, GPO, Bhopal 462001 India

P-ISSN: 0974-6455 O-ISSN: 2321-4007 CODEN USA: BBRCBA

---

## (AUTHOR ETHICAL STATEMENT / COPYRIGHT FORMS / PLAGIARISM CHECK REPORT)

Articles must be submitted by only the corresponding author of the manuscript, and should not be submitted by anyone on behalf. The corresponding author may submit this Copyright/ Ethical Statement Form along with the manuscript, on behalf of all the co-authors (if any). The author (s) will confirm that the manuscript (or any part of it) has not been published previously or is not under consideration for publication elsewhere. Furthermore, any illustrations, structures or tables that have been published elsewhere must be properly reported with citations/ and or, copyright permission for reproduction must be obtained.

1. I / (We) confirm that the enclosed article entitled-----Authored by: 1.....2.....3.....  
.....4.....5.....et al has not been published previously in whole or in part, nor is being considered elsewhere for publication, and if accepted for publication in Biosc.Biotech.Res.Comm, it will not be published elsewhere.
2. I / We acknowledge that on the condition of acceptance, Biosc Biotec Res Comm and its authors will have the copyright of the scholarly work which grants usage rights to others using an open license ([Creative Commons](https://creativecommons.org/licenses/by/4.0/)) allowing for immediate free access to the work, provided it is properly cited as per standard guidelines. Financial support / fundings have been duly acknowledged.
3. I / We also confirm that all necessary permissions, ethical considerations for animal and human rights for experimentation to carry out this research have been obtained by the authors from the concerned authorities.
4. **It is also certified that the manuscript has been prepared as per instructions to the authors, complying all the author instructions, policies of plagiarism, its check and ethical statement as required by Biosc Biotec Res Comm. All authors have seen the final manuscript and approve its publication.**
5. **We also certify that the similarity / plagiarism levels of the attached manuscript have been checked using Ithenticate /Turnitin software. It has been found to be less than 20% as per international standards and the certificate of same is duly attached with the manuscript.**

Corresponding Author Name.....Orcid Id.....Signature.....

Date.....

Department..... Institution:.....City:.....Country.....

Email:.....

Author 2 Name.....Orcid Id.....Signature.....

Address.....Email.....

Author 3 Name.....Orcid Id.....Signature:.....

Address.....Email.....

Use Extra Space if required.