

Role of Protochlorophyllide oxidoreductase C in protection of plants from singlet oxygen-induced oxidative stress

Garima Chauhan and Baishnab C Tripathy*

School of Life Sciences, Jawaharlal Nehru University, New Delhi 110067, India

ABSTRACT

Exposure of plants to extreme environmental conditions greatly reduces their growth by increasing the concentration of reactive oxygen species (ROS). The elevation in the ROS accumulation directly impacts the photosynthetic machinery of the plant. Among the ROS, singlet oxygen (1O_2) cannot be detoxified by any antioxidative enzyme-mediated reactions. Therefore, 1O_2 is the major cause of damage to plants during day time. Spraying plants with a chlorophyll precursor 5-aminolevulinic acid (ALA) increases the accumulation of Chl biosynthetic intermediate protochlorophyllide (Pchlde) which acts as a photosensitizer. Upon light exposure of ALA-treated plants, overaccumulated Pchlde in chloroplasts gets excited and transfer their absorbed energy to oxygen to generate 1O_2 via type-II photosensitization reaction. The 1O_2 immediately damages thylakoid membranes and induces wilting in the whole plant. The *porC-2* mutant that lacks the photoenzyme protochlorophyllide oxidoreductase C (PORC) fail to photo-transform overaccumulated Pchlde to Chlide upon light exposure therefore, leading to the excess generation of 1O_2 . This leads to severe damage to photosynthetic apparatus that destroys chlorophylls and the photosynthetic reactions as indicated from Chl a fluorescence imaging, reduced *Fv/Fm* and *Fv/Fo* ratio, the quantum yield of PSII and electron transport rate. Conversely, plants overexpressing protochlorophyllide oxidoreductase C (PORCx) are capable of efficiently photo-converting photodynamic Pchlde to Chlide that reduces the 1O_2 production in ALA-treated and light-exposed plants. Reduced 1O_2 produced in PORCx plants causes less damage to the photosynthetic machinery and plants do not bleach. Results demonstrate the pivotal role of protochlorophyllide oxidoreductase C in the protection of plants from 1O_2 -induced oxidative stress during day time.

KEY WORDS: *ARABIDOPSIS THALIANA* · IMAGING PAM · OXIDATIVE STRESS · PROTOCHLOROPHYLLIDE OXIDOREDUCTASE C · SINGLET OXYGEN

Article Information:*Corresponding Author: baishnabtripathy@yahoo.com

Received 15/03/2019 Accepted after revision 25/06/2019

Published: 30th June 2019 Pp-504-511

This is an open access article under Creative Commons License,

Published by Society for Science & Nature, Bhopal India.

Available at: <https://bbrc.in/>

Article DOI: <http://dx.doi.org/10.21786/bbrc/12.2/38>

INTRODUCTION

In a cell, metabolic reactions are tightly regulated to minimize endogenous ROS production. Several antioxidant enzymes and antioxidant molecules work in coordination to regulate the redox status in plants. However, this balance gets disrupted when plants are exposed to extreme environmental conditions (Dat *et al.*, 2000). Excess light exposure over reduces the photochemical electron transport chain, thereby reducing the overall photosynthetic efficiency of the plant (Nishiyama *et al.*, 2001). Apart from chlorophyll molecules present in the antenna and reaction center complex, the membrane-bound chlorophyll (Chl) biosynthetic intermediates can also absorb excess excitation energy and gets excited. Since they are not directly connected to the reaction centers, the energy absorbed by the intermediates cannot be channeled to the photochemical reaction (Tripathy, Mohapatra and Gupta, 2007). Instead, they return to their ground state by transferring their energy to molecular oxygen to give rise to 1O_2 via type II photosensitization reaction (Foote, 1991; Montoya *et al.*, 2005). Accumulation of 1O_2 results into peroxidation of membrane lipids (Triantaphylides *et al.*, 2008), the irreversible oxidation of D1 proteins (Krieger-Liszkay, Fufezan and Trebst, 2008) and gravely damages PS I and PS II (Tripathy and Pattanayak, 2010). Experiments conducted on *chlorina1* mutant (lack chlorophyll b) shows severe phenotypic damage when exposed to light (Ramel *et al.*, 2013). In the absence of chlorophyll b, the photosystem II chlorophyll protein antenna complex becomes non-functional and produces 1O_2 at the natural site of their production due to naked PS II centers. HPLC analysis of enzymatic and nonenzymatic mediated lipid peroxidation products such as octadecatrienoic hydroxyl acids (HOTEs) increased when exposed to high light for 2 days (Ramel *et al.*, 2013; Yang *et al.*, 2019).

Carotenoids are the natural quencher of singlet oxygen, which are present in close abundance in the antenna molecule to dissipate the excess energy absorbed as heat (Baroli *et al.*, 2000). However, in the reaction center, the distance between the reaction center and beta carotene molecule is too large to allow energy transfer, therefore results in the formation of 1O_2 . (Laloi and Havaux, 2015). The Chl biosynthetic intermediates are bound to the plastidic membrane (Mohapatra and Tripathy, 2007). The carotenoids present in the thylakoid and envelope membranes are not in close proximity of Chl biosynthetic intermediates. Therefore, carotenoids fail to quench the 1O_2 from the excited states of Chl biosynthetic tetrapyrroles and allow the production of intermediate derived 1O_2 in response to high light stress. Instead, 1O_2 oxidized carotenoid metabolites actively participate in downstream regulation of gene expression in

response to high light stress, (D'Alessandro and Havaux 2019).

Application of tetrapyrrole precursor, ALA induce accumulation of chlorophyll biosynthetic intermediate, Pchlide (Tripathy and Chakraborty, 1991; Chakraborty and Tripathy, 1992; Fujii *et al.*, 2017). Its conversion into nonphotodynamic intermediate is catalyzed by a light-dependent enzyme, Protochlorophyllide oxidoreductase (POR) which requires light and NADPH to reduce Pchlide to Chlide at the C17 and C18 on the D ring (Armstrong *et al.*, 1995; Oosawa *et al.*, 2000; Pattanayak and Tripathy, 2002). The evolution of POR enzyme occurred very early among oxygenic photosynthetic organisms (Vedalkar and Tripathy 2019). External application of ALA induces production of Pchlide which cannot be converted into Chlide due to the limited supply of POR in WT plants. In rice plants, two isoforms of POR are present, in which POR A is expressed during early development whereas, PORB is present throughout maturity (Kwon *et al.*, 2017).

In the model plant *Arabidopsis thaliana*, multiple isoforms of POR enzyme exist, namely POR A, POR B, and POR C (Armstrong *et al.*, 1995; Reinbothe *et al.*, 1996; Oosawa *et al.*, 2000; Su, Armstrong and Apel, 2001; Pattanayak and Tripathy, 2002). Among them, POR C is highly expressed in response to light and is chiefly present in the photosynthesizing tissues (Oosawa *et al.*, 2000; Masuda *et al.*, 2003). Therefore, overexpression of POR C rapidly converts the accumulated Pchlide into Chlide and thereby reducing the possibility of Pchlide derived 1O_2 oxygen production (Pattanayak and Tripathy, 2011). In contrast, POR C knock-down mutants (*porC-2*) (Masuda *et al.*, 2003) have a limited supply of PORC. We have taken the mutant to modulate Pchlide derived 1O_2 generation in plants to ascertain the role of PORC during oxidative stress. In the present study, WT, PORC overexpressors (PORCx) (Pattanayak and Tripathy, 2011) and *porC-2* mutants were used to modulate the Pchlide sensitized 1O_2 production to study its impact on the photosynthetic efficiency of plants. We have demonstrated the pivotal role of protochlorophyllide oxidoreductase C in the protection of plants from 1O_2 -induced oxidative stress in light.

MATERIALS AND METHODS

Plant material and growth conditions: *Arabidopsis* seeds were soaked in double distilled water for 48h at 4°C for seed stratification. Seeds were sterilized with 4% sodium hypochlorite solution and washed 5 times with autoclaved double distilled water. The sterilized seeds were plated on half-strength Murashige and Skoog media (Sigma). After 12 days of plating, the seedlings were transferred to autoclaved agropete and vermiculite

potting mixture (6:1) and grown at 100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ at 21°C under 14 h light/ 10 h dark photoperiod.

ALA treatment: 1 mM aqueous ALA solution was used to spray on three-week-old pot grown Arabidopsis plants. Later, plants were covered with aluminum foil and kept in the dark for 12 hours to accumulate Pchl_{id}. Control plants were sprayed with distilled water.

Light treatment: After 12 h of dark incubation, ALA-treated and untreated plants were exposed to low light (75 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) for different time intervals.

Imaging PAM: After completion of light treatment, plants were again incubated in the dark for 20 min to open all the reaction centers. Images were captured from IMAGING PAM MAXI chlorophyll fluorometer (Walz, Germany) and the fluorescence parameters were calculated from ImagingWin software (Walz, Germany). Chlorophyll fluorescence parameters such as the maximum quantum yield of PS II (F_v/F_m) was determined by the equation $F_v/F_m = (F_m - F_o)/F_m$ where F_o is the dark fluorescence yield, F_m is the maximum fluorescence yield and $(F_m - F_o)$ is the variable fluorescence (F_v). Maximum efficiency of the water diffusion reaction is analyzed from the ratio of variable to minimum fluorescence yield (F_v/F_o). Other parameters such as Electron transport rate of photosystem II was calculated from the equation $\text{ETR (II)} = \phi \text{ PS II} \times \text{PAR} \times 0.5 \times 0.84$ where $\phi \text{ PS II}$ is effective PSII quantum yield (calculated by $(F'_m - F_t)/F'_m = \Delta F/F'_m$ where F'_m is referred as the maximum fluorescence yield when the samples are illuminated, and F_t is the fluorescence yield at any given time (t). PAR abbreviates for photosynthetically active radiation, 0.5 is the factor of the ratio of PS II and PS I (1:1), 0.84 is the value that correlates with the percentage of incident photons are absorbed by the leaf to drive photosynthesis. The non-photochemical quenching of the maximum fluorescence

was calculated by $(F_m - F'_m)/F'_m$. Quantum yield of nonregulated energy dissipation $Y(\text{NO})$ is calculated by the equation $1/(NPQ + 1 + qL (F_m/F_o - 1))$, where qL represents the fraction of reaction centers that are open according to the lake model (Baker, 2008).

RESULTS AND DISCUSSION

Overexpression of PORC protect plants from 5-Aminolevulinic acid (ALA) induced oxidative stress: WT, PORCx, and *porC-2* plants were grown under the low light regime (100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$). After three weeks, plants were sprayed with 1 mM aqueous ALA solution and covered with aluminum foil and kept in the dark for 12 h to allow phototransformation and were kept under low light (75 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) for 1 or 2 h. Upon exposure to light, due to the limited presence of POR enzyme, plants failed to photo-transform overaccumulated Pchl_{id} to Chl_{id} which resulted in the build-up of excess Pchl_{id} pool that acted as a photosensitizer. The excess Pchl_{id} absorbed a supra-optimal amount of light that could not be transferred to the photosynthetic reaction centers to synthesize reducing equivalents. Instead, the absorbed light energy was transferred to molecular oxygen to generate $^1\text{O}_2$ via type II photosensitization reaction of tetrapyrroles (Tripathy and Chakraborty, 1991; Chakraborty and Tripathy, 1992).

The severity of $^1\text{O}_2$ -induced damage to plants was monitored by IMAGING PAM that measured Chl fluorescence. Figure 1 shows false color images of chlorophyll fluorescence at a time t (F_t), from WT, PORCx overexpressor, and its knock-down mutant *porC-2* plants treated without and with 1 mM ALA. In ALA-treated *porC-2* mutants, the fluorescence was seen from only a few partially green patches of bleached leaves.




FIGURE 1. Overexpression of PORC protect plants from 5-Aminolevulinic acid (ALA) induced oxidative stress.

The *porC-2* mutant plants due to lack of PORC enzyme were the most affected as they failed to photo-transform overaccumulated Pchl_{ide} to Chl_{ide}. Under identical conditions, the overexpressor plants had a lot of fluorescence emanating from greener leaves whose images were recorded by the fluorometer.

WT, PORC overexpressor and its knock-down mutant (*porC-2*) plants of three week stage was sprayed with 1mM ALA or distilled water and kept in the dark for 12 h to accumulate photosensitizer Pchl_{ide}. Chlorophyll a fluorescence images at time t (F_t) was imaged by using Imaging-PAM (Walz, Germany) after 1 h and 2 h of light exposure. The color code beneath the image depict values from 0 to 1.

Singlet oxygen accumulation decreases chlorophyll fluorescence yield: External application of tetrapyrrole precursor ALA to three-week-old WT, PORC_x and *porC-2* plants induced oxidative stress when ALA-treated dark incubated plants were exposed to low light (75 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$), i.e., lower than the intensity at which they were grown (100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$). To understand if ALA-induced oxidative stress influence the maximum PS II quantum yield in dark incubated plants, the *F_v/F_m* ratio was analyzed in both control and ALA-treated samples after 1 h and 2 h of light exposure (Fig.2a). The *F_v/F_m* ratio of dark incubated ALA-treated WT and *porC-2* plants was reduced by 8% and 14% after 1 h of light treatment. Under identical conditions, the *F_v/F_m* ratio of PORC_x plants was unaffected. After 2 h of light treatment, the *F_v/F_m* ratio declined by 14% in WT, 20% in *porC-2* and to a small extent (5%) in PORC overexpressors.

To ascertain the intensity of damage caused by ¹O₂ to the oxygen evolution complex, the *F_v/F_o* ratio in ALA-treated dark incubated plants was monitored upon light exposure (Fig. 2b). The *F_v/F_o* ratio denotes the activity of water splitting complex of PSII. Due to the limited availability of PORC enzyme in *porC-2* and WT plants, the accumulation of Pchl_{ide} caused a burst in ¹O₂ production that significantly damages the oxygen-evolving complex. The maximum decrease in *F_v/F_o* ratio was observed in *porC-2* plants followed by WT whereas PORC_x plants had smaller impairment of photosynthetic oxygen evolution machinery. The *F_v/F_o* ratio declined by 23% and 18% after 2 h of light exposure in *porC-2* and WT plants respectively. However, under identical conditions, minimal damage was observed with respect to the *F_v/F_o* ratio of PORC_x plants.

To understand the effect of accumulated singlet oxygen due to ALA treatment on the quantum yield of PS II, the chlorophyll a fluorescence parameters were analyzed in ALA or distilled water treated dark incubated WT, PORC_x and *porC-2* mutant plants after light exposure. (a) the ratio of variable to maximum fluorescence




FIGURE 2. Pchl_{ide} derived singlet oxygen decreases chlorophyll fluorescence yield.

(*F_v/F_m*) represents the maximum photochemical efficiency of PS II (b) *F_v/F_o* denotes the relative activity of water splitting complex on the donor side of PS II. Each data point is the average of 5 replicates and error bars represent \pm SE. Asterisks indicate significant differences determined by t test (**P*<0.05).

ALA-induced oxidative stress decreased the quantum yield of PS II photochemistry and electron transport rates: The impact of ¹O₂-induced oxidative stress on the overall photochemical quantum yield of PS II (ϕ PSII) was determined at 530 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ light intensity using the equation described in materials and methods. In ALA-treated dark incubated *porC-2* plants, the effective quantum yield of PS II decreased by 50.3% within an hour of light exposure that later declines to 72.6 % after 2 h of light treatment (Fig.3a). In the WT plants, the effective quantum yield decreases from 35% and 63% due to 1h and 2h of light treatment respectively. However, in the PORC overexpressors, the decline in ϕ PSII was 17 % after 2 h of light exposure. The decrease in the electron transport rate (ETR) of PS II measured at 530 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ light intensity was determined in control, and ALA-treated plants. ALA sprayed *porC-2*, and WT plants, exposed to light 75 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ for 2 h had 50% and 35 % reduction in ETR respectively (Fig. 3b). Under identical conditions, the ETR of PORC_x plants was declined by 17.41% after 2 h of light exposure.




FIGURE 3. Exogenous application of ALA decreases quantum yield and electron transport rates of *porC-2* mutant.

Pulse amplitude modulated chlorophyll a fluorescence parameters were analyzed from ALA or distilled water treated samples. (a) Quantum yield of photosystem II (b) Electron transport rate of photosystem II was calculated using Imaging Win Software. Each data point is the average of 5 replicates and error bars represent \pm SE. Asterisks indicate significant differences determined by t test (* $P < 0.05$, ** $P < 0.01$). Activation of heat dissipation machinery in PORCx in response to 102-induced oxidative stress: NPQ is an indicator of activation of heat dissipation machinery in response to stress. Increased NPQ in PORC overexpressor indicates the activation of gradient-dependent heat dissipation machinery (Fig. 4a). Whereas, in *porC-2* and WT, the decrease in NPQ indicate the absence of activation of the protective mechanism. In the control conditions, the Y(NO), that denotes quantum yield of energy dissipation in PS II, were similar in all the three types of plants (Fig. 4b). After ALA treatment, the increase in Y(NO) in *porC-2* and WT indicates that both the processes of photochemical energy conversion and protective regulatory mechanisms are unable to cope up even with the low light ($75 \mu\text{mol photons m}^{-2} \text{s}^{-1}$). Under identical conditions in PORCx plants, the decrease in the Y(NO) indicates that due to excess of PORC enzyme, the Y(NO) remained unaffected after 2 h of light exposure.




FIGURE 4. Activation of heat dissipation machinery in PORCx in response to ALA-induced oxidative stress.

Pulse amplitude modulated Chlorophyll a fluorescence parameters in ALA and distilled water treated samples were determined by Imaging PAM. (a) Non photochemical quenching (NPQ) (b) Quantum yield of non regulated energy dissipation in PS II. Each data point is the average of 5 replicates and error bars represent \pm SE. Asterisks indicate significant differences determined by t test (* $P < 0.05$). In a plant cell, $^1\text{O}_2$ is produced as a byproduct of cellular metabolic function. The limited amount of $^1\text{O}_2$ can be easily quenched by carotenoids (Ramel *et al.*, 2012). However, when plants are exposed to photooxidative conditions, the balance between $^1\text{O}_2$ productions and quenching gets perturbed, resulting in an elevation in $^1\text{O}_2$ level. The accumulation of $^1\text{O}_2$ not only oxidizes several biomolecules but severely impact the metabolic processes (Tripathy and Pattanayak, 2010). In plants, chloroplasts are the major source of $^1\text{O}_2$ generation as the tetrapyrroles that act as photosensitizers are exclusively located in the organelle (Chakraborty and Tripathy, 1992; Triantaphylides *et al.*, 2008; Ambastha *et al.*, 2017).

ALA is the precursor of tetrapyrroles such as chlorophyll, haem, and phycobilins. In the dark, the accumulation of Pchlide modulates ALA synthesis by a feedback mechanism. Regulation of ALA synthesis is essential to prevent the accumulation of photodynamic chlorophyll metabolic intermediates. This enables plants to evade photooxidative stress (Rebeiz *et al.*, 1988; Tripathy and Chakraborty, 1991; Chakraborty and Tripathy, 1992). Unlike other isoforms of POR, which are prominently present in the etiolated seedlings, PORC expres-

sion is present throughout the leaf development and it increases in response to high light. In PORCx plants, the rapid conversion of Pchl_{ide} to Chl_{ide} takes place due to maximum availability of PORC enzyme (Pattanayak and Tripathy, 2002, Pattanayak and Tripathy, 2011). Chl_{ide} produced as a result of phototransformation is immediately converted into chlorophyll that associates itself to the chlorophyll binding proteins present in the thylakoid membranes and transfers the absorbed energy to the reaction centers to drive photosynthetic reactions.

To understand the effect of Pchl_{ide} derived ¹O₂ on the photosynthetic machinery of plants, their chlorophyll fluorescence parameters were studied by IMAGING-PAM (Kandoi *et al.*, 2016). The energy absorbed by Chl can be utilized through either one of the three processes. Most primarily, the energy absorbed by the Chl molecule is directed to initiate the photochemical reactions. Secondly, the absorbed energy is dissipated in the form of heat, and third, the excited Chl molecules return to the ground state by fluorescence. These are three competing processes. Analysis of modulated chlorophyll fluorescence in response to different light pulses provide valuable information regarding the photosystem II activity in a quick and non-invasive manner. The ratio of variable to maximal fluorescence (*F_v/F_m*) indicates the health status of dark-adapted plants. Under control conditions, a dark-adapted, healthy and non-stressed plant emits maximum fluorescence (*F_m*) after illuminated with a short pulse of saturating light. Therefore the ratio of variable to maximum fluorescence was closer to 0.78 in non-stressed leaves, which is an effective parameter to study the maximal quantum efficiency of PS II (Björkman and Demmig, 1987). Similarly, under stress, a decline in the light-adapted maximal fluorescence (*F'_m*) indicates the release of the absorbed energy as heat due to activation of nonphotochemical quenching (NPQ). Likewise, other parameters of chlorophyll a fluorescence can be interpreted to obtain information regarding the quantum yield and PSII-dependent electron transport rate.

In this study, we have observed that plants over-expressing PORC enzyme were capable of tolerating ALA-induced oxidative stress. The *porC-2* mutant has T-DNA inserted in the 4th exon of PORC gene (Masuda *et al.*, 2003). The unavailability of PORC enzyme reduces efficient phototransformation of Pchl_{ide} to Chl_{ide}. The nontransformed Pchl_{ide} upon receiving light gets excited like a Chl molecule and returns to the ground state by generating singlet oxygen via type-II photosensitization process of Pchl_{ide} (Tripathy and Chakraborty, 1991; Chakraborty and Tripathy, 1992). Further, spraying *porC-2* plants with ALA increases the accumulation of Pchl_{ide} and Pchl_{ide} derived ¹O₂. Therefore, maximum damage to the photosynthetic machinery was observed

in the *porC-2* mutant. Enhanced level of ¹O₂ can reduce photosynthetic efficiency by oxidizing structural and functional proteins associated with the PS II. During electron transport across the PS II, D1/D2 heterodimer constitutes the core of reaction center. D1 protein participates in the charge separation and electron transfer events during the electron transport. Production of ¹O₂ in the plastidic membrane due to the accumulation of membrane-bound Chl biosynthetic intermediates (Pchl_{ide}) can easily target D1 protein. When the oxidative damage caused by ¹O₂ increase beyond the D1 repair machinery, impairment in the reduction of P680 directly influences the electron transport rate and quantum yield of PS II (Edelman and Mattoo, 2008).

The increased production of ¹O₂ in *porC-2* mutant bleached substantial amounts of Chls leaving a few green patches in the leaves. The fluorescence was emitted only from a few green patches that were not severely bleached. The WT plants using its endogenous POR could photo transform some of the Pchl_{ide} to Chl_{ide}, resulting in lower Pchl_{ide} accumulation than *porC-2* mutant. Therefore, WT had lesser bleaching of leaves due to reduced generation of ¹O₂ than *porC-2* mutants, and consequently, the fluorescence was emitted from a higher number of green leaves. The fluorescence imaging clearly reveals that the PORCx plants were protected from photooxidative damage having fluorescence emission from a large number of leaves due to the minimal generation of ¹O₂ in light.

The *F_v/F_m* ratio, a measure of the quantum efficiency of PSII, in dark-adapted leaves, declined in WT and *porC-2* plants. The PORCx plants always had higher *F_v/F_m* ratio than that of the WT and *porC-2* plants. In the same vein, the *F_v/F_o* that measures the oxygen-evolving activity in the oxidizing side of PSII implies the activity of water splitting complex, which is very sensitive to redox changes, was lower in light exposed *porC-2* plants. It was due to ¹O₂-induced impairment of oxygen-evolving complex associated with PSII. Similarly, the quantum yield of PSII and the estimated PSII-dependent ETR were higher in PORCx plants due to reduced oxidative damage to the oxygen-evolving complex. Conversely, the heat dissipation of absorbed photons measured as NPQ and quantum yield of non regulated energy dissipation Y (NO) (Genty, Briantais and Baker, 1989) was higher *porC-2* mutants exposed to light. Present results demonstrate the important role of protochlorophyllide oxidoreductase C in the protection of plants from 102-induced oxidative damage in light.

ACKNOWLEDGEMENT

This work was supported by BSR Faculty Fellowship from the University Grants Commission to BCT.

Conflict of interest :The authors declare that they have no conflict of interest

REFERENCES

- Ambastha V Sopory SK Tiwari BS *et al* (2017) Photo-modulation of programmed cell death in rice leaves triggered by salinity Apoptosis Springer US 22(1) pp 41–56 doi: 10.1007/s10495-016-1305-7
- Armstrong GA Runge S Frick G *et al* (1995) Identification of NADPH:Protochlorophyllide Oxidoreductases A and B: A Branched Pathway for Light-Dependent Chlorophyll Biosynthesis in *Arabidopsis thaliana* Plant Physiology 108(4) pp 1505–1517 doi: 10.1104/pp.108.4.1505
- Baker NR (2008) Chlorophyll Fluorescence: A Probe of Photosynthesis In Vivo Annual Review of Plant Biology 59 (1) pp 89–113 doi: 10.1146/annurev.arplant.59.032607.092759
- Baroli I Niyogi KK (2000) Molecular genetics of xanthophyll-dependent photoprotection in green algae and plants Philosophical Transactions of the Royal Society B: Biological Sciences 355(1402) pp 1385–1394. doi: 10.1098/rstb.2000.0700
- Björkman O and Demmig B (1987) Photon yield of O₂ evolution and chlorophyll fluorescence characteristics at 77 K among vascular plants of diverse origins Planta 170(4) pp. 489–504 doi: 10.1007/BF00402983.
- Chakraborty N and Tripathy B C (1992) Involvement of Singlet Oxygen in 5-Aminolevulinic Acid-Induced Photodynamic Damage of Cucumber (*Cucumis sativus* L.) Chloroplasts Plant Physiology 98(1) pp 7–11 doi: 10.1104/pp.98.1.7
- Dat J Vandenabeele S Vranova' E *et al* (2000) Dual action of the active oxygen species during plant stress responses Cellular and Molecular Life Sciences 57(5) pp 779–795 doi: 10.1176/ajp.88.1.103
- D'Alessandro S and Havaux M (2019) Sensing β -carotene oxidation in photosystem II to master plant stress tolerance New Phytologist doi: 10.1111/nph.15924
- Edelman M and Mattoo AK (2008) D1-protein dynamics in photosystem II: The lingering enigma, Photosynthesis Research 98(1–3) pp 609–620 doi: 10.1007/s11120-008-9342-x
- Foote CS (1991) Definition of Type I and Type I1 Photochemistry and Photobiology 54(5) p 659 doi: 10.1111/j.1751-1097.1991.tb02071.x
- Foyer CH and Shigeoka S (2011) Understanding Oxidative Stress and Antioxidant Functions to Enhance Photosynthesis Plant Physiology 155(1) pp 93–100 doi: 10.1104/pp.110.166181
- Fujii S Kobayashi K Nagata N *et al* (2017) Monogalactosyldiacylglycerol Facilitates Synthesis of Photoactive Protochlorophyllide in Etioplasts Plant Physiology 174(4) pp 2183–2198 doi: 10.1104/pp.17.00304
- Genty B Briantais J and Baker N (1989) The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence, Biochimica et Biophysica Acta 990(1) pp 87–92 doi: 10.1016/S0304-4165(89)80016-9
- Kandoi D Mohanty S Govindjee *et al.* (2016) Towards efficient photosynthesis : overexpression of Zea mays phosphoenolpyruvate carboxylase in *Arabidopsis thaliana* Photosynthesis Research Springer Netherlands doi: 10.1007/s11120-016-0224-3
- Krieger-Liszky A Fufezan C and Trebst A (2008) Singlet oxygen production in photosystem II and related protection mechanism Photosynthesis Research 98(1–3) pp 551–564 doi: 10.1007/s11120-008-9349-3
- Kwon C T Kim S H Song G *et al* (2017) Two NADPH: Protochlorophyllide Oxidoreductase (POR) Isoforms Play Distinct Roles in Environmental Adaptation in Rice Rice 10(1) pp 1–14 doi: 10.1186/s12284-016-0141-2
- Laloi C and Havaux M (2015) Key players of singlet oxygen-induced cell death in plants Frontiers in plant science 6 pp 1–9 doi: 10.3389/fpls.2015.00039
- Masuda T Fusada N Oosawa N *et al* (2003) Functional Analysis of Isoforms of NADPH:Protochlorophyllide Oxidoreductase (POR) PORB and PORC in *Arabidopsis thaliana* Plant and Cell Physiology 44(10) pp 963–974 doi: 10.1093/pcp/pcg128
- Mohapatra A and Tripathy BC (2007) Differential distribution of chlorophyll biosynthetic intermediates in stroma Molecular genetics of xanthophyll-dependent photoprotection in green algae and plants Philosophical Transactions of the Royal Society envelope and thylakoid membranes in Beta vulgaris Photosynthesis Research 94(2–3) pp 401–410 doi: 10.1007/s11120-007-9209-6
- Montoya S C N Comini L R Sarmiento M *et al* (2005) Natural anthraquinones probed as Type I and Type II photosensitizers: singlet oxygen and superoxide anion production Journal of Photochemistry and Photobiology B: Biology 78(1) pp 77–83 doi: https://doi.org/10.1016/j.jphotobiol.2004.09.009
- Nishiyama Y Yamamoto H Allakhverdiev S I *et al* (2001) Oxidative stress inhibits the repair of photodamage to the photosynthetic machinery 20(20) pp 1–8
- Oosawa N Masuda T Awai K. *et al* (2000) Identification and light-induced expression of a novel gene of NADPH-protochlorophyllide oxidoreductase isoform in *Arabidopsis thaliana* FEBS Letters 474(2–3) pp 133–136. doi: 10.1016/S0014-5793(00)01568-4
- Pattanayak GK and Tripathy BC (2002) Catalytic function of a novel protein protochlorophyllide oxidoreductase C of *Arabidopsis thaliana* Biochemical and Biophysical Research Communications 291(4) pp 921–924 doi: 10.1006/bbrc.2002.6543
- Pattanayak GK and Tripathy BC (2011) Overexpression of Protochlorophyllide Oxidoreductase C Regulates Oxidative Stress in *Arabidopsis* PLoS one 6(10) doi: 10.1371/journal.pone.0026532
- Ramel F Birtic S Cuine S *et al* (2012) Chemical quenching of singlet oxygen by carotenoids in plants Plant physiology 158(3) pp 1267–78 doi: 10.1104/pp.111.182394
- Ramel F Ksars B Akkari E *et al* (2013) Light-induced acclimation of the Arabidopsis chlorina1 mutant to singlet oxygen The Plant cell 25(4) pp 1445–1462 doi: 10.1105/tpc.113.109827

- Rebeiz C A Zouhior A M Mayasich J M *et al* (1988) Photodynamic herbicides Recent developments and molecular basis of selectivity *Critical Reviews in Plant Sciences* 6(4) pp 385–436 doi: 10.1080/07352688809382256
- Reinbothe S Reinbothe C Lebedev N *et al* (1996) POR A POR B Two light-dependent Protochlorophyllide-Reducing Enzymes of Angiosperm Chlorophyll Biosynthesis *The Plant Cell* 8 pp 763–769
- Su Q Armstrong G and Apel K (2001) POR C of *Arabidopsis thaliana* : a third light- and NADPH-dependent protochlorophyllide oxidoreductase that is differentially regulated by light *Plant Molecular Biology* 47(2001) pp 805–813
- Triantaphylides C Krischke M Hoerberichts F A *et al* (2008) Singlet Oxygen Is the Major Reactive Oxygen Species Involved in Photooxidative Damage to Plants *Plant Physiology* 148(2) pp 960–968 doi: 10.1104/pp.108.12569
- Tripathy B C and Chakraborty N (1991) 5-Aminolevulinic Acid Induced Photodynamic Damage of the Photosynthetic Electron Transport Chain of Cucumber (*Cucumis sativus* L.) Cotyledons *Plant Physiology* 96(3) pp 761–767 doi: 10.1104/pp.96.3.761
- Tripathy BC Mohapatra A and Gupta I (2007) Impairment of the photosynthetic apparatus by oxidative stress induced by photosensitization reaction of protoporphyrin IX *Biochimica et Biophysica Acta - Bioenergetics* 1767(6) pp 860–868 doi: 10.1016/j.bbabi.2007.03.008
- Tripathy BC and Oelmüller R (2012) Reactive oxygen species generation and signaling in plants *Plant Signaling and Behavior* 7(12) pp 1621–1633 doi: 10.4161/psb.22455
- Tripathy BC and Pattanayak Gk (2010) Singlet Oxygen-Induced Oxidative Stress in Plants In Rebeiz CA *et al* (eds) *The Chloroplast: Basics and Applications* Dordrecht: Springer Netherlands pp 397–412 doi: 10.1007/978-90-481-8531-3_25
- Vedalankar P and Tripathy BC (2019) Evolution of light-independent protochlorophyllide oxidoreductase *Protoplasma* pp 293–312
- Yang B Tang J Yu Z *et al* (2019) Light Stress Responses and Prospects for Engineering Light Stress Tolerance in Crop Plants *Journal of Plant Growth Regulation* Springer US 0(0) p. 0. doi: 10.1007/s00344-019-09951-8