

Antimicrobial and Antioxidant Potential of Endophytic Bacteria Isolated from *Emilia sonchifolia* (Linn.)DC.

Sithara K Urumbil¹, Jesy EJ² and Anilkumar M^{3*}

¹Department of Botany, Little Flower College, Guruvayoor, Kerala, Pin 680523

²Department of Chemistry, Little Flower College, Guruvayoor, Kerala, Pin 680523

³Cell Culture Lab, Department of Botany, Union Christian College, Aluva, Ernakulam, Pin-683 102, Kerala, India.

ABSTRACT

Endophytes are symbiotic bacteria that inhabit plant tissues and they were recognised as beneficial microorganisms which does not cause any disease symptoms or adverse effect on the host plant. They may be associated with the production of metabolites that either directly or indirectly influence the medicinal properties of plants. The antimicrobial and antioxidant property of the endophytic bacterial species associated with *E. sonchifolia* can be beneficial for the identification and isolation of valuable bioactive compounds. Six endophytic bacteria were isolated from *Emilia sonchifolia* (Linn.)DC. and their ethyl acetate extract was prepared. This extract was used for the study of antimicrobial and antioxidant properties. The total phenol and flavanoid contents of the bacterial extracts were estimated and antioxidant activity by DPPH, ferric ion reducing, nitric oxide scavenging and cupric ion reducing assays were done. Antioxidant analysis revealed the potential antioxidant property of endophytic bacterial isolates. The Isolate ES1 indicated the highly efficient antioxidant property. GC-MS and LC-MS analyses were employed for the identification of compounds which imparts antimicrobial and antioxidant property of the endophytic bacterial isolate ES1 this pointed the presence of bioactive compounds like surfactin, fengycin, iturin from the bacterial extract.

KEY WORDS: *EMILIA SONCHIFOLIA*, ANTIMICROBIAL, ANTIOXIDANT, ENDOPHYTE, SURFACTIN, ITURIN.

INTRODUCTION

The rapid emergence of multiple drug resistant strains of pathogens demanded the need for the development of plant based antibiotics. Extensive screening of medicinal plants for unexplored metabolites is fast progressing in this area. Many of them harbour bacterial or fungal endophytes that support the production of metabolites. Endophytes

are mutualistic organisms residing inside the plant and produce compounds of pharmacological importance (Pezzuto, 1996). In the recent years, studies on these endophytic microorganisms as an alternative source of various bioactive metabolites resulted in the development of many probiotic and antibiotic compounds. The interest on endophytes associated with medicinal plants led to the identification of different antimicrobial compounds producing strains of microbes. There were many reports on natural products like alkaloids, flavanoids, phenolic compounds, peptides and steroids obtained from medicinal plants that harboured endophytes. In most of the cases the endophytes may either participate or gain some genetic information that led to the production of such metabolites, (Shukla et al., 2015, Ahemad and Kibret, 2014, Gond et al., 2015, Sulistyani et al., 2016, Mohamad et al., 2018, Hazarika et al., 2019).

Article Information:*Corresponding Author: drmakumar@gmail.com

Received 02/04/2020 Accepted after revision 30/06/2020

Published: 30th June 2020 Pp-620-626

This is an open access article under Creative Commons License,.

Published by Society for Science & Nature, Bhopal India.

Available at: <https://bbrc.in/>

Article DOI: <http://dx.doi.org/10.21786/bbrc/13.2/39>

Along with the production of antimicrobial compounds endophytes are reported to produce potent antioxidants (Zheng et al., 2016, Bintang et al., 2015). Health care and food processing industries demand novel antioxidant compounds that rapidly reduce oxidation and manage reactive oxygen species causing damage to cells. Endophytes play a major role either directly or indirectly by enhancing the growth of the host plant and thus the plant growth promotion. Plants harbouring endophytes shows increased growth rate due to the production of various phytohormones, ameliorating the stress tolerance or by checking pathological conditions (Swarnalatha et al., 2015). Therefore metabolites produced by endophytes gained attention as future prospective antibiotics and antioxidants, (Hazarika et al., 2019, Hanif et al 2019).

The present investigation was carried out in *Emilia sonchifolia* (Linn.)DC., a member of Asteraceae family of Dicots. In Ayurveda the plant is highly reputed for its medicinal uses and is an important member in Dasapushpa (Raj et al., 2013). This medicinal plant was reported to produce a broad spectrum of therapeutically active metabolites possessing antimicrobial (Latha et al., 2009, Thenmozhi et al., 2013), antioxidant (Sophia et al., 2011) and anti-inflammatory (Essien et al., 2009) activities. It has been used in the treatment of asthma, inflammatory disorders, cuts and wounds. In ethnomedicine, the plant reported for sore throats, eye, ear ailments, malaria, and measles (Kumar et al., 2015). In this regard enquiry of bacterial endophytes in *Emilia sonchifolia* (Linn.)DC. has been undertaken along with their antioxidant and antimicrobial studies.

MATERIAL AND METHODS

Endophytic bacteria were isolated and identified by 16S rDNA sequencing from *Emilia sonchifolia* (Linn.) DC. (Urumbil and Anilkumar, 2019). The sequence data were submitted to NCBI Gen Bank. The endophytic bacteria isolated were *Bacillus subtilis* strain UCCBOT-ES1 (MG692780), *Paenibacillus sp* strain UCCBOT-ES2 (MG692781), *Microbacterium sp.* strain UCCBOT-ES3 (MG692782), *Bacillus cereus* strain UCCBOT-ES4 (MG692783), *Bacillus aryabhatai* strain UCCBOT-ES5 (MG692784) and *Micrococcus sp.* strain UCCBOT-ES6 (MH027648).

Preparation of extract: The endophytic bacterial isolates were inoculated into 500ml nutrient broth and incubated at 25±2°C for 5 days. The cultures were centrifuged at 8000 rpm for 10 minutes and the supernatant was extracted with double the volume of ethyl acetate and concentrated to a powder form. It was further dissolved in methanol and used for the antimicrobial and antioxidant analysis.

Antimicrobial analysis: The antimicrobial activity of endophytic bacterial extracts were tested against human pathogenic bacteria such as *Escherichia coli* (MTCC 40), *Salmonella typhi* (MTCC 426), *Klebsiella pneumoniae* (MTCC 109) and *Proteus vulgaris* (MTCC 3220), and fungi

such as *Candida albicans* (NCIM 3102) and *Aspergillus niger* (NCIM619) by co culture and disc diffusion methods. The plates were incubated overnight at 37°C and observed for the growth of pathogenic bacteria on either side of the isolates. Further antimicrobial activity of the extracts was studied by disc diffusion method. Streptomycin (10µg/ml) and itraconazole (50µg/ml) was used as antibacterial and antifungal drugs respectively.

Antioxidant analysis: Estimation of total phenol: The total phenol content of the endophytic bacterial extract was calculated by Folin and Ciocalteu method (Swarnalatha et al., 2015). The experiments were repeated thrice and mean value was calculated. The phenolic content was expressed as Gallic acid equivalents in µg/ml.

Estimation of total flavonoid: The total flavanoid content was measured by aluminium chloride colorimetric assay (Kamtekar et al., 2014). The total flavanoid content was expressed as µg/ml of quercetin equivalent.

DPPH free radical scavenging Assay: 50µl of extract in methanol was added to 100µl of DPPH solution and 850µl methanol, so that the final volume was 1ml. Nutrient broth was used as the control and methanol as blank. Percentage of scavenged DPPH radical was calculated using following formula

$$\% \text{ Scavenging} = \frac{A_c - A_1}{A_c} \times 100$$

Where, A_c is the absorbance of control and A_1 is the absorbance of sample. Ascorbic acid was used as standard, (Sulistiyanani et al., 2016).

Ferric ion reducing assay: Extracts prepared from the endophytic bacterial isolates were mixed with 2.5ml of phosphate buffer (0.2M, pH 6.6) and 2.5ml potassium ferricyanide (1%w/v). Method reported by Jayanthi and Lalitha, 2011 using trichoro acetic acid and ferric chloride solution was employed for the ferric iron reducing assay. Ascorbic acid at various concentrations was used as standard.

Nitric oxide radical scavenging assay: 1ml of different concentrations of the extract was mixed with 0.5ml of 10mM sodium nitropruside in phosphate buffered saline and incubated at 25°C for 180 minutes. After incubation the extract was mixed with an equal volume of freshly prepared Griess reagent. Control samples without the extract but with an equal volume of buffer were prepared similar to test sample (Boora et al., 2014). Ascorbic acid was used as positive control. The percentage of nitric oxide radical scavenging activity of the endophytic bacterial extract was calculated.

Cupric iron reducing antioxidant capacity assay (CUPRAC Assay): Cupric iron reducing capacity was measured in accordance to the method of Apak et al., (2008). 1ml of crude extract in ethanol was added with 1ml of 7.5×10⁻³M Neocuproine (Nc) solution, 1ml of 1×10⁻²M CuCl₂ Solution, 1ml of ammonium acetate buffer (pH=7) and

1ml water. Incubate for 30minute at 25°C and measure the absorbance at 450nm and ascorbic acid was used as standard.

Statistical analysis: All data were expressed as mean ±SD. The mean values were statistically analysed using One-way analysis of variance (ANOVA) using the graph pad instat software package.

GC-MSAnalysis and identification: Ethyl acetate fraction of the endophytic bacterial extract with greater antioxidant property was used for GC-MS analysis (Model Number: QP2010S), Column (Rxi-5Sil MS,30 meter length,0.25 mm ID, 0.25 µm thickness), GCMS Software (GCMS Solutions) and Libraries (NIST 11 & WILEY 8).

LCMS/MS-Q-TOF analysis: The crude ethyl acetate fraction of bacterial extract was subjected to liquid chromatography coupled with mass analyser (Waters Xevo G2 QTOF-MS/MS). The separation was carried out in a Acquity BEH C18 column and the two component solvent system contained 90% water (Acidified with 1% formic acid) and 10% acetonitrile with a flow rate of 0.3 ml/min.

RESULTS AND DISCUSSION

Endophytic bacteria are one of the unexplored, promising and relevant producers of metabolites useful in pharmaceutical, health care and agricultural industries. Endophytic bacteria especially many members from the *Bacillus* species, are reported to possess antimicrobial

activity as an indirect mechanism for plant growth promotion by secretion of compounds to check phytopathological conditions (Ahemad and Kibret, 2014, Gond et al., 2015, Mohamad et al., 2018). In the present study, among the different isolates, ES4, ES5 and ES6 exhibited clear growth inhibition against all tested bacterial strains. At the same time ES1, ES2 and ES3 showed growth inhibition against *P.vulgaris* and *S.typhi* only (Fig.1, 2). When disc diffusion method was performed to assess the antibacterial activity, the strain ES4 (*Bacillus cereus* strain), ES5 (*Bacillus aryabhatai* strain) and ES6 (*Micrococcus sp.*strain) produced zone of inhibition against all bacterial strains and maximum zone of inhibition was reported against *P.vulgaris* and *S.typhi* (Table 1, Fig.3, 4). On the contrary strain ES1, ES2 and ES3 showed negligible zone of inhibition against the tested pathogens. Sunkar and Nachiyar (2012) reported the use of *Bacillus cereus* isolated from *Garcinia xanthochymus* for the synthesis of antibacterial silver nanoparticle indicated its significance as an antibacterial agent.

Of the six isolates tested for antifungal activity, ES2 (*Paenibacillus sp.*) produced a zone of inhibition of 15mm which is greater than the standard drug(itraconazole,11mm) (Table 1, Fig.5) Sulistiyani et al., (2016) reported an endophytic *Paenibacillus sp.* with antifungal property from *Curcuma longa* rhizome. A particular peptide showing antifungal properties was isolated from *Paenibacillus sp.* (Alkotaini et al., 2014). Anandaraj et al., (2009) isolated two antibacterial peptides Paenibacillin P and Paenibacillin N from *Paenibacillus sp.*

Table 1. Antimicrobial activity of the isolates

Pathogenic strains	Inhibitory activity as per co-culture method						Zone of inhibition(in mm) in disc diffusion method							Std.
	ES1	ES2	ES3	ES4	ES5	ES6	ES1	ES2	ES3	ES4	ES5	ES6		
<i>E.coli</i>	-	-	-	-	+	-	0	0	0	0	9	0	25	
<i>S.typhi</i>	+	+	+	+	+	+	0	0	0	24	19	17	21	
<i>K.pneumoniae</i>	-	-	+	+	+	+	9	8	0	9	9	9	22	
<i>P.vulgaris</i>	+	+	+	+	+	+	8	9	8	20	15	12	20	
<i>C.albicans</i>	-	-	-	-	-	-	0	0	0	0	0	0	13	
<i>A.niger</i>	-	-	-	-	-	-	0	15	0	0	0	0	11	

Figure 1: Total phenol content of the isolates at different concentrations of extract




Figure 2: Total flavanoid content of the isolates at different concentrations of extract.





Figure 4: Antifungal activity of isolate ES2 against *A.niger*



Antimicrobial activity of endophytic bacterial extract-Fig.1 & 2-growth inhibition activity of endophytic bacterial isolate in co culture, Fig.3-Antibacterial activity of isolates against *Proteus vulgaris*, Fig.4-Antifungal activity of isolate ES2 against *A.niger*

Sophia et al., (2012) reported the antioxidant property of *E. sonchifolia* the medicinal plant used for the isolation of endophytes in the present study. In the present attempt the antioxidant activity of the endophytic bacterial extracts showed positive results indicating that the isolate as well as host plant synthesized some metabolites with common functions that might be due to the plant microbe interactions. Estimation of phenolic content revealed that ES1 produced maximum amount (156µg/ml), while ES4 produced the lowest (Graph 1). Total flavanoid content of the extracts showed that the isolate ES1 contained 37.3µg/ml quercetin equivalent of flavanoid and ES2 and ES5 have comparable flavanoid contents of 29.6µg/ml, 18.6µg/ml of quercetin equivalent respectively (Graph 2). *Bacillus sp.* have been preferred as a probiotic bacteria in the feed industry because of its antioxidant property (Wang et al., 2017).

The most accurate method used for antioxidant analysis was found to be the DPPH assay. Total phenol and flavanoid estimation showed higher amount of these components in *Bacillus subtilis* strain. Free radical scavenging activity was measured in terms of IC50 value and lowest IC50 indicates a highest antioxidant activity. The highest antioxidant activity by DPPH assay was observed in ES1 (IC50 0.825µg/ml), followed by ES5 (IC50 1.19µg/ml), and ES6 (IC50 1.044µg/ml). Reports by Nongkhilaw and Joshi (2015) specified that L-Asparaginase production was directly linked with antioxidant property and the *Bacillus subtilis* strain cenB associated with *Centella asiatica* showed positive results for L-Asparaginase production and antioxidant property.

The results of each antioxidant assay may be different among the isolates based on the presence of various metabolites produced by the isolates and it cannot be comparable among the assays (Rafat et al., 2012). When ferric ion reducing power analysis was conducted, isolate ES1 (IC₅₀ 0.33µg/ml) showed highest reducing power. The extracts of isolates, ES5 (IC₅₀ 0.43µg/ml) and ES6 (IC₅₀ 0.68µg/ml) also showed higher antioxidant property in accordance with the principles of Ferric reducing power assay. The IC₅₀ value of Nitric oxide radical scavenging assay showed that isolates ES6 (IC₅₀ 2.79µg/ml), ES3 (IC₅₀ 3.14µg/ml), ES4 (IC₅₀ 3.12 µg/ml) had significant antioxidant activity. Among the isolates ES5 showed highest cupric ion reducing property with lowest IC₅₀ value (0.103µg/ml). The statistical analysis was carried out by one way ANOVA followed by Tukey test (Table 2).

Table 2. IC50 values in antioxidant assays

	DPPH Assay	Ferric ion reducing power	CUPRAC Assay	Nitric oxide radical Scavenging assay
Std	1.654	0.84	0.138	3.68
ES1	0.825***	0.33***	0.66ns	3.49ns
ES2	2.23***	1.55*	0.147***	5.6 ns
ES3	2.17***	1.25**	0.266**	3.14 ns
ES4	2.9***	2.21ns	0.254**	3.12 ns
ES5	1.19***	0.43**	0.103***	3.47 ns
ES6	1.044***	0.68**	0.423ns	2.79**











***p < 0.001- highly significant, **p < 0.05-significant, nsp > 0.05-not significant

Antioxidant and antimicrobial studies revealed that the isolate ES1 has significant activity and hence GC-MS Analysis of the ES1 extract was carried out. It revealed nine detectable compounds (Table-3). The major compound detected in the GCMS analysis in accordance with area percentage is Hexadecanoic Acid, 15-Methyl-, Methyl Ester (Common name-Methyl isoheptadecanoate) and it was reported to have antioxidant, nematocidal (Imran et al, 2007, Zayed et al., 2014), antifungal and antibacterial properties (Ali et al., 2017). It was for first time we report the presence of this compound from the extract of endophytic bacteria that may contribute in the bioactivity. We conclude that the endophytic bacteria isolated from *E. Sonchifolia* possess significant antimicrobial and antioxidant properties.

Bioactivity studies revealed that *Bacillus subtilis* strain UCC BOT ES1 was the most promising among all isolates. LCMS/MS analysis was employed to screen the bioactive compounds produced by this isolate. *Bacillus subtilis* was already reported as one of the frequently identified bacterial endophytes known to be efficient in plant growth enhancement (Malfanova et al., 2011) and biocontrol activities (Luo et al., 2015, Kim et al., 2017).

Liquid chromatography with mass spectral analysis of different *Bacillus* members revealed high potential for the production of natural bioactive compounds (Hazarika et al., 2019).

Table 3. GC-MS identification of compounds in extract of isolate ES1

Peak#	R.Time	Area	Area%	Height	Height%	Name	Base m/z	
1	7.618	116466	18.15	33533	17.29	Cyclotrisiloxane, Hexamethyl-	207.00	
2	28.493	162474	25.32	50188	25.87	Hexadecanoic Acid, 15-Methyl-, Methyl Ester	74.10	
3	32.317	26412	4.12	11411	5.88	Hexanoic Acid, 2-Methyl-	74.05	
4	32.403	137413	21.41	45488	23.45	2,2',2''-Nitrilotriethanol, triethyl ether	174.10	
5	44.641	55758	8.69	8570	4.42	[1,1'-Bicyclopropyl]-2-octanoic acid, 2'-hexyl-,methyl ester	149.15	
6	45.232	6727	1.05	8144	4.20	Thiofanox	145.15	
7	46.539	108149	16.85	18678	9.63	1-Heptatriacotanol	161.10	
8	48.131	20271	3.16	9335	4.81	Propenone, 1-(1-hydroxycyclohexyl)-3-(4-hydroxyphenyl)-	81.10	
9	49.166	8002 641672	1.25 100.00	8651 193998	4.46 100.00	15-(4-HYDROXYANILINO) RETINAL #	209.05	

A cyclic lipopeptide (cLP) biosurfactant 'surfactin' was reported from *Bacillus* species (Haddad et al., 2008). Apparent surfactin peaks were noticed in the current study and were comparable with the earlier reports (Sarwar et al., 2018). Peaks corresponding to the molecular mass 992, 1007, 1022 and 1035 therefore indicated the surfactin biosynthetic potential of *Bacillus subtilis* strain UCC BOT ES1. Peaks observed at 1083.58 [M+H⁺] was noted to be that of another non-ribosomal peptide iturin as per the previous reports. Iturin category of compounds was generally considered as antifungal compounds (Arrebola et al., 2010). Fengycin is another category of lipopeptide with wide range of

applications as antifungal compounds and they were efficient in preventing the growth of filamentous fungi (Malfamova et al., 2011). Presence of fengycin was reported from different members of endophytic *Bacillus* (Hanif et al., 2019). The molecular weight observed in the present study [M+H⁺] 1417.78 and 1415.82 are in accordance with fengycin A (Table 4). Non-ribosomally synthesised lipopeptides showed greater degrees of structural similarity and the difference among them was confined to the fatty acid chain (Roongsawang et al., 2010). However efficient purification and screening methods were required to separate and analyse the different class of lipopeptide bioactive compounds from the bacterial extract.

Table 4. LC-MS/MS Analysis – Compounds identified in extract of isolate ES1

Lipopeptide	Molecular Mass	[M+H ⁺]	[M+NH ₄]	[M+2K+H]	[M+Cl]	[M+CH ₃ O H+H+]	[M+Na-H ⁺]	[M-H ⁺]
Surfactin	992.51	993.51	1011.53	1069.56	-	-	-	-
	1007.52		1025.53	1083.58	1041.5469	-	-	-
	1022.52	1023.53	1041.54	-	-	-	-	-
	1035.56	1036.56	1053.56	-	-	1069.56	-	-
Iturin	1082.57	1083.58	1101.57	-	-	-	1129.56	-
Fengycin	1414.41	1415.82	1432.03	-	-	-	1058.59	-
	1416.77	1417.78					1461.77	1415.76

ACKNOWLEDGEMENTS

The first author thanks University Grants Commission, New Delhi, for providing financial assistance by Faculty Development Programme.

Conflict of Interest: The authors has no conflict of interest

REFERENCES

- Ahemad M Kibret M (2014) Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective Journal of King Saud University Science Vol 26 Pages1-20.
- Ali A Javaid A Shoaib A (2017) GCMS analysis and antifungal activity of methanolic root extract of *Chenopodium album* against *Sclerotium rolfsii* Planta Daninha Vol35 No e017164713.
- Alkotaini B Anuar N Kadhum AAH Sani AAA (2014) Isolation and identification of a new intracellular antimicrobial peptide produced by *Paenibacillus alvei* AN5. World Journal of Microbiology and Biotechnology Vol 30 Pages 1377-1385.
- Anandaraj B Vellaichamy A Kachman M Selvamanikandan A Pegu S Murugan V (2009) Co-production of two new peptide antibiotics by a bacterial isolate *Paenibacillus alvei* NP75 Biochemical and Biophysical Research Communication Vol 379 No2 Pages 179-85.
- Apak R Güçlü K Ozyürek M Celik S E (2008) Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay Microchimica Acta Vol160 Pages 413-419.
- Arrebola E Jacobs R Korsten L (2010) Iturin A is the principal inhibitor in the biocontrol activity of *Bacillus amyloliquefaciens* PPCB004 against postharvest fungal pathogen Journal of Applied Microbiology Vol108 Pages 386-395
- Bintang M Purwanto UMS Kusumawati DE Yang JJ (2015) Study of endophytic bacteria as a novel source of antioxidant agent based on GC-MS analysis International Journal of chemical, environmental and biological science Vol 3No5 Pages 368-369.

- Boora F Chirisa E Mukanganyama S (2014) Evaluation of Nitrite Radical Scavenging properties of selected Zimbabwean plant extracts and their phytoconstituents. Journal of Food Processing Vol 2014 Pages 1-8
- Essien GE Nwidi LL Nwafor PA (2009) Anti-Inflammatory and Analgesic Potential of Methanolic Extract of *Emilia sonchifolia* (Compositae) Leaves in Rodents African Journal of Biomedical Research Vol12 No3 Pages199-207
- Gond SK Bergen MS Torres MS White JF Jr (2015) Endophytic *Bacillus* sp. produce antifungal lipopeptide and induce host defence gene expression in Maize Microbiological research Vol172 Pages 79-87
- Haddad N Liu X Yang S Mu B (2008) Surfactin isoforms from *Bacillus subtilis* HSO121: separation and characterization Protein and Peptide Letters Vol15 No 3 Pages 265-269
- Hanif A Zhang F Li P Li C Xu Y Zubair M Zhang M Jia D Zhao X Liang J Majid T Yan J Farzand A Wu H Gu Q Gao X (2019) Fengycin produced by *Bacillus amyloliquefaciens* FZB42 Inhibits *Fusarium graminearum* growth and mycotoxins biosynthesis Toxins Vol11No295 Pages1-11
- Hazarika DJ Goswami G Gautom T Parveen A Das P Barooah M Boro RC (2019) Lipopeptide mediated biocontrol activity of endophytic *Bacillus subtilis* against fungal phytopathogens BMC Microbiology Vol 19 No71 Pages 1-13
- Imran M Ullah F Ayaz M Sadiq A Shah M R Jan M S Ullah F (2017) Anticholinesterase and antioxidant potentials of *Nonea micrantha* Bioss & Reut along with GCMS analysis BMC Complementary and Alternative Medicine Vol17 No499 Pages 1-12
- Jayanthi P Lalitha P (2011) Reducing power of the solvent extracts of *Eichhornia crassipes* (mart.) Solms. International Journal of Pharmacy and Pharmaceutical Sciences Vol 3 Pages 126-128
- Kamtekar S Keer V Patil V (2014) Estimation of Phenolic content, Flavonoid content, Antioxidant and Alpha amylase Inhibitory Activity of Marketed Polyherbal Formulation Journal of Applied Pharmaceutical Science Vol 4 No 09 Pages 61-65
- Kim YT Park BK Kim SE Lee WJ Moon JS Cho MS

- Park H-Y Hwang I Kim SU (2017) Organization and characterization of genetic regions in *Bacillus subtilis* subsp. *krietiensis* ATCC55079 associated with the biosynthesis of iturin and surfactin compounds PLoS ONE Vol12 No e0188179
- Kumar DG Syafiq AM Ruhaiyem Y (2015) Traditional uses, phytochemical and pharmacological aspects of *Emilia sonchifolia* (L.)DC. International Journal of Research in Ayurveda and Pharmacy Vol 6 No 4 Pages 551-556
- Latha YL Darah I Sasidharan S Jain K (2009) Antimicrobial activity of *Emilia sonchifolia* DC., *Tridax procumbence* L., and *Vernonia cineria* L., of Asteraceae family, potential as food preservatives Malasian Journal of Nutrition Vol15 No2 Pages 223-231
- Luo C Zhou H Zou J Wang X Zhang R Xiang Y Chen Z (2015) Bacillomycin L and surfactin contribute synergistically to the phenotypic features of *Bacillus subtilis* 916 and the biocontrol of rice sheath blight induced by *Rhizoctonia solani*. Applied Microbiology and Biotechnology Vol 99 Pages1897-1910
- Malfanova N Kamilova F Validov S Shcherbakov A Chebotar V Tikhonovich I Lugtenberg B (2011) Characterization of *Bacillus subtilis* HC8, a novel plant-beneficial endophytic strain from giant hogweed Microbial Biotechnology Vol 4 No4 Pages 523-32
- Mohamad OAA Li L Ma JB Hatab S Xu L Guo JW Rasulov BA Liu YH Hedlund BP Li WJ (2018) Evaluation of the Antimicrobial Activity of Endophytic Bacterial Populations from Chinese Traditional Medicinal Plant Licorice and Characterization of the Bioactive Secondary Metabolites Produced by *Bacillus atrophaeus* against *Verticillium dahliae*. Frontiers in Microbiology Vol 9 No 924 Pages1-14
- Nongkhlaw F M W Joshi SR (2015) L-Asparaginase and antioxidant activity of endophytic bacteria associated with ethnomedicinal plants Indian Journal of Biotechnology Vol14 Pages 59-64
- Pezzuto J (1996) Taxol production in plant cell culture comes of age Nature Biotechnology Vol14 no9 Page1083
- Rafat A Philip K Muniandy S (2012) A novel source of bioactive compounds: Endophytic bacteria isolated from *Centella asiatica*. Journal of Pure and Applied Microbiology Vol6 No1Pages1-10
- Raj A G R Shailaja U Prasanna RN Ajayan S (2013) The therapeutic potential of ten sacred plants (Dashapushpa) of Kerala State of Southern India Journal of Ayurveda and Holistic Medicine Vol1No3 Pages1-16
- Roongsawang N Washio K Morikawa M (2010) Diversity of non-ribosomal peptide synthetases involved in the biosynthesis of lipopeptide biosurfactants International Journal of Molecular Sciences Vol12 No1Pages141-172.
- Sarwar A Brader G Corretto E Aleti G Abaidullah M Sessitsch A Hafeez FY (2018) Qualitative analysis of biosurfactants from *Bacillus* species exhibiting antifungal activity PLoS ONE Vol13 No 6 Pages1-15
- Shukla S Naik G Mishra SK (2015) Potential antimicrobial activity of bacterial endophytes isolated from *Flacourtia jangomas* (Lour.) Raesch, a less explored medicinal plant Journal of Microbiology Biotechnology Food Science Vol4 No6 Pages 473-477
- Sophia D Ragavendra P Arulraj C Gopalakrishnan VK (2011) Invitro antioxidant activity and HPTLC determination of n-hexane extract of *Emilia sonchifolia* (L.)DC. Vol 2 No 4 Pages179-183
- Sulistiyani Ardyati T Winarsih S (2016) Antimicrobial and antioxidant activity of endophyte bacteria associated with *Curcuma longa* Rhizome Journal of Experimental Science Vol6 No1Pages 45-51
- Sunkar S Nachiyar CV (2012) Biogenesis of antibacterial silver nanoparticles using the endophytic bacterium *Bacillus cereus* isolated from *Garcinia xanthochymus* Asian Pacific Journal of Tropical Biomedicine Vol2 No12 Pages 953-959
- Swarnalatha Y Saha B Choudary L Y (2015) Bioactive compound analysis and antioxidant activity of endophytic bacterial extract from *Adhathoda beddomei*. Asian Journal of Pharmaceutical and Clinical Research Vol 8 No1 Pages 70-72.
- Thenmozhi K Saradha M Manian S Paulsamy S [2013] In vitro antimicrobial potential of root extracts of the medicinal plant species, *Emilia sonchifolia* (Linn.)DC. Asian Journal of Pharmaceutical and Clinical Research. Vol 6 No 3 Pages 149-151
- Urumbil S K Anilkumar M [2019] Isolation, identification and characterisation of plant growth promoting bacterial endophytes from *Emilia sonchifolia*(Linn.) DC. Asian Journal of Microbiology, Biotechnology and Environmental Sciences Pages 648-655
- Wang Y Wu Y Wang Y Xu H Mei X Yu D Wang Y Li W (2017) Antioxidant properties of probiotic bacteria Nutrients Vol9 No521Pages1-15
- Zayed MZ Ahmad FB Ho WS Pang SH (2014) GCMS analysis of phytochemical constituents in leaf extracts of *Neolamarckia cadamba* (Rubiaceae) from Malaysia. International Journal of Pharmacy and pharmaceutical Science Vol6 No9 Pages123-127
- Zheng LP Zou T Ma YJ Wang JW Zhang YQ (2016) Antioxidant and DNA Damage protecting activity of exopolysaccharides from the endophytic bacterium *Bacillus Cereus* SZ1 Molecules Vol 21 No174 Pages 1-15